



Revista Ceres

ISSN: 0034-737X

ceresonline@ufv.br

Universidade Federal de Viçosa
Brasil

Vrba Brandão, Juliana Andréia; Ribeiro Casimiro Lopes-Assad, Maria Leonor; Ceccato-Antonini,
Sandra Regina

Solubilization of diabase and phonolite dust by filamentous fungus

Revista Ceres, vol. 61, núm. 5, septiembre-octubre, 2014, pp. 740-745

Universidade Federal de Viçosa

Vicosa, Brasil

Available in: <http://www.redalyc.org/articulo.oa?id=305232579018>

- How to cite
- Complete issue
- More information about this article
- Journal's homepage in redalyc.org

redalyc.org

Scientific Information System

Network of Scientific Journals from Latin America, the Caribbean, Spain and Portugal

Non-profit academic project, developed under the open access initiative

Solubilization of diabase and phonolite dust by filamentous fungus¹

Juliana Andréia Vrba Brandão², Maria Leonor Ribeiro Casimiro Lopes-Assad³,
Sandra Regina Ceccato-Antonini⁴

<http://dx.doi.org/10.1590/0034-737X201461050018>

ABSTRACT

The objective of this study was to evaluate the effect of the fungus *Aspergillus niger* strain CCT4355 in the release of nutrients contained in two types of rock powder (diabase and phonolite) by means of *in vitro* solubilization trials. The experimental design was completely randomized in a 5 x 4 factorial design with three replications. It was evaluated five treatments (phonolite dust + culture medium; phonolite dust + fungus + culture medium; diabase powder + culture medium; diabase powder + fungus + culture medium and fungus + culture medium) and four sampling dates (0, 10, 20 and 30 days). Rock dust (0.4% w/v) was added to 125 mL Erlenmeyer flasks containing 50 mL of liquid culture medium adapted to *A. niger*. The flasks were incubated at 30°C for 30 days, and analysis of pH (in water), titratable acidity, and concentrations of soluble potassium, calcium, magnesium, zinc, iron and manganese were made. The fungus *A. niger* was able to produce organic acids that solubilized ions. This result indicates its potential to alter minerals contained in rock dust, with the ability to interact in different ways with the nutrients. A significant increase in the amount of K was found in the treatment with phonolite dust in the presence of the fungus. The strain CCT4355 of *A. niger* can solubilize minerals contained in these rocks dust.

Key words: *Aspergillus niger*, bio-solubilization, rock powder, stonemeal.

RESUMO

Solubilização de pós de diabásio e de fonolito por fungo filamentososo

Este trabalho objetivou avaliar o efeito do fungo *Aspergillus niger*, linhagem CCT4355, na liberação de nutrientes contidos em pós de dois tipos de rocha (fonolito e diabásio), por meio de ensaios de solubilização *in vitro*. O delineamento experimental foi o inteiramente casualizado, em esquema fatorial 5x4, com três repetições. Foram avaliados cinco tratamentos (pó de fonolito + meio; pó de fonolito + fungo + meio, pó de diabásio + meio, pó de diabásio + fungo + meio e fungo + meio) e quatro épocas de amostragens (0, 10, 20 e 30 dias). Adicionou-se pó de rocha (0,4% m/v) a Erlenmeyers de 125 mL, com 50 mL de meio de cultura líquido adaptado para *A. niger*. Os frascos foram incubados a 30 °C, por 30 dias, e foram feitas análises de pH em água, acidez titulável e teores de potássio, cálcio, magnésio, zinco, ferro e manganês solúveis. O *A. niger* foi capaz de produzir ácidos orgânicos que solubilizaram íons. Este resultado indica o potencial de *A. niger* para alterar minerais contidos em pós de rocha, com capacidade de interagir de diferentes formas com os nutrientes. Houve aumento significativo da quantidade de K solúvel, no tratamento com pó de fonolito, em presença do fungo. Conclui-se que a linhagem CCT4355 de *A. niger* pode solubilizar minerais contidos nesses pós de rochas.

Palavras-chave: biossolubilização, *Aspergillus niger*, pó de rocha, rochagem.

Received: 3/18/2013; Approved: 7/18/2014.

¹ Part of the Master's thesis submitted by the first author. Financial support: FINEP and CNPq.

² Agronomist, Master of Science. Estrada Outeiro Santos 1867, casa 11, Taquara, 22.713-168, Rio de Janeiro, Rio de Janeiro, Brazil. vrba@uol.com.br

³ Agronomist, Doctor of Science. Departamento de Recursos Naturais e Proteção Ambiental, Universidade Federal de São Carlos, Campus Araras, Via Anhanguera, km 174, Caixa Postal 153, 13600-970, Araras, São Paulo, Brazil. assad@cca.ufscar.br (corresponding author).

⁴ Biologist, Doctor of Science. Departamento de Tecnologia Agroindustrial e Socioeconomia Rural, Universidade Federal de São Carlos, Via Anhanguera, Km 174, Caixa Postal 153, 13600-970, Araras, São Paulo, Brazil. antonini@cca.ufscar.br

INTRODUCTION

Brazil has tropical soils formed from different origin matter, often highly weathered, with accelerated removal of nutrients, which eventually promote high acidity and aluminum toxicity, besides the high capacity of phosphorus fixation (Lopes & Guilherme 2007). However, their low natural fertility can be modified by means of appropriate management techniques for conservation of their physical, chemical and biological characteristics. Noteworthy are the reduction of the intensity of tillage, crop rotation, intercropping cultivation, planting on contour lines, use of mulch, green manure and fertilizer to supply the nutrients demanded for the proper development of plants.

According to the Associação Nacional para Difusão de Adubos – ANDA (National Association for the Promotion of Fertilizers) (2014), Brazil imported more than 21.5 million tons of products to manufacture fertilizers in 2013. Because such fertilizer uses fossil fuel for its processing and transportation, its market value is directly influenced by currency fluctuations and fluctuations in the price of a barrel of oil. Moreover, because they are highly soluble, these fertilizers can be easily leached, both in rainfed agricultural systems as in irrigated systems. Furthermore, soluble fertilizers are not permitted under the current technical standards for organic agriculture, since soil biological activity and water quality must be maintained and incremented in this type of agriculture (Brasil, 1999; FAO, 2007).

Studies have been made in Brazil to alternative sources of nutrients, using organic matter or minerals that have not been commercially exploited due to the lack of knowledge on the potential use, lack of product positioning in the market or lack of interest by the fertilizer industry (Benites *et al.*, 2010). Among these studies, the use of *in natura* rock powder directly in the soil has been considered of strategic importance for the country since it contributes to the reduction of negative impacts on the Brazilian trade balance, caused by the import of fertilizers (Martins *et al.*, 2008; Nascimento & Loureiro, 2009).

However, the processes of chemical decomposition of minerals and the release of nutrients from the rock dust to the soil solution, in appropriate way form and amount for plant absorption, are relatively slow. In order to solve this issue, rock dust with finer grain size is searched because the stability of the minerals that compose rocks depends on its structure, its solubility, and its specific surface area among other factors (Kämpf *et al.*, 2009). Promising results have been obtained in experiments with different micro-organisms (filamentous fungi, bacteria and yeasts) capable of

promoting the solubilization of rock dust, accelerating chemical modification processes of silicate contained in the dust (Sheng, 2005; Sugumaran & Janarthnam, 2007; Hamdali *et al.*, 2008; Lian *et al.*, 2008; Xiao *et al.*, 2008).

Saprophytic filamentous fungi present, as part of their metabolic activity, a potent secretory system of acids and enzymes able to solubilize materials and make nutrients available (Grimm *et al.*, 2005; Guebel & Darías, 2001). Among the ascomycetes filamentous fungi, *Aspergillus niger* has been extensively used in studies of rock powder solubilization (Lopes-Assad *et al.*, 2006; Lopes-Assad *et al.*, 2010). The biosolubilization promoted by *Aspergillus niger* occurs in an indirect manner, by the action of reducing acids (citric and oxalic acids), which are produced by the metabolism of the fungus (Mehta *et al.*, 2010) and released into the solution. It is found in leaf litter in the nature and its growth occurs in wide ranges of temperature (0 to 47°C) (Schuster *et al.*, 2002).

The objective of this study was to evaluate the effect of the fungus *Aspergillus niger*, strain CCT4355, on the release of nutrients contained in phonolite and diabase dust by means of *in vitro* trials.

MATERIAL AND METHODS

The experiment was carried out in the Laboratory of Agricultural and Molecular Microbiology (LAMAM) and nutrient analysis was performed at the Soil and Plant Chemical Analysis Laboratory, at the Center for Agriculture Sciences of Universidade Federal de São Carlos, Araras Campus in the State of São Paulo.

The phonolite dust was provided by a mining company from Poços de Caldas, State of Minas Gerais, and diabase dust was supplied by a quarry from Araras, State of São Paulo. Both presented median amounts of silica (SiO₂) (Table 1).

In the solubilization tests, it was used the fungus *Aspergillus niger*, strain CCT-4355, isolated from sugar cane vinasse (Ceccato, 1989), of the LAMAM microorganism bank. Until the beginning of the tests, the fungus remained stored in the refrigerator (6-8°C) in test tubes, in MYPG inclined culture medium. Regarding production of inoculum, MYPG medium (3 g L⁻¹ of meat extract, 3g L⁻¹ yeast extract, 8 g L⁻¹ peptone, 10 g L⁻¹ glucose and 20 g L⁻¹ agar) was used in culture in the test tube in inclined medium; the tubes were incubated at 30°C for 5 days. A volume of 4 mL Tween 80 (0.1%) was added to each culture tube and the spores were scraped with a platinum loop for their release. A suspension from 1.0 to 1.8 x 10⁷ spores mL⁻¹ was obtained, after counting in a Neubauer chamber. For

setting up the experiment, it was used 50 mL of the liquid culture medium adapted for *A. niger* (2.85 g L⁻¹ sodium citrate, 1 g L⁻¹ ammonium phosphate, 0.5 g L⁻¹ magnesium sulfate, 0.132 g L⁻¹ calcium chloride and 10 g L⁻¹ glucose and pH adjusted to 7.0), according to Cerezine *et al.* (1988). This means that in 50 mL of culture medium, there was 2.35 mg calcium, 38 mg sodium, 239.5 mg of carbon, 6.08 mg nitrogen, 13.48 mg phosphorus, 5 mg magnesium, 6.65 mg sulfur and 4.2 mg chlorine. The cultivation was performed in 125-mL Erlenmeyer flasks, with addition or not of 0.2 g of rock dust (0.4% w/v), with particle size smaller than 0.053 mm (270 mesh) and inoculation or not of 0.5 mL of the spore suspension, according to the treatment.

The *in vitro* experiment was carried out in a completely randomized design in a 5 x 4 factorial design with three replications. The treatments were as follows: *A. niger* + medium (MA); phonolite dust + medium (FM); phonolite dust + *A. niger* + medium (FMA); diabase dust + medium (DM) and diabase dust + *A. niger* + medium (DMA). The flasks were incubated at 30°C, 160 rpm, for 30 days, with four sampling times (0, 10, 20 and 30 days). The contents of the flasks were filtered under vacuum, using no. 1 Whatman filter paper. The following were analyzed in the filtrate: water pH and titratable acidity by samples titration up to pH 7, using NaOH 0.05 mol L⁻¹ were analysed in the filtrate. Contents of potassium (K) were determined by flame emission photometry and contents of soluble calcium (Ca), zinc (Zn), manganese (Mn) and iron (Fe) by atomic absorption spectrophotometry.

The results of *in vitro* tests were submitted to analysis of variance (F test) and means were compared by Tukey test at 5% probability or by regression depending on the origin of the data, quantitative or qualitative, using the statistical software Statistica 6.0 (2001).

RESULTS AND DISCUSSION

The phonolite rock dust in the presence of the fungus provided greater amount of solubilized K (Table 2); however, the action of *A. niger* on the diabase rock powder did not provide significant difference in the amount of solubilized K. The amount of K from the phonolite rock dust is ten times greater than that of diabase rock dust (Table 1), which explains the result

obtained in phonolite + medium + fungus treatment (AMF), which had a high release of the soluble K into the medium, demonstrating the solubilizing potential of the fungus.

Potassium present in feldspars is not readily available because it is strongly bound to oxygen molecules of tetrahedrons of SiO₄ and AlO₄ (Curi *et al.*, 2005). Therefore, the release of K⁺ ions requires the dissolution of feldspars, by the action of H⁺ ions from the acids produced by *A. niger*. The reaction can be expressed in a simplified way, as $\text{KAlSi}_3\text{O}_8 + \text{H}^+ \rightarrow \text{HAlSi}_3\text{O}_8 + \text{K}^+$.

Contents of Ca were higher in treatments with rock powder, without the presence of fungus, with no difference between the two types of dust (Table 2), indicating a likely intake by the fungus. This is because Ca silicates and aluminosilicates are sufficiently soluble in low pH, but the concentration of soluble Ca decreases as pH increases (Mello & Perez, 2009). The small amounts of Ca found in the treatment with phonolite rock dust, which showed no calcium oxide in its composition (Table 1), can be explained by the addition of calcium chloride in the culture medium.

The fungus *A. niger* had no effect on the solubilization of Mg, Zn, Fe and Mn in the treatments with both studied rock dust (Table 2), and the amount of solubilized Fe was higher in diabase rock dust treatment. Phonolite is an alkaline leucocratic rock whereas the evaluated diabase (geologically classified as a lati-andesite) is a basic mesocratic rock (Machado *et al.* 2005), a reflex of the mafic mineral contents that have dark color since they contain iron, magnesium, titanium, manganese, and so on. The iron content of the diabase rock was higher than that of the phonolite rock (Table 1). Therefore, the amount of that ion in the medium was higher in the diabase + medium + *A. niger* (DMA) treatment than in the phonolite + medium + *A. niger* (FMA) treatment.

Significant results at 5% of probability were found in the time and treatment interaction, for all attributes, except for pH and titratable acidity, which showed high variance. Data regarding treatments with diabase + *A. niger* + medium (DMA) and phonolite dust + *A. niger* + medium (FMA) are discussed in this paper.

The levels of soluble Ca (Figure 1A) decreased until the twentieth day of the trial; after this period, the concentration of Ca increased, but this increase was not

Table 1. Chemical characterization of phonolite and diabase dust used in *in vitro* solubilization trial

Rock dust	SiO ₂	Fe ₂ O ₃	K ₂ O	Al ₂ O ₃	Na ₂ O	MgO	CaO	TiO ₂	MnO
	%								
Phonolite ¹	54.8	3.29	8.46	21.3	6.07	-	-	-	-
Diabase ²	50.31	12.62	0.84	13.94	2.24	6.91	10.16	1.02	0.19

¹ Supplied by minning company Curimbaba Ltda; ² Supplied by Acme Analytical Laboratories LTD.

due to the action of *A. niger* (Table 2). This behavior is in part due to the consumption of this element by the microorganism, in part due to the presence of organic acids produced by *A. niger*, especially citric acid ($C_6H_8O_7$), which presents a chelating activity by Ca^{+2} (Carvalho *et al.*, 2005).

The content of potassium showed an increase, over time, not reaching its maximum value (Figure 1B) being much more expressive in phonolite dust, richer in K_2O than diabase dust (Table 1). After 30 days, it was found that the amount of K, solubilized by *A. niger*, in the treatment with phonolite dust + medium + fungus (FMA) has not reached its maximum degree (Figure 1B). Because each flask with phonolite dust contained $7.2 \text{ mmol}_c \text{ dm}^{-3}$ K, the content of soluble K, after 30 days of incubation, was $1.21 \text{ mmol}_c \text{ dm}^{-3}$ in the phonolite + medium + fungus treatment. This corresponds to a solubilization rate of 16.8% whereas this rate remained at 7.2% in the medium + phonolite treatment. Using the same strain of *A. niger*, Lopes-Assad *et al.* (2006) obtained solubilization rates of 54.7% for the alkaline ultramafic rock dust, containing 3.32% of K_2O and 33.26% SiO_2 , and 11.8% for phlogopitite dust, containing 5.13% of K_2O , and 47.12% SiO_2 at 21 days of incubation. These results suggest that the rate of solubilization obtained with *A. niger* is probably related to the amounts of SiO_2 and K_2O , present in rocks and their mineralogical characteristics.

The behavior of ion Mg (Figure 2A) in the treatments with diabase dust was similar to the behavior presented by Ca, with a decrease in the content until day 20, probably due to consumption by the microorganism, and a subsequent increase in which it was not possible to state that the maximum solubilization was reached at the end of the trial. The behavior of the ions Zn, Fe and Mn in the treatments with diabase rock dust was similar over the 30 days of the trial (Figures 2b, c and d), with greater release between the tenth and twentieth day of the trial.

Fungus *A. niger* releases reducing acids (citric and oxalic acids), which convert the insoluble forms Fe (III) and MnO_2 into soluble forms Fe (II) and Mn (II) (Mehta *et al.*, 2010). Some factors have an influence in the submerged fermentation of citric acid (type and




Figure 1. Variation in the concentration of calcium (A) and potassium (B) over 30 days of solubilization trials of diabase dust (DMA) and phonolite dust (FMA) using *Aspergillus niger* in liquid culture medium.

Table 2. Chemical characteristics determined at 30 days of *in vitro* trials of solubilization of phonolite and diabase dust by the fungus *Aspergillus niger* strain CCT4355

Characteristics	Treatments				
	DM	DMA	FM	FMA	MA
pH	6.64 b	7.15 a	6.71 b	5.95 c	3.53 d
Titrateable acidity ($\text{mmol}_c \text{ dm}^{-3}$)	2.76 bc	1.00 c	2.00 c	8.16 b	52.00 a
Potassium ($\text{mmol}_c \text{ dm}^{-3}$)	0.36 c	0.26 c	0.50 b	0.77 a	0.30 c
Calcium ($\text{mmol}_c \text{ dm}^{-3}$)	0.29 a	0.08 b	0.26 a	0.08 b	0.06 b
Magnesium ($\text{mmol}_c \text{ dm}^{-3}$)	0.39 a	0.33 a	0.38 a	0.35 a	0.38 a
Zinc (mg dm^{-3})	0.19 a	0.21 a	0.13 bc	0.16 ab	0.09 c
Iron (mg dm^{-3})	64.40 a	64.47 a	12.25 bc	16.77 b	9.37 c
Manganese (mg dm^{-3})	287 a	2.21 ab	2.58 ab	2.41 b	0.12 c

DM = diabase dust in liquid culture medium; DMA = diabase dust and *A. niger* in liquid culture medium; FM = phonolite dust in liquid culture medium; FMA = phonolite dust and *A. niger* in liquid culture medium; MA = *A. niger* in liquid culture medium. Means on the lines, followed by an equal letter at least, do not differ by Tukey test at 5% probability




Figure 2. Variation in the contents of magnesium (A), zinc (B), iron (C) and manganese (D) over 30 days of solubilization trials of diabase (DMA) dust and phonolite dust (FMA) using *Aspergillus niger* in liquid culture medium.

concentration of the C source, N and phosphate limitation, pH, aeration, concentration of trace elements and morphology of the producer organism). Among the metals that need to be in a limit amount are Zn, Mn, Fe, Cu, alkali metals and heavy metals (Papagianni, 2007). In the solubilization of silicates contained in rock dust, the concentrations of Fe and Mn in the medium decrease at first, possibly due to the formation of oxides and hydroxides of Fe and Mn with low solubility (Oliveira Junior *et al.*, 2000; Mello & Perez, 2009).

Plant ashes are two sources of potassium commonly used in organic agriculture, with about 10% of K, and the decomposed cattle manure, with about 5% of K (Souza & Alcântara, 2008). Thus, phonolite dust, with about 8% of K, may be an alternative to organic fertilization. The biotechnology pathway by incubation of rock dust with *A. niger* and production of biofertilizer is a promising alternative in order to accelerate the release of nutrients.

CONCLUSIONS

The fungus *Aspergillus niger* promoted the solubilization of the elements contained in the dust of diabase and phonolite rocks.

The dust of the evaluated phonolite released soluble potassium in the culture medium in a significant way after 30 days of incubation.

Over the 30 days of trial, there were no significant variations in pH and exchangeable acidity in the performed treatments; however, there was effect on the levels of the soluble ions studied.

ACKNOWLEDGMENTS

To the CNPq, for the scholarship granted to the first two authors and to FINEP for the financial support by the ROCKBIOCOM – Ref. Finep 0944/08 project.

REFERENCES

- ANDA - Associação Nacional para Difusão de Adubos (2014) Principais indicadores do setor de fertilizantes. Available at: <<http://www.anda.org.br/estatisticas.aspx>>. Accessed in May 20th, 2014.
- Benites VM, Polidoro JC & Resende AV (2010) Oportunidades para a inovação tecnológica no setor de fertilizantes no Brasil. Boletim Informativo da Sociedade Brasileira de Ciência do Solo, 35:18-21.
- Brasil (1999) Ministério da Agricultura, Pecuária e Abastecimento. Instrução Normativa nº 7, de 17 de maio de 1999. Diário Oficial [da] República Federativa do Brasil. Brasília, DF, 19 de maio de 1999, Seção 1. 11p.
- Carvalho W, Silva DDV, Canilha L & Mancilha IM (2005) Aditivos alimentares produzidos por via fermentativa parte I: ácidos orgânicos. Revista Analytica, 18:70-76.
- Ceccato SR (1989) Produção de biomassa e depuração biológica da vinhaça de cana de açúcar por cultura mista de fungos filamentosos e leveduras. Dissertação de Mestrado. Universidade Estadual Paulista, Rio Claro. 159p.
- Cerezine PC, Nahas E & Banzatto DA (1988) Phosphate accumulation by *Aspergillus niger* from fluorapatite. Applied Microbiology and Biotechnology, 29:501-505.
- Curi N, Kämpf N & Marques JJ (2005) Mineralogia e formas de potássio em solos brasileiros. In: Yamada T & Roberts TL (Eds.) Potássio na agricultura brasileira. Piracicaba, POTAFOS. p.71-91.
- FAO World Health Organization (2007) Guidelines for the production, processing, labelling and marketing of organically produced foods. Rome, FAO-WHO. 39p. (Codex Alimentarius. CAC/GL, 32).
- Grimm LH, Kelly S, Krull R & Hempel DC (2005) Morphology and productivity of filamentous fungi. Applied Microbiology and Biotechnology, 69:375-384.
- Guebel DV & Darias NVT (2001) Optimisation of citric acid production by *A. niger* through a metabolite flux balance model. Process Biotechnology, 4:01-11.
- Hamdali H, Hafidi M, Virolle MJ & Ouhdouch Y (2008) Rock phosphate-solubilizing actinomycetes: screening for plant growth-promoting activities. World Journal of Microbiology and Biotechnology, 24:2565-2575.
- Kämpf N, Curi N & Marques JJ (2009) Intemperismo e ocorrência de minerais no ambiente do solo. In: Melo, VF & Alleoni LR (Eds.) Química e mineralogia do solo. Viçosa, SBCS. p.333-379.
- Lian B, Wang B, Pan M, Liu C & Teng HH (2008) Microbial release of potassium from K-bearing minerals by thermophilic fungus *Aspergillus fumigatus*. Geochimica et Cosmochimica Acta, 72:87-98.
- Lopes AS & Guilherme LRG (2007) Fertilidade do Solo e Produtividade Agrícola. In: Novais RF, Alvarez V VH, Barros NF, Fontes RLF, Cantarutti RB, Neves JCL (Eds.) Fertilidade do Solo. Viçosa, SBCS. p.1-64.
- Lopes-Assad ML, Rosa MM, Erler G & Ceccato-Antonini SR (2006) Solubilização de pó-de-rocha por *Aspergillus niger*. Espaço & Geografia, 9:1-17.
- Lopes-Assad ML, Avansini SH, Rosa MM, Carvalho JRP & Ceccato-Antonini SR (2010) The solubilization of potassium-bearing rock powder by *Aspergillus niger* in small-scale batch fermentations. Canadian Journal of Microbiology, 56:598-605.
- Machado FB, Nardy AJR, Melo RP, Oliveira MAF & Squisato E (2005) As rochas intrusivas da formação Serra Geral na porção leste da bacia do Paraná no estado de São Paulo: aspectos petrográficos e geoquímicos – resultados preliminares. Geociências, 24:5-17.
- Martins ES, Oliveira CG, Resende AV & Matos MSF (2008) Agrominerais – Rochas Silicáticas como Fontes Mineraias de Potássio para a Agricultura. In: Luz AB & Lins FAF (Eds.) Rochas e Minerais Industriais: Usos e Especificações. Rio de Janeiro, CETEM. p.205-223.
- Mehta KD, Chitrangada DAS & Pandey BD (2010) Leaching of copper, nickel and cobalt from Indian Ocean manganese nodules by *Aspergillus niger*. Hydrometallurgy, 105:89-95.
- Mello JWV de & Perez DV (2009) Equilíbrio Químico das Reações no Solo. In: Melo V de F, Alleoni LR (Eds.) Química e Mineralogia do Solo. Viçosa, SBCS. p.151-249.
- Nascimento M & Loureiro FEL (2009) O potássio na agricultura brasileira: fontes e rotas alternativas. In: Lapido-Loureiro FE, Melamed R, Neto JF (Eds.) Fertilizantes: agroindústria e sustentabilidade. Rio de Janeiro, CETEM. p.305-335.
- Oliveira Junior JA de, Malavolta E & Cabral CP (2000) Efeitos do manganês sobre a soja cultivada em solo de Cerrado do Triângulo Mineiro. Pesquisa Agropecuária Brasileira, 35:1629-1636.
- Papagianni M (2007) Advances in citric acid fermentation by *Aspergillus niger*: biochemical aspects, membrane transport and modeling. Biotechnology Advances, 25:244-263.
- Schuster E, Dunn-Coleman N, Frisvad JC & Van Dijck PWM (2002) On the safety of *Aspergillus niger* – a review. Applied Microbiology and Biotechnology, 59:426-435.
- Sheng XF (2005) Growth promotion and increased potassium uptake of cotton and rape by a potassium-releasing strain of *Bacillus edaphicus*. Soil Biology and Biochemistry, 37:1918-1922.
- Souza RB de & Alcântara FA (2008) Adubação no sistema orgânico de produção de hortaliças. Brasília, Embrapa Hortaliças. 8p. (Circular técnica, 65).
- Sugumaran P & Janarthanam B (2007) Solubilization of potassium-containing minerals by bacteria and their effect on plant growth. World Journal of Agricultural Sciences, 3:350-355.
- Xiao CQ, Chi RA, Huang XH, Zhang WX, Qiu GZ & Wang DZ (2008) Optimization for rock phosphate solubilization by phosphate-solubilizing fungi isolated from phosphate mines. Ecological Engineering, 33:187-193.