



Revista Ceres

ISSN: 0034-737X

ceresonline@ufv.br

Universidade Federal de Viçosa  
Brasil

Avelar Dornelas, Gabriel; Ampélio Pozza, Edson; de Souza, Paulo Estevão; Vêras da Costa, Rodrigo; Alexandre Pozza, Adélia Aziz; Alvarenga Santos, Leandro  
Nitrogen and potassium fertilization on the yield and intensity of the maize white spot  
Revista Ceres, vol. 62, núm. 4, julio-agosto, 2015, pp. 351-359  
Universidade Federal de Viçosa  
Viçosa, Brasil

Available in: <http://www.redalyc.org/articulo.oa?id=305241508004>

- How to cite
- Complete issue
- More information about this article
- Journal's homepage in redalyc.org

redalyc.org

Scientific Information System

Network of Scientific Journals from Latin America, the Caribbean, Spain and Portugal

Non-profit academic project, developed under the open access initiative

# Nitrogen and potassium fertilization on the yield and intensity of the maize white spot<sup>1</sup>

Gabriel Avelar Dornelas<sup>\*2</sup>, Edson Ampélio Pozza<sup>3</sup>, Paulo Estevão de Souza<sup>3</sup>, Rodrigo Vêras da Costa<sup>4</sup>, Adélia Aziz Alexandre Pozza<sup>5</sup>, Leandro Alvarenga Santos<sup>6</sup>

<http://dx.doi.org/10.1590/0034-737X201562040004>

## ABSTRACT

A plant's nutritional balance can influence its resistance to diseases. In order to evaluate the effect of increasing doses of N and K on the yield and severity of the maize white spot, two experiments were installed in the field, one in the city of Ijaci, Minas Gerais, and the other in the city of Sete Lagoas, Minas Gerais. The experimental delimitation was in randomized blocks with 5 x 5 factorial analysis of variance, and four repetitions. The treatments consisted of five doses of N (20; 40; 80; 150; 190 Kg ha<sup>-1</sup> of N in the experiments 1 and 2) and five doses of K (15; 30; 60; 120; 180 Kg ha<sup>-1</sup> of K in experiment 1 and 8.75; 17.5; 35; 50; 100 Kg ha<sup>-1</sup> of K in experiment 2). The susceptible cultivar 30P70 was planted in both experiments. The plot consisted of four rows 5 meters long, with a useful area consisting of two central rows 3 meters each. Evaluations began 43 days after emergence (DAE) in the first experiment and 56 DAE in the second one. There was no significant interaction between doses of N and K and the disease progress. The effect was only observed for N. The K did not influence the yield and the severity of the disease in these experiments. Bigger areas below the severity progress curve of the white spot and better yield were observed with increasing doses of N. Thus, with increasing doses of N, the white spot increased and also did the yield.

**Keywords:** Nutritional stress, *Zea mays* L., foliar diseases.

## RESUMO

### Efeito da adubação nitrogenada e potássica na produtividade e na intensidade da mancha branca do milho

O equilíbrio nutricional de plantas pode influenciar a resistência a doenças. Com o objetivo de avaliar o efeito de doses crescentes de N e de K na produtividade e na severidade da mancha branca do milho, foram instalados em campo, dois experimentos, um em Ijaci, MG e o outro em Sete Lagoas, MG. O delineamento experimental foi em blocos casualizados com esquema de análise de variância fatorial 5 x 5 e quatro repetições. Os tratamentos consistiram de 5 doses de N (20; 40; 80; 150; 190 Kg ha<sup>-1</sup> de N nos experimentos 1 e 2) e de 5 doses de K (15; 30; 60; 120; 180 Kg ha<sup>-1</sup> de K no experimento 1 e 8,75; 17,5; 35; 50; 100 Kg ha<sup>-1</sup> de K no experimento 2). Foi plantado nos dois experimentos o cultivar suscetível 30P70. As parcelas foram constituídas de quatro fileiras de 5 m de comprimento, sendo a área útil composta por duas linhas centrais com 3 m cada. As avaliações iniciaram-se aos 43 dias após emergência (DAE) no primeiro e aos 56 DAE, no segundo experimento. Não houve interação significativa entre as doses de N e de K e o

Submitted on 04/23/2014 and approved on 07/08/2015.

<sup>1</sup> This paper is part of the first author's master dissertation.

<sup>2</sup> Universidade Federal de Lavras, Departamento de Fitopatologia, Lavras, Minas Gerais, Brazil. gabrieldornelas@yahoo.com.br

<sup>3</sup> Universidade Federal de Lavras, Departamento de Fitopatologia, Lavras, Minas Gerais, Brazil. eapozza@dfp.ufla.br; pauleste@dfp.ufla.br

<sup>4</sup> Embrapa Milho e Sorgo, Gurupi, Tocantins, Brazil. rodrigo.veras@embrapa.br

<sup>5</sup> Universidade Federal de Lavras, Departamento de Ciência do Solo, Minas Gerais, Brazil. adelia.pozza@des.ufla.br

<sup>6</sup> Universidade Estadual do Centro Oeste, Departamento de Agronomia, Guarapuava, Paraná, Brazil. leandro.alvarenga.s@hotmail.com

\* Corresponding author: gabrieldornelas@yahoo.com.br

progresso da doença. O efeito foi observado apenas para o N. O K não influenciou a produtividade e a severidade da doença nesses experimentos. Observaram-se maiores áreas abaixo da curva de progresso da severidade da mancha branca e maior produtividade com o aumento das doses de N. Sendo assim, com o aumento das doses de N houve aumento da mancha branca do milho e também da produtividade.

**Palavras-chave:** Estresse nutricional, *Zea mays* L., doenças foliares.

## INTRODUCTION

The white spot - *Pantoea ananatis* (Paccola-Meirelles *et al.*, 2001; Bomfeti *et al.*, 2008), is one of the most important diseases in corn crop. Historically, it only occurred at the end of the cycle of the crop; however, high severity was observed in young plants, during the vegetative phase, resulting in premature wilt of plants in susceptible genotypes (Mendes & Tebaldi, 2011). The disease occurs in temperature and relative humidity higher than or equal to 14° C and 60%, respectively (Rolim *et al.*, 2007) in regions above 700 m.a.s.l. (altitude) (Silva & Menten, 1997), due to the longer duration of leaf wetness period, provided by dew. In these conditions, damages can cause losses of up to 60% of corn yield (Rolim *et al.*, 2007).

Among environmental factors that influence the occurrence of the disease, some studies have highlighted the important role of temperature and water (Fernandes *et al.*, 1995; Fantin *et al.*, 2005). However, few studies have described the interaction of mineral nutrition with this disease. The nutritional status of plants is an environmental factor that can be manipulated with relative ease, via fertilization. The supply of nutrients in a balanced way assists in building of several horizontal resistance barriers, including the wax layer (Pozza *et al.*, 2004) and the cell wall, responsible for slowing the penetration and colonization of the pathogen, reducing the epidemic progress rate (Pozza & Pozza, 2012).

In the literature, there are several reports about how the effect of macro and micro nutrients promote reduction of the intensity of diseases in different crops, such as cercosporiosis (Pozza *et al.*, 2001; 2004; Garcia Júnior *et al.*, 2003) and the stain of Phoma in coffee trees (Lima *et al.*, 2010), anthracnose in strawberries (Tanaka *et al.*, 2002), Phoma in Brassicas (Sochting & Verret, 2004) and the perforations in plum leaves (Tutida *et al.*, 2007) among others.

When it comes to corns, nutritional imbalances between N and K predisposed plants of two cultivars to

infection by *Colletotrichum graminicola* (Ces.) Wils. (SensuArx, 1957). In both cultivars, the amount of damaged leaf area depended on the interaction between the nutrients. The lowest severity levels were observed at the lowest dose of N, combined to the highest dose of K, and nitrogen fertilization influenced the K content negatively in the upper canopy. Caldwell *et al.* (2002) also report an increase in the severity of cercosporiosis (*Cercospora zea maydis* Tehon & Daniels) of corn with increased doses of N (0, 60 and 120 Kg ha<sup>-1</sup>) and K (0, 25, 50 and 150 Kg ha<sup>-1</sup>) in the soil. However, there is still no information about the influence of N and K have upon the maize white spot.

Therefore, this study was performed in order to assess the effect of nitrogen and potassium fertilization on the yield and intensity of the white spot infection of corn.

## MATERIAL AND METHODS

The experiments were installed in two field areas. The first one was implemented on 11/25/2011, in an area located on *Palmital* farm of the Universidade Federal de Lavras (UFLA), in the city of Ijaci, Minas Gerais, located at 21°09'50"S, 44°55'09"W and 837 m.a.s.l. (altitude). The second experiment was implemented on 11/18/2011, in *EMBRAPA Milho e Sorgo's* experimental area, in the city of Sete Lagoas, Minas Gerais, 19°26'00"S, 44°10'59"W and 732 m.a.s.l. (altitude). In both experiments, the Pioneer® 30P70 corn was cultivated, a simple hybrid likely to develop the white spot. The experiments were conducted under natural infection. The soil preparation was done by minimum cultivation with scarification and harrowing on the sowing eve.

The experimental design was installed in randomized blocks, in 5 x 5 factorial design, and four repetitions. The treatments corresponded to five doses of N and five doses of K combined with each other, in each area. The N doses were 20, 40, 80, 150 and 190 Kg ha<sup>-1</sup>. The doses of K were calculated individually in each experimental area, according to the results of soil analysis (Table 1) and

following the recommendations for the crop in Minas Gerais (Ribeiro *et al.*, 1999). The doses of K were 15, 30, 60, 120 and 180 Kg ha<sup>-1</sup> of K<sub>2</sub>O for experiment 1 and 8.75, 17.5, 35, 50 and 100 Kg ha<sup>-1</sup> of K<sub>2</sub>O for experiment 2. The parts consisted of four rows of 5m long and the useful area by two central rows of 3 m each. The sources of N and K were urea and potassium chloride, respectively. Twenty-five percent and 50% of doses of N and K respectively were applied at sowing, and the rest was divided in two side dressings, one 35 days after sowing and the other 15 days after the first one.

In addition to N and K, 120 and 70 Kg ha<sup>-1</sup> of P<sub>2</sub>O<sub>5</sub> were applied in experiments 1 and 2, respectively and, 2 Kg ha<sup>-1</sup> of Zn in the form of zinc sulphate and 30 Kg ha<sup>-1</sup> of calcium sulfate (14% of S) in the form of phosphogypsum, according to the results of soil analysis and recommendations of Ribeiro *et al.*, (1999), at sowing date. Two micronutrient sprays of 2.0 L h<sup>-1</sup> containing 5% Zn, 3% Mn, 0.50% of B, 0.60% Cu, 0.06% Mo and 4% S in 200 L ha<sup>-1</sup> of water were also applied, in stages V3 and V6, (Ritchie *et al.*, 2003), that is, 30 and 45 days after planting.

Planting fertilization and zinc sulfate and phosphogypsum were applied in the furrow. Sowing was performed manually, at a spacing of 0.70 x 0.20 m. For weed control, applications of glyphosate (2.0 L ha<sup>-1</sup>) were used at pre-emergence and nicosulfuron (1.5 L ha<sup>-1</sup>) mixed with atrazine (5.0 L ha<sup>-1</sup>) as post-emergence herbicides.

The evaluation of the disease severity was initiated at 43 and 56 days after emergence (DAE) in experiment 1 and 2, respectively, with the appearance of the first symptoms. From then on, evaluations were carried out every 10 days, totaling six and seven evaluations, in areas 1 and 2, respectively. These severity evaluations were performed with the help of diagrammatic scale (Agrocères, 1996). The data was used to calculate the area under the curve of progress of disease severity (AUCPS), according to Shanner & Finney (1977).

After the manual harvest the grains were threshed, weighted and the humidity content was determined. Grain yield data were refined to 13% of humidity content and expressed in t ha<sup>-1</sup>. Climatic variables were monitored

by minimum, maximum and average temperatures, average relative humidity and precipitation, in both places with Campbell Scientific® weather stations implemented near the experiments.

Data of the AUCPS and of grain yield, from each experiment, were submitted to analysis of variance. Then, the significant variables in the F test were submitted to the adjustment of linear regression models. Pearson's correlation analysis were performed between yield and the AUCPS of both experiments. Analysis were conducted in the SAS® program (SAS Institute Inc., 1996).

## RESULTS AND DISCUSSION

The average of the maximum, average and minimum daily temperatures, during the conduction period of the experiments were 28, 22 and 18 °C. Average relative humidity was of 76% and precipitation of 1078 mm in experiment 1 (Figure 1A). In experiment 2, the maximum, average and minimum daily temperatures were 29, 22 and 18° C, and the average relative humidity was 74% and the precipitation was 1225 mm (Figure 1B). According to Oliveira *et al.* (2004) and Reis *et al.* (2004) the weather conditions favorable to the occurrence of white spot are relative humidity above 60%, high precipitation and moderate temperatures between 20 to 25 °C.

### *Area under the curve of progress of disease severity (AUCPS)*

In both experiments there was no significant interaction ( $P \geq 0,05$ ) between N and K to the AUCPS of the white spot (Table 2). However, the increasing doses of N influenced the AUCPS.


With the increase of N doses, a linear rise of AUCPS was observed. There was an increase of 23.8 (Figure 2A) and 51.7% (Figure 2B) in disease severity from 20 to 190 Kg ha<sup>-1</sup> of N in experiments 1 and 2, respectively. The contribution of nitrogen fertilizer on plants' resistance to diseases varies, among other factors, depending on the pathogen, genotype, dose and source of the nutrient used and on the interaction between nutrients (Pozza & Pozza, 2012). Appropriate doses of

**Table 1:** Results of chemical analyses of the soils in Sete Lagoas – MG (Soil 1) and Ijaci- MG (Soil 2) prior to the establishment of the experiment

Soil	pH	P-rem mg L <sup>-1</sup>	P mg dm <sup>-3</sup>	K mg dm <sup>-3</sup>	Ca	Mg cmol <sub>c</sub> dm <sup>-3</sup>	Al	H+Al	SB cmol <sub>c</sub> dm <sup>-3</sup> %	T	V
Soil1 <sup>(1)</sup>	5.8	15.6	3.4	47	2.0	0.3	0.1	3.2	2.4	5.7	42.8
Soil2 <sup>(2)</sup>	6.8	20.1	22.0	107	4.7	0.6	0.0	2.1	5.6	7.7	72.9

<sup>(1)</sup>Soil Analysis Laboratory of the Department of Soil Science (UFLA).

<sup>(2)</sup>Soil Analysis Laboratory of the Department of Soil (UFV).



**Figure 1:** A) Total precipitation, relative humidity, maximum temperature, average temperature and minimum temperature in Ijaci (experiment 1) of 01/17/2012 to 03/14/2012 and in B) Sete Lagoas (experiment 2). Data obtained in the Bioclimatology sectors of UFLA, and EMBRAPA

N contributes to the synthesis of lignin, phytoalexins and tannins, but in excess, N reduces the production of these compounds, due to the demand of carbon in photosynthesis via cycle of Krebs. Thus, the synthesis of the secondary metabolites via shikimic acid is compromised and the turgor cell also raises, providing greater amounts of water and also the production of carbohydrates, such as glucose (Huber & Thompson, 2007; Taiz & Zieger, 2013), essential to the infection process of the pathogen. In addition, horizontal resistance barriers as the layer of wax and also the cell

wall may be compromised due to the rapid cell growth, making them thinner and less resistant to fungal penetration (Pozza & Pozza, 2012).

Similarly, Fidelis *et al.* (2003), relate the increased severity of the white spot of corn in early stages of growth with high doses of N compared to lower doses. The reaction of 23 cultivars of corn, submitted to low N (32 Kg ha<sup>-1</sup> of N at sowing) and to high N supply (32 Kg ha<sup>-1</sup> at sowing + 90 Kg ha<sup>-1</sup> of N on side dressing), was 6 and 30% of the severity of white spot, respectively, at 30 days after flowering.

**Table 2:** Summary of the analysis of variance for severity of white spot and yield of corn, according to the doses of N and K in experiment 1 (Ijaci) and in experiment 2 (Sete Lagoas)

#### Experiment 1


FV	GL	Severity			Productivity		
		QM <sub>Sev</sub>	Fc	Pr > Fc	QM <sub>Prod</sub>	Fc	Pr > Fc
Doses of N	4	1614496.9	17.60	0.00**	58.53	59.69	0.00**
Doses of K	4	82454.8	0.90	0.46 <sup>ns</sup>	1.15	1.17	0.33 <sup>ns</sup>
Doses N x K	16	68376.7	0.75	0.73 <sup>ns</sup>	2.74	2.79	0.00 <sup>ns</sup>
Block	3	129095.6	1.41	0.24 <sup>ns</sup>	0.14	0.14	0.93 <sup>ns</sup>
Error	72	91717.8	-	-	0.98	-	-
CV (%)				9.97			13.85

#### Experiment 2

FV	GL	Severity	Productivity				
		QM <sub>Sev</sub>	Fc	Pr > Fc	QM <sub>Prod</sub>	Fc	Pr > Fc
Doses of N	4	5524974.8	52.85	0.00**	64.55	22.03	0.00**
Doses of K	4	180333.8	1.73	0.15 <sup>ns</sup>	2.99	1.02	0.40 <sup>ns</sup>
Dose N x K	16	86976.5	0.83	0.64 <sup>ns</sup>	1.08	0.37	0.98 <sup>ns</sup>
Block	3	333440.7	3.19	0.03*	3.78	1.29	0.28 <sup>ns</sup>
Error	72	104537.1	-	-	2.93	-	-
CV (%)				11.04			27.45

<sup>ns</sup> non-significant

\*\* and \* significant, probability of 1 e 5%, by F test, respectively.



**Figure 2:** Area under the curve of progress of disease severity (AUCPS) of the white spot in the corn crop depending on the N doses, (A) experiment 1 (Ijaci) and (B) experiment 2 (Sete Lagoas).



However, other authors, studying the isolated effect of N on the severity of the white spot on the corn, found no significant effects of nitrogen fertilization and the intensity of the disease. Pegoraro *et al.* (2001) relate no substantial differences among six corn hybrids to the reaction of the white spot with two doses of nitrogen fertilization (42.5 and 85 Kg ha<sup>-1</sup> of N). According to the authors, these N doses were high enough to ensure the nutrition of plants not favoring high intensities of the white spot on the corn. Meanwhile, these doses were lower than the last two used in experiments of this study and, also lower than the highest dose used in the experiment of Fidelis *et al.* (2003). The results obtained by Pegoraro *et al.* (2001) also agree with Duarte *et al.* (1999) and Souza & Duarte (2002) in experiments with lower doses of N (8 and 32 Kg ha<sup>-1</sup>), in which changes in the susceptibility of cultivars to the white spot were not observed. But these authors also did not study how much the dose can increase without causing increment in the intensity of the disease, as studied in this work.

In other pathosystems, N can both reduce and increase diseases. The tropical rust (*Physopella zeae* (Mains) Cummins & Ramachar) of the corn (Tomazela *et al.*, 2006), for example, increased with the addition of 200 Kg ha<sup>-1</sup> of N, also followed by increased yield. This study reports the importance of nitrogen fertilization favoring foliar diseases of the corn, although it did not study the interaction with other nutrients and did not quantify its increment in relation to higher doses over time.

Still in relation to the corn, Carvalho *et al.* (2013), studying the interaction of doses of N (75, 150, 300, 600 and 1200 mg dm<sup>-3</sup>) and K (63, 125, 250, 500 and 1000 mg dm<sup>-3</sup>) in pots, and two cultivars, one moderately resistant and the other highly susceptible, report an interaction between N and K. According to these authors, for both cultivars, the largest leaf area injured (LAI) was obtained in the treatment with the lowest doses of N and K (75 mg dm<sup>-3</sup> de N and 63 mg dm<sup>-3</sup> of K, respectively). The increasing of the doses of N-NH<sub>4</sub><sup>+</sup> reduced the K content in the upper canopy, for both cultivars.

In this paper, the amount of K that already present in both soils (Table 1), may have contributed to supply the plants without needing to provide the doses via fertilization. This can be the cause of the lack of interaction between the two nutrients on the severity of the disease. For future researches, it is necessary to find soils with low levels of K.

### Grain Yield

For grain yield, the interaction between the doses of N and K was not significant ( $P \geq 0,05$ ) for both

experiments, but the doses of N ( $P < 0,01$ ) influenced this variable (Table 2).


In experiment 1, the quadratic effect of the N dose on grain yield was observed. The increment of nitrogen fertilization to dose 170.5 Kg ha<sup>-1</sup> (Figure 3A) increased yield. The response of the N doses compared to the lowest dose (20 Kg ha<sup>-1</sup>) was 18.8, 48.2, 73.7, 73.4% in the doses 40, 80, 150 and 190 Kg ha<sup>-1</sup>, respectively. For each Kg of N applied to the soil, an increase of 47.0, 40.2, 28.4, 21.6 Kg ha<sup>-1</sup> of grain was obtained, in the doses 40, 80, 150 and 190 Kg ha<sup>-1</sup>, respectively, compared to the lowest dose. In the treatment with the highest dose of N (190 Kg ha<sup>-1</sup>), yield was 3,677 Kg ha<sup>-1</sup> greater than the lower dose (40 Kg ha<sup>-1</sup>).

In experiment 2, a rising linear effect to the N (Figure 3B) was observed. The increase in grain yield compared to the lowest dose (20 Kg ha<sup>-1</sup>) was 0.56% per Kg of N applied. That is, there was an increase of 11.3, 34.0, 73.7, 96.4% in the doses 40, 80, 150 and 190 Kg ha<sup>-1</sup>, respectively. For each Kg of N applied, an increase of 24.6 Kg ha<sup>-1</sup> of grain was obtained. In the treatment with the highest dose of N (190 Kg ha<sup>-1</sup>) the yield was 4,200 Kg ha<sup>-1</sup> more grains than the lowest dose (20 Kg ha<sup>-1</sup>).

The use of high doses of N has contributed to the increment in the yield of grain in the majority of the researches done, being fundamental in the composition of the yield of the corn crop. An example of this was the increased yield of corn with 147 Kg ha<sup>-1</sup> of N, in direct planting, without irrigation, in the Brazilian “cerado”, obtained by Fernandes *et al.* (1999). Also Silva *et al.* (2005), in the region of Ilha Solteira, São Paulo, obtained the maximum yield of rain-fed corn, achieved with the dose of 166 Kg ha<sup>-1</sup> of N. In another experiment in the same region, the maximum efficiency of N was achieved with doses between 144 and 174 Kg ha<sup>-1</sup> and with quadratic response to doses of N (Silva *et al.*, 2006).

According to Sousa & Lobato (2004,) corn plants require around 20 Kg ha<sup>-1</sup> of N per ton of grains produced. Thus, in the present experiments, income below the expected to the largest dose was obtained, that is, in experiment 1, 21.90 Kg per ton of corn was spent, which is equivalent to a loss of 9%, and in experiment 2, 22.20 Kg per ton of corn were spent, equivalent to a 10% of loss. Nevertheless, this loss can be related to the severity of the white spot, which contributed to the reduction of yield.

Once K did not contribute to an increase in yield, Pavinato *et al.* (2008) conducted an experiment with six doses of N (0, 80, 120, 160, 200 and 240 Kg ha<sup>-1</sup>) in the form of urea and four doses of K<sub>2</sub>O (0, 40, 80 and



**Figure 3:** Yield of corn kernels (t ha<sup>-1</sup>) in relation to increasing doses of N (A) in experiment 1 (Ijaci) and (B) in experiment 2 (Sete Lagoas).

120 Kg ha<sup>-1</sup>), allocated in bi-factorial design, and also did not observe an increase in yield of corn when higher doses of K were applied. They relate that yield was not affected by the application of K and justified this by the high levels of K available in the soil. However, there was an increment with the doses of N and there was no significant interaction observed between N and K fertilizers. These results are similar to those found in experiments 1 and 2 described above, in which doses of K did not affect the yield of grain, which is only dependent of increased doses of N. The lack of interaction between the two nutrients (Table 2) can also be justified by “medium and good” rates of the levels of K in the soil (47 and 107 mg dm<sup>-3</sup>), in experiments 1 and 2 respectively (Table 1) according to the tables of interpretation of results of soil analysis presented by Ribeiro *et al.*, (1999).

Positive and significant values were observed for correlations between the severity of the disease (AUCPS) and grain yield (t ha<sup>-1</sup>), that is, with both N or K increasing (Table 3).

In the biggest AUCPS, grain yield was higher, this could be explained by the fact that N is the main nutrient for increasing yield and it is required in greater amounts by the crop, but it also encourages the severity of the pathogen. In this pathosystem, other studies showed different correlations, perhaps due to the low doses of N

used in these researches. For example, Souza & Duarte (2002) with doses of N (8 Kg ha<sup>-1</sup> and 32 Kg ha<sup>-1</sup>) found no correlation between white spot and corn yield. On the other hand, Brito *et al.* (2012), in an experiment with chemical control of white spot, with the same fertilization (36 Kg ha<sup>-1</sup> at sowing + 90 Kg ha<sup>-1</sup> of N as side dressing), found a negative correlation between the severity of white spot and yield, -0.41, -0.37, -0.46 in the cities of Lavras, Passos and Patos de Minas, respectively.

In another pathosystem, however, Vaz-de-Melo *et al.* (2010) observed the reaction of corn hybrids to *Curvularia ssp* on two levels of fertilization with N (high - 80 Kg ha<sup>-1</sup> and low - 40 Kg ha<sup>-1</sup>) and found a significant correlation between the severity of the *Curvularia* spot with yield. Nevertheless, the correlation between *Curvularia* severity and yield was weak and negative (-0.178) under high N conditions but negative and stronger under low N supply. Such results points out the effects of the disease severity on yield under limiting N conditions.

Therefore, the use of balanced mineral nutrition can help establish integrated management program allowing greater efficiency to control the white spot on the corn, in addition to strategy based on evasion and protection, in order to reduce the damage of this disease in plantations. Although N has raised the severity of the disease, yield also increased in response to the doses applied in the soil, requiring recommendations of additional management practices for better efficiency.

## CONCLUSIONS

Under the conditions that the experiments were conducted, the increase of dose of nitrogen favored the yield and the severity of the white spot on the corn.

**Table 3:** Pearson's correlation coefficients between grain yield and the area below the disease progress curve, considering both experiments (1 – Ijaci and 2 – Sete Lagoas)

Treatments	AUCPS	
	Experiment 1	Experiment 2
Yield of grains	0.42**	0.90**

\*\* Significant, probability of 1% ( $H_0: r=0$ )



## REFERENCES

- Agrocere (1996) Guia Agrocere de Sanidade. 2ª ed. São Paulo, Sementes Agrocere. 72p.
- Bomfeti CA, Souza-Paccola EA, Júnior Massola NS, Marriel IE, Meirelles WF, Casela CR & Paccola-Meirelles LD (2008) Localization of *P. ananatis* inside lesions of maize white spot disease using transmission electron microscopy and molecular techniques. *Tropical Plant Pathology*, 33:63-66.
- Brito AH, Pereira JLAR, Von Pinho RG & Balestre M (2012) Controle químico de doenças foliares e grãos ardidos em milho (*Zea mays* L.). *Revista Brasileira de Milho e Sorgo*, 11:49-69.
- Caldwell PM, Ward MJ, Miles N & Laing MD (2002) Assessment of the effects of fertilizer applications on gray leaf spot and yield in maize. *Plant Disease*, 86:859-866.
- Carvalho DO, Pozza EA, Casela CR, Costa RV, Pozza AAA & Carvalho CO (2013) Adubação nitrogenada e potássica na severidade da antracnose em dois cultivares de milho. *Revista Ceres*, 60:380-387.
- CONAB - Companhia Nacional de Abastecimento (2014) 5º Levantamento – Safra Agrícola de Grãos 2013/2014. Disponível em: <http://www.conab.br> Acessado em: 21 de fevereiro de 2014.
- Duarte JM, Souza JC & Corte JR (1999) Reação de cultivares de milho a *Phaeosphaeria maydis*. In: Reunião Latino Americana Del Maiz, Sete Lagoas. Memoriais, Embrapa/CNPMS. p. 405-412.
- Fantin GM, Duarte AP, Castro JL, Dudienas C, Pereira JOF, Junior AP, Geller C, Bragato EL, Kirnew PA & Cruz FA (2005) Severidade em cultivares de milho safrinha na região paulista do Vale do Paranaíba em 2004 e 2005. In: Seminário Nacional de Milho Safrinha. Campinas. Anais, Instituto Agrônomo. p. 309-318.
- Fernandes FT, Lima P, Sans LMA & Pinto NFJA (1995) Influência de fatores climáticos na ocorrência de *Phyllosticta* em milho. *Fitopatologia Brasileira*, 20:309-318.
- Fernandes LA, Vasconcellos CA, Furtini Neto AE, Roscoe R & Guedes GAA (1999) Preparo do solo e adubação nitrogenada na produção de grãos e matéria seca e acúmulo de nutrientes pelo milho. *Pesquisa Agropecuária Brasileira*, 34:1691-1698.
- Fidelis RR, Miranda GV, Souza LV, Coimbra RR, Melo AV & Galvão JCC (2003) Reação de cultivares de milho à mancha de *Phaeosphaeria* em estresse de nitrogênio. *Bioscience Journal*, 19:27-34.
- Garcia Júnior D, Pozza EA, Pozza AAA, Souza PE, Carvalho JG & Balieiro AC (2003) Incidência e severidade da cercosporiose do café em função do suprimento de potássio e cálcio em solução nutritiva. *Fitopatologia Brasileira*, 28:286-291.
- Huber DM & Thompson IA (2007) Nitrogen and plant disease. In: Datnoff LE, Elmer WH & Huber DM (Eds.) Mineral nutrition and plant disease. Saint Paul, APS Press. p.31-44.
- Lima LM, Pozza EA, Torres HN, Pozza AAA, Salgado M & Pfenning LH (2010) Relação nitrogênio/potássio com a mancha de Phoma e nutrição de mudas de café em solução nutritiva. *Tropical Plant Pathology*, 35:223-228.
- Mendes LS & Tebaldi ND (2011) *Pantoea ananatis*: importância, isolamento, cultivo em meio de cultura, inoculação, sobrevivência, disseminação, penetração, colonização e controle para cultura do milho (*Zea mays* L.). *Revisão Anual de Patologia de Plantas*, 19:386-399.
- Oliveira E, Fernandes FT, Casela CR, Pinto NFJA & Ferreira AS (2004) Diagnóstico e controle de doenças na cultura do milho. In: Galvão JCC & Miranda GV (Eds.) Tecnologias de produção do milho. Viçosa, UFV. p. 227-267.
- Paccola-Meirelles LD, Ferreira AS, Meirelles WF, Marriel IE & Casela CR (2001) Detection of bacterium associated with a leaf spot disease of maize in Brazil. *Journal of Phytopathology*, 149:275-279.
- Pavinato PS, Ceretta CA, Giroto E & Moreira ICL (2008) Nitrogênio e potássio em milho irrigado: análise técnica e econômica da fertilização. *Ciência Rural*, 38:358-364.
- Pegoraro DG, Vacaro E, Nuss CN, Soglio FK & Barbosa Neto JF (2001) Efeito de época de semeadura e adubação na mancha foliar de *Phaeosphaeria* em milho. *Pesquisa Agropecuária Brasileira*, 36:1037-1042.
- Pozza AAA, Alves E, Pozza EA, Carvalho JG, Montanari M, Guimarães PTG & Botelho DMS (2004) Efeito do silício no controle da cercosporiose em três variedades de café. *Fitopatologia Brasileira*, 29:185-188.
- Pozza AAA, Martinez HEP, Caixeta SL, Cardoso AA, Zambolim L & Pozza EA (2001) Influência da nutrição mineral na intensidade da mancha olho pardo em mudas de café. *Pesquisa Agropecuária Brasileira*, 36:53-60.
- Pozza EA & Pozza AAA (2012) A Nutrição Mineral no Manejo de Doenças de Plantas. In: Grupo de Estudos Avançados em Fitopatologia (Ed.). Nutrição no manejo de doenças de plantas. Viçosa, GEAFIP. p.177-212.
- Reis EM, Casa RT & Bresolin ACR (2004) Manual de diagnóstico e controle de doenças do milho. 2ª ed. Lages, Graphel. 141p.
- Ribeiro AC, Guimarães PTG & Alvarez VVH (1999) Recomendações para uso de corretivos e fertilizantes em Minas Gerais – 5ª Aproximação. Viçosa, Universidade Federal de Viçosa. 359p.
- Ritchie SW, Hanway JJ & Benson GO (2003) Como a planta de milho se desenvolve. Piracicaba, Potafos. 20p. (Informações Agrônomicas, 103).
- Rolim GS, Pedro Júnior MJ, Fantin GM, Brunini O, Duarte AP & Dudienas C (2007) Modelo agrometeorológico regional para a estimativa da severidade da mancha de *Phaeosphaeria* em milho safrinha no estado de São Paulo, Brasil. *Bragantia*, 66:721-728.
- SAS Institute Inc. (2002) Statistical Analysis System user's guide. Version 9.0. Cary, Statistical Analysis System Institute. 513p.
- Shanner G & Finney RE (1977) The effect of nitrogen fertilization on the expression of slow-mildewing resistance in Knox wheat. *Phytopathology*, 67:1051-1056.
- Silva EC, Buzetti S, Guimarães GL, Lazarini E & Sá ME (2005) Doses e épocas de aplicação de nitrogênio na cultura do milho em plantio direto sobre Latossolo Vermelho. *Revista Brasileira de Ciência do Solo*, 29:353-362.
- Silva EC, Muraoka T, Buzetti S & Trivelin PCO (2006) Manejo de nitrogênio no milho em Latossolo Vermelho sob plantio direto com diferentes plantas de cobertura. *Pesquisa Agropecuária Brasileira*, 41:477-486.
- Silva HP & Menten JOM (1997) Manejo integrado de doenças na cultura do milho. In: Fancelli & Dourado Neto (Eds.) Simpósio sobre a cultura do milho. Piracicaba, ESALQ/FEALQ. p.40-56.
- Sochting HP & Verret JA (2004) Effects of cultivation systems: soil management, nitrogen fertilization on the epidemics of fungal diseases in oil seed rape (*Brassica napus* L. var. *napus*). *Journal of Plant Diseases and Protection*, 111:01-29.
- Sousa DMG & Lobato E (2004) Calagem e adubação para culturas anuais e semiperenes. In: Sousa DMG & Lobato E (Eds.) Cerrado: correção do solo e adubação. Planaltina, Embrapa Cerrados. p. 283-315.
- Souza JC & Duarte JM (2002) Reação de cultivares de milho a *Phaeosphaeria maydis*. *Ciência e Agrotecnologia*, 26:325-331.
- Taiz L & Zeiger E (2013) Fisiologia Vegetal. 5ª ed. Porto Alegre, Artmed. 918p.
- Tanaka MAS, Passos FA, Feitosa CT & Tanaka RT (2002) Efeito da adubação mineral e orgânica do morangueiro sobre a antracnose do rizoma, causada por *Colletotrichum fragariae*. *Summa Phytopathologica*, 28:236-241.

- Tomazela AL, Favarin JL, Fancelli AL, Martin TN, Dourado Neto D & Reis AR (2006) Doses de nitrogênio e fontes de Cu e Mn suplementar sobre a severidade da ferrugem e atributos morfológicos do milho. *Revista Brasileira de Milho e Sorgo*, 5:192-201.
- Tutida I, May-De Mio LL, Motta ACV & Rosa JMC (2007) Incidência e severidade do “furo de bala” em folhas da ameixeira sob doses de nitrogênio e potássio. *Ciência Rural*, 37:1227-1234.
- Vaz-de-Melo A, Afférri FS, Dotto MA, Peluzio JM, Santos GR & Carvalho EV (2010) Reação de híbridos de milho à *Curvularia ssp* sob dois níveis de adubação com nitrogênio no sul do Tocantins. *Scientia Agrária*, 11:149-154.