



Revista Argentina de Cardiología

ISSN: 0034-7000

revista@sac.org.ar

Sociedad Argentina de Cardiología
Argentina

DIONISIO, GABRIEL F.; GÓMEZ, GASTÓN; MACÍN, STELLA MARIS; GERONAZZO,
RICARDO J.; KEVORKIAN, RUBÉN; FERREIRÓS, ERNESTO; BLANCO, PATRICIA;
HIGA, CLAUDIO

Comparison of two Registries of Acute Coronary Syndrome in Argentina: STRATEG-SIA
and SCAR (1999 – 2011)

Revista Argentina de Cardiología, vol. 83, núm. 4, agosto, 2015, pp. 299-303

Sociedad Argentina de Cardiología
Buenos Aires, Argentina

Available in: <http://www.redalyc.org/articulo.oa?id=305341287007>

- How to cite
- Complete issue
- More information about this article
- Journal's homepage in redalyc.org

redalyc.org

Scientific Information System

Network of Scientific Journals from Latin America, the Caribbean, Spain and Portugal

Non-profit academic project, developed under the open access initiative

Comparison of two Registries of Acute Coronary Syndrome in Argentina: STRATEG-SIA and SCAR (1999 – 2011)

Comparación de dos Registros de Síndrome Coronario Agudo en Argentina: STRATEG-SIA y SCAR (1999 – 2011).

GABRIEL F. DIONISIO^{†, 1}, GASTÓN GÓMEZ^{†, 2}, STELLA MARIS MACÍN³, RICARDO J. GERONAZZO^{MTSAC, 4}, RUBÉN KEVORKIAN^{MTSAC, 5}, ERNESTO FERREIRÓS^{MTSAC, 6}, PATRICIA BLANCO^{MTSAC, 7}, CLAUDIO HIGA^{MTSAC, 8}

ABSTRACT

Background: The influence of clinical evidence on strategies implemented in the treatment of non-ST-segment elevation acute coronary syndromes (NSTEMACS) is not known in our setting.

Objectives: The aim of this study was to evaluate the differences in clinical characteristics, strategies adopted, therapeutic management and in-hospital events of NSTEMACS in participating centers from two registries in Argentina.

Methods: Patients included in participating centers of the STRATEG-SIA registry (1999) and SCAR registry (Síndromes Coronarios Agudos en Argentina - 2011) were compared.

Results: We analyzed 238 patients of the STRATEG-SIA registry and 452 of the SCAR registry in 36 centers. Most patients were men and <65 years (SCAR 57%, STRATEG-SIA 54%; $p=ns$). The SCAR group presented higher prevalence of hypertension (75% vs. 60%; $p=0.001$), dyslipidemia (63% vs. 51%; $p=0.003$), chronic heart failure (10.5% vs. 4.6%; $p=0.02$) and history of myocardial revascularization (30% vs. 17%; $p=0.001$). In the SCAR registry, the proportion of moderate and high-risk patients (TIMI risk score 3-4: 48% vs. 37%; 5-7: 18% vs. 8%; $p=0.0001$) was higher and coronary angiography was more frequent (71% vs. 50%; $p=0.0001$), with a twofold increase in the proportion of percutaneous coronary interventions and 50% reduction in the number of myocardial revascularization surgeries. There were no significant differences in the rate of mortality and myocardial infarction during hospitalization (7.2% vs. 5.9%; $p=ns$).

Conclusions: Patients of the SCAR (2011) registry represent a group at higher risk. The differences in the rates of in-hospital events were not statistically significant.

Key words: Acute Coronary Syndrome - Registries - Coronary Artery Disease

RESUMEN

Introducción: En nuestro medio se desconoce cuál ha sido la influencia de la evidencia clínica sobre las estrategias implementadas en el tratamiento de los síndromes coronarios agudos sin elevación del segmento ST (SCASEST).

Objetivos: Evaluar la variación de las características clínicas, las estrategias adoptadas, las conductas terapéuticas y los eventos hospitalarios de los SCASEST en centros que participaron en dos registros realizados en la Argentina.

Material y métodos: Se compararon pacientes incluidos en centros que participaron en los registros STRATEG-SIA (1999) y SCAR (Síndromes Coronarios Agudos en Argentina - 2011).

Resultados: Se analizaron 238 pacientes del registro STRATEG-SIA y 452 del SCAR incluidos en 36 centros. La mayoría eran de género masculino y menores de 65 años (SCAR 57%, STRATEG-SIA 54%; $p=ns$). El grupo SCAR presentó mayor prevalencia de hipertensión arterial (75% vs. 60%; $p=0.001$), dislipidemia (63% vs. 51%; $p=0.003$), insuficiencia cardíaca crónica (10,5% vs. 4,6%; $p=0,02$) y revascularización coronaria previa (30% vs. 17%; $p=0,001$). Con una proporción mayor de puntaje TIMI de riesgo moderado y alto (3-4: 48% vs. 37%; 5-7: 18% vs. 8%; $p=0,0001$), la coronariografía fue más frecuente en el SCAR (71% vs. 50%; $p=0,0001$), duplicándose la angioplastia coronaria y reduciéndose a la mitad las cirugías de revascularización miocárdica. No hubo diferencias significativas en la tasa intrahospitalaria de muerte e infarto (7,2% vs. 5,9%; $p=ns$).

Conclusiones: Los pacientes del registro SCAR (2011) representan un grupo de mayor riesgo. Las diferencias en las tasas de eventos hospitalarios no fueron estadísticamente significativas.

Palabras clave: Síndromes coronarios agudos - Registros - Enfermedad de la arteria coronaria

REV ARGENT CARDIOL 2015;83:299-303. <http://dx.doi.org/10.7775/rac.v83.i4.6218>

Received: 04/01/2015 - Accepted: 06/11/2015

Address for reprints: Gabriel Fernando Dionisio - Azcuénaga 980/4 - (C1115AAD) CABA, Argentina – Phone: 11 5949 3309 - e-mail: gfdionisio75@yahoo.com.ar

Research Area and Council on Cardiovascular Emergency Care of the Argentine Society of Cardiology
^{MTSAC} Full Member of the Argentine Society of Cardiology

[†] To apply as Full Member of the Argentine Society of Cardiology

¹ Cardiac Catheterization Laboratory, Hospital General de Agudos Donación Francisco Santojanni

² Coronary Care Unit, Hospital General de Agudos Donación Francisco Santojanni

³ Coronary Care Unit, Instituto de Cardiología J. F. Cabral, Corrientes

⁴ Department of Cardiology INCOR, La Rioja

⁵ Department of Cardiology, Hospital General de Agudos Donación Francisco Santojanni

⁶ Boehringer Ingelheim Pharma GmbH & Co. KG. Former director of the Council of Clinical Cardiology, Argentine Society of Cardiology.

Former Scientific Secretary of the Research Area, Argentine Society of Cardiology

⁷ Hospital Naval

⁸ Hospital Alemán

Abbreviations

| | | | |
|--------------|--------------------------------------|----------------|--|
| CA | Coronary angiography. | MI | Myocardial infarction |
| CABGS | Coronary artery bypass graft surgery | NSTEACS | Non-ST segment elevation acute coronary syndrome |

INTRODUCTION

Cardiovascular disease is the most relevant health care problem in developed countries. (1, 2) According to the World Health Organization, cardiovascular disease represents the leading cause of morbidity and mortality worldwide, accounting for 16% of deaths in developed countries and 12% in developing or under-developed countries. (3) Despite identifying and treating cardiovascular risk factors as smoking, cholesterol levels, blood pressure, diabetes and body weight have significantly reduced the incidence of mortality and myocardial infarction (MI), the prevalence of ischemic events related to disease progression is still important. (4, 5)

Different registries have reported that unstable angina is the leading cause of hospitalization due to coronary artery disease in our country. (5-8) The explosive development of innovative technologies and drugs, and the implementation of new strategies for diagnosis and treatment has generated not only a con-

stant review of the management adopted but also an update of the national and international guidelines and recommendations. (9-12) However, the impact in our setting has not been consistently evaluated.

The goal of this analysis was to compare the differences in the clinical characteristics, management strategies and rate of in-hospital events in patients with non-ST-segment elevation acute coronary syndrome (NSTEACS) enrolled in centers participating in two registries performed with a 12-year difference (STRATEG-SIA in 1999 and SCAR in 2011).

METHODS


This comparative analysis of non-ST-segment elevation acute coronary syndromes was performed by members of the Research Area and of the Council on Cardiovascular Emergency Care of the Argentine Society of Cardiology. Patients were enrolled in 36 nationwide centers which had participated in both registries.

The STRATEG-SIA registry was developed between

Table 1. Baseline characteristics of both populations

| Variable | STRATEG-SIA | SCAR | p |
|--|-------------|------------|-------|
| Age, years | 64 (55-74) | 66 (56-73) | ns |
| Female gender, n (%) | 29 (68) | 26.3 (119) | ns |
| Diabetes, % (n) | 24 (57) | 23.9 (108) | ns |
| Smoking habits, n (%) | 35.7 (85) | 39.4 (178) | ns |
| Dyslipidemia, n (%) | 51 (121) | 63 (285) | 0.003 |
| Hypertension, n (%) | 60 (143) | 75 (339) | 0.001 |
| Previous heart failure, % (n) | 4.6 (11) | 10.5 (47) | 0.02 |
| Previous myocardial revascularization, % (n) | 17 (40) | 30 (136) | 0.001 |
| Chronic stable angina, % (n) | 34 (81) | 12 (54) | 0.001 |

Fig. 1. Distribution of the population by TIMI risk score



| Variable | STRATEG-SIA | SCAR | p |
|---------------------------------|-------------|----------|---------|
| Patients studied with CA, n | 116 | 310 | |
| Reason | | | |
| High-risk at admission, % (n) | 32 (38) | 50 (156) | ns |
| Recurrent ischemia, % (n) | 22 (26) | 9 (29) | < 0.001 |
| Positive functional test, % (n) | 22 (26) | 9 (29) | < 0.04 |
| Medical decision, % (n) | 14 (17) | 19 (59) | ns |
| PCI <6 months, % (n) | 0 (0) | 3 (9) | ns |
| Diagnosis, % (n) | 5 (6) | 0 (0) | ns |
| Only positive troponin, % (n) | 0 (0) | 7.4 (23) | ns |
| No data, % (n) | 5 (3) | 2.6 (5) | ns |

CA: Coronary angiography. PCI: Percutaneous coronary intervention.

Table 2. Reasons for indicating coronary angiography

| Variable | STRATEG-SIA | SCAR | p |
|-----------------------------|-------------|-----------|--------|
| Mortality and AMI, % (n) | 5.9 (14) | 7.2 (32) | 0.23 |
| Mortality, % (n) | 1.7 (4) | 3.5 (16) | 0.18 |
| AMI and reinfarction, % (n) | 5 (12) | 3.8 (17) | 0.43 |
| Refractory angina, % (n) | 7.1 (17) | 1.2 (5) | 0.0001 |
| Recurrent angina, % (n) | 20.2 (48) | 10.2 (46) | 0.0001 |

AMI: Acute myocardial infarction.

Table 3. Comparison of the rate of events in both registries

March and September 1999. The registry included 492 patients from 77 institutions with the goal of making a situational diagnosis of the strategies used for the management of patients with NSTEMACS in Argentina, defining the factors associated and/or the determinants which influence decision-making, and to know their impact. (6)

The SCAR registry was performed between June and August 2011 with the participation of 87 centers including 868 patients with NSTEMACS.

Inclusion criteria for the analysis

The patients included were enrolled in centers participating of the STRATEG-SIA and SCAR registries with a final diagnosis of NSTEMACS. In order to attenuate the biases arising while comparing patients enrolled in two registries during two different periods, the same inclusion criteria and exclusion criteria of the STRATEG-SIA were used. In the same sense, the same definitions of coronary risk factors (diabetes, hypertension, high cholesterol levels and smoking habits) and of in-hospital events as myocardial infarction (using the definition of the World Health Organization) of the STRATEG-SIA registry were applied to the database of the SCAR registry. (6)

Study design

Both registries (STRATEG-SIA and SCAR) were cross-sectional observational studies which enrolled consecutive patients with NSTEMACS. A comparative model adjusted for the type of center was used for this analysis. The patients were categorized in tertiles according to the TIMI risk score: low risk (0-1), moderate risk (2-4) and high risk (TIMI 5-7). (13)

Statistical analysis

The distribution of continuous variables was evaluated with

the Kolmogorov-Smirnov test, with measurement of kurtosis and skewness. Continuous variables were presented as mean \pm standard deviation, or median and interquartile range (IQR 25-75), according to their distribution. Comparisons were made using Student's t test and ANOVA, or the Mann-Whitney-Wilcoxon test, as applicable.

Discrete variables were expressed as percentages, and were compared using the chi square test. A two-tailed p value < 0.05 was considered statistically significant. All the statistical analyses were performed using SPSS 19.0 statistical package for Windows (SPSS-IBM, 2010, IL, US).

Ethical considerations

The protocol was approved by SAC's Ethics Committee. As no personal or follow-up data were required, it was not necessary to include an informed consent in the registry, in accordance to the National Habeas Corpus Act Number 25326 (Protection of Personal Data).

RESULTS

Population characteristics

The database consisted of 690 patients, 452 SCAR registry patients and 238 STRATEG-SIA patients, which fulfilled the criteria to be incorporated in the analysis of 36 centers participating in both studies.

Table 1 shows baseline population characteristics. Median age was similar in both studies: 64 years (55-74) in the STRATEG-SIA registry and 66 years (56-73) in the SCAR registry. The proportion of patients >65 years was also similar: 46% in the STRATEG-SIA registry vs. 43% SCAR registry (p=ns). There were no significant gender differences between both registries.

The patients enrolled in 2011 had higher prevalence of hypertension, dyslipidemia, heart failure and history of myocardial revascularization. The patients enrolled in the 1999 registry had higher prevalence of chronic stable angina. The characteristics of both populations were not significantly different from those reported by international NSTEMI registries. (14-17)

ST-segment depression was more frequent in patients enrolled in 2011 (25% vs. 18%; $p=0.05$). There were no statistically significant differences in the other electrocardiographic abnormalities. A high TIMI risk score was more common in patients of the SCAR registry (TIMI 0-2: 36% vs. 55%; TIMI 3-4: 46% vs. 37%; TIMI 5-7: 18% vs. 8%; $p=0.0001$) (Figure 1).

Treatment

Anticoagulant therapy (heparin) was more commonly used in the SCAR registry [328/452 patients (72.5%)] than in the STRATEG-SIA registry [123/238 patients (51.6%); $p=0.001$] with a greater use of low-molecular-weight heparin [SCAR: 268/328 patients (81.7%); STRATEG-SIA: 56/123 patients (45.5%); $p=0.002$].

In both registries, high risk at admission was the most frequent indication of coronary angiography (CA) increasing from 33% in the STRATEG-SIA registry to 50% in the SCAR registry. Recurrent ischemia and a high-risk functional test were the second most frequent indications in the STRATEG-SIA registry while medical team decision was the second reason in the SCAR registry (Table 2).

In 1999, one out of every two patients underwent CA, and by 2011 the procedure was performed to almost three out of every four patients (STRATEG-SIA 50%; SCAR 71%; $p=0.0001$). During the same period, the comparison between both registries showed a significant reduction in the implementation of coronary artery bypass graft surgery (CABGS) (13% vs. 7.5%, $p=0.0001$) and the rate of percutaneous coronary interventions evidenced a two fold increase (22% vs. 45%; $p=0.0001$).

The incidence of recurrent and refractory angina was significantly lower in the SCAR versus the STRATEG-SIA registry. There were no significant differences in the incidence of other cardiovascular events (Table 3)

DISCUSSION

The present analysis compares demographic and clinical characteristics, diagnostic and therapeutic management and the outcome of NSTEMI in patients enrolled in the participating centers of two registries conducted by the Argentine Society of Cardiology in 1999 and 2011.

After 12 years, the demographic pattern of the population studied does not seem to be significantly different; most patients are still middle-aged men. However, the prevalence of some cardiovascular risk factors, particularly hypertension and dyslipidemia, seems to be higher. Probably, this finding could be at-

tributed to a greater capability of detecting these factors in the population, as it happened with surveys performed by the National Ministry of Health. (18, 19)

The patients enrolled in the SCAR registry had history of higher cardiovascular risk and presented with a more severe ischemic risk.

The diagnostic and therapeutic management was more aggressive in the SCAR registry, and the use of anticoagulation therapy was more common, particularly with low-molecular-weight heparins. This approach is consistent with the one reported by large international multicenter registries. The indication of invasive treatment was higher and also similar to the rate indicated by international reports. (15-17, 20)

Interestingly, percutaneous coronary intervention was the treatment of choice, and the indication of CABGS was lower than in 1999.

The first reason to indicate CA was high clinical risk at admission in both registries. The second reason was recurrent ischemia or high-risk functional test in the STRATEG-SIA registry, and decision of the medical team in the SCAR registry (see Table 2).

There were no significant differences in terms of mortality, myocardial infarction or reinfarction between both registries.

Study limitations

The presence of differences in the indication of procedures or the absence of differences in mortality in observational studies cannot be considered definitive, particularly when the analysis was not adjusted due to the low number of events

CONCLUSIONS

The burden of coronary risk factors and of history of heart disease associated with a higher prevalence of moderate and high TIMI risk score would indicate that the risk of patients enrolled in the SCAR registry was higher.

The diagnostic and therapeutic approach was different in the period between both registries.

Coronary angiography increased with a two-fold rise in the number of percutaneous coronary interventions and a reduction in the number of patients undergoing CABGS that is similar to the most recent clinical evidence.

Despite the risk profile was different in both populations, the differences in the rate of in-hospital events were not significant.

Conflicts of interest

None declared

(See author's conflicts of interest forms in the web / Supplementary Material)

REFERENCES

1. World Health Organization. "Ten leading causes of death in 2008". 2011 July 18. Disponible en: http://gamapserver.who.int/gho/interactive_charts/mbd/cod_2008/graph.html

2. Araújo F, Gouvêas C, Fontes F, La Vecchia C, Azevedo A, Lunet N. Trends in cardiovascular diseases and cancer mortality in 45 countries from five continents (1980-2010). *Eur J Prev Cardiol* 2013;21:1004-17. <http://doi.org/5vn>
3. Doval HC, Tajer CD, Ferrante D. Evidencias en Cardiología III. De los ensayos Clínicos a las Conductas Terapéuticas. Buenos Aires: GEDIC; 2003. p. 43-4.
4. Vetrovec GW. Optimizing percutaneous coronary intervention outcomes. The next steps. *Circulation* 2005;111:125-6. <http://doi.org/dccx7m>
5. Encuesta Nacional de Unidades Coronarias. *Rev Argent Cardiol* 1991;59(Suppl 2):7-40.
6. Ferreirós ER, Fuselli JJ, Guetta J, Boissonnet CP, Di Toro D, Cragnoletto R et al. Results of the First National, Multicenter and Prospective Study on Management Strategies of Acute Ischemic Syndromes without ST-Segment Elevation in Argentina (STRATEG-SIA Study). *Rev Argent Cardiol* 2001;69:11-33.
7. Piombo A, Charask A, Botto F, Llois S, Fuselli JJ. Acute Ischemic Syndromes without ST Segment Elevation. First Internet Survey of Coronary Care Units. *Rev Argent Cardiol* 2003;71:396-401.
8. Mauro V, Charask A, Gitelman P, Salzbeg S, Bruno C, Prieto N, et al. Analysis of the Evolution of the AMI in the Past 14 years in Argentina. Therapeutic Indications. *Rev Argent Cardiol* 2001;69:591-601.
9. Sans S, Kesteloot H, Kromhout D. The burden of cardiovascular diseases mortality in Europe. Task Force of the European Society of Cardiology on Cardiovascular Mortality and Morbidity Statistics in Europe. *Eur Heart J* 1997;18:1231-48. <http://doi.org/5vp>
10. Bertrand ME, Simoons ML, Fox KA, Wallentin LC, Hamm CW, McFadden E, et al. Management of acute coronary syndromes in patients presenting without persistent ST-segment elevation. *Eur Heart J* 2002;23:1809-40. <http://doi.org/bkb9ck>
11. Amsterdam EA, Wenger NK, Brindis RG, Casey DE Jr, Ganiats TG, Holmes DR Jr, et al. 2014 AHA/ACC guideline for the management of patients with non-ST-elevation acute coronary syndromes: executive summary: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines. *Circulation* 2014;130:2354-94. <http://doi.org/5vq>
12. Sociedad Argentina de Cardiología. Consenso para el Manejo de Pacientes con Síndrome Coronario Agudo sin Supradesnivel del Segmento ST (Angina Inestable e Infarto de Miocardio sin elevación del ST). *Rev Argent Cardiol* 2014;82(Supl 1):1-44.
13. Antman EM, Cohen M, Bernink PJ, McCabe CH, Horacek T, Papuchis G, et al. The TIMI risk score for unstable angina/non-ST elevation MI: a method for prognostication and therapeutic decision making. *JAMA* 2000;284:835-42. <http://doi.org/fngdg5>
14. Nicolau JC, Franken M, Lotufo PA, Carvalho AC, Marin Neto JA, Lima FG, et al. Use of Demonstrably Effective Therapies in the treatment of acute coronary syndromes: comparison between different Brazilian regions. Analysis of the Brazilian Registry on Acute Coronary Syndromes (BRACE). *Arq Bras Cardiol* 2012;98: 282-9. <http://doi.org/5vr>
15. Al-Aqeedi R, Sulaiman K, Al Suwaidi J, Alhabib K, El-Menyar A, Panduranga P, et al. Characteristics, management and outcomes of patients with acute coronary syndrome and prior coronary artery bypass surgery: findings from the second Gulf Registry of Acute Coronary Events. *Interact Cardiovasc Thorac Surg* 2011;13:611-8. <http://doi.org/dnm68m>
16. Boersma E, Pieper KS, Steyerberg EW, Wilcox RG, Chang WC, Lee KL, et al. Predictors of outcome in patients with acute coronary syndromes without persistent ST-segment elevation: Results from an international trial of 9461 patients. *Circulation* 2000;101:2557-67. <http://doi.org/5vs>
17. Fox KA, Goodman SG, Anderson FA Jr, Granger CB, Moscucci M, Flather MD. From guidelines to clinical practice: the impact of hospital and geographical characteristics on temporal trends in the management of acute coronary syndromes. The Global Registry of Acute Coronary Events (GRACE). *Eur Heart J* 2003;24:1414-24. <http://doi.org/bnwdmh>
18. Segunda Encuesta Nacional de Factores de Riesgo Para Enfermedades no Transmisibles. Disponible en: http://www.msal.gov.ar/ent/images/stories/vigilancia/pdf/fr_encuesta-nacional-factores-riesgo-2011
19. Tercera Encuesta Nacional de Factores de Riesgo para Enfermedades no Transmisibles. Presentación de Principales Resultados. Disponible en: <http://www.msal.gov.ar/images/stories/publicaciones/pdf/11.09.2014-tercer-encuentro-nacional-factores-riesgo>
20. Gierlotka M, Gasior M, Wilczek K, Wasilewski J, Hawranek M, Tajstra M, et al. Temporal trends in the treatment and outcomes of patients with non ST-segment elevation myocardial infarction in Poland from 2004-2010 (from the Polish Registry of Acute Coronary Syndromes). *Am J Cardiol* 2012;109:779-86. <http://doi.org/ffrgcp>