



Acta Scientiarum. Health Sciences

ISSN: 1679-9291

eduem@uem.br

Universidade Estadual de Maringá

Brasil

Garcia, Flávio Roberto Mello; Lise, Fernanda

Ants associated with pathogenic microorganisms in brazilian hospitals: attention to a silent vector

Acta Scientiarum. Health Sciences, vol. 35, núm. 1, enero-junio, 2013, pp. 9-14

Universidade Estadual de Maringá

Maringá, Brasil

Available in: <http://www.redalyc.org/articulo.oa?id=307226203002>

- How to cite
- Complete issue
- More information about this article
- Journal's homepage in redalyc.org

redalyc.org

Scientific Information System

Network of Scientific Journals from Latin America, the Caribbean, Spain and Portugal

Non-profit academic project, developed under the open access initiative



## Ants associated with pathogenic microorganisms in Brazilian hospitals: attention to a silent vector

Flávio Roberto Mello Garcia\* and Fernanda Lise

Laboratório de Ecologia de Insetos, Departamento de Zoologia e Genética, Instituto de Biologia, Universidade Federal de Pelotas, Cx. Postal 354, 96010-900, Pelotas, Rio Grande do Sul, Brazil. \*Author for correspondence. E-mail: [flavio.garcia@pq.cnpq.br](mailto:flavio.garcia@pq.cnpq.br)

**ABSTRACT.** The presence of ants in hospitals is a threat to the health of the hospital community, given the potential to carry pathogenic microorganisms. This study was developed in order to comment on publications related to the occurrence of ants associated with pathogenic microorganisms in hospitals in Brazil. Studies indicated that ants were collected in several wards of 13 hospitals in five States from South and Southeast regions. The microbiological analysis has identified on the ants, microorganisms such as *Staphylococcus*, *Streptococcus*, *Klebsiella*, *Escherichia coli*, *Pseudomonas*, and others with microbiological importance.

**Keywords:** *Monomorium pharaonis*, *Staphylococcus*, hospital infection, public health.

## Formigas associadas a microorganismos patogênicos em hospitais brasileiros: a atenção para um vetor

**RESUMO.** A presença de formigas em hospitais constitui risco a saúde da comunidade hospitalar, uma vez que possuem o poder de carrear microorganismos patogênicos. Este estudo foi realizado com o objetivo de comentar publicações referentes à ocorrência de formigas associadas a microorganismos patogênicos em hospitais no Brasil. Os estudos indicam que foram coletadas formigas em diversas unidades hospitalares de 13 hospitais, em cinco estados nas regiões Sul e Sudeste. A análise microbiológica permitiu identificar nas formigas microorganismos como *Staphylococcus*, *Streptococcus*, *Klebsiella*, *Escherichia coli*, *Pseudomonas*, *Streptococcus* entre outros de importância microbiológica.

**Palavras-chave:** *Monomorium pharaonis*, *Staphylococcus*, infecção hospitalar, saúde pública.

### Introduction

Ants are social insects found almost everywhere and have higher diversity in tropical regions. These insects cause problems in hospitals worldwide (BEATSON, 1972), was the first reporting the occurrence of ants in nine hospitals in the United Kingdom, then in England (EDWARD; BACKER, 1981), Chile (IPINZA-REGLA et al., 1981); Germany (EICHELER; 1990) and Trinidad (CHADEE; MAITRE, 1990), Colombia (OLAYA-MASMELA et al., 2005), Spain (ESPALDER; ESPEJO, 2002), United States (KLOTZ et al., 1995; NELDER et al., 2006), Malaysia (NA; LEE, 2001), and Korea (KIM et al., 2005).

In Brazil, studies have initiated in the 90's with the identification of 14 ant species (FOWLER et al., 1993), Bueno and Fowler (1994) investigated 20 hospitals in Brazil, and the ant fauna in each hospital consisted of between 10 and 23 species.

Among the factors affecting the presence of ants in hospitals, highlights the architectural structure, proximity to residences, weather interferences,

temperature variations that stimulate the migration of these insects to electronic devices searching for thermal stability and/or packages of drugs that provide suitable conditions for nesting, besides foodstuffs that function as extra attractive (BEATSON, 1972; ZARZUELA et al., 2002).

With the growing concern about the occurrence of ants in these environments, several studies have been undertaken in Brazil aiming to verify the potential of these organisms to carry pathogenic microorganisms, and the results have indicated their role as mechanical vector, contaminating the environment and collaborating to nosocomial infections (BUENO; FOWLER, 1994; FOWLER et al., 1993, MOREIRA et al., 2005). This discussion is justified because it is essential that health professionals to know and be aware of the need to control and monitor the presence of these insects, since currently hospital infections are a serious public health problem in the country.

Although ants are not considered the major source of hospital infections, are viewed as any other vehicle of transmission (FOWLER et al., 1993). For

immunosuppressed patients, susceptible to develop infections, this exposure may change the prognosis, increasing the length and cost of hospitalization (IPINZA-REGLA et al., 1981).

This study examined the occurrence of ants in Brazilian hospital environments over the last decade, by means of a literature review, focusing on the distribution and potential of ants to carry pathogenic microorganisms, once ants in hospital settings may transport diverse bacterial species (gram-positive and gram-negative), considering that some strains may be multidrug-resistant. The Table 1 lists results from 13 studies performed in Brazil, with the species of ants collected in hospitals, microorganisms carried by them, and hospital wards in which they were collected.

## Material and methods

For this study we used the systematic review using meta-analysis to integrate the results, allowing to describe current knowledge, indicating those aspects that are science-based and those that do not have a solid base of support and require further investigation.

The criteria for selection and inclusion of articles has occurred as the objectives of this study to maintain the rigor and uniformity in the choice of these were complied with some criteria: articles that address the topic, articles indexed in databases LILACS (Latin American Literature in Health Sciences) and MEDLINE (Medical Literature Analysis and Retrieval Sistem online); articles published in national journals within the defined time period from 2001 to 2011.

The key words used for this literature in LILACS and MEDLINE were ants, hospital and vectors.

## Results and discussion

The ant diversity registered in Brazil with potential to act as mechanical vector of pathogenic microorganisms in hospital environments is significantly higher than in European countries. *M. pharaonis* is the species found in European hospitals (BEATSON, 1972; EICHLER, 1990; ULLOA, 2003). Fowler et al. (1993) considered that the ant species predominant in Brazil are the exotic *M. pharaonis* and *T. melanocephalum*.

Surveys in five Brazilian states indicated that some species are predominant in hospitals, and the occurrence of *T. melanocephalum* was predominant (TANAKA et al., 2007; TEIXEIRA et al., 2009), followed by *P. longicornis*. These and other species listed in Table 1 are mechanical vectors for nosocomial infections, since were identified pathogens such as *Staphylococcus*, *Enterococcus*,

*Klebsiella*, *Escherichia coli*, *Pseudomonas* and *Streptococcus* carried by them in different hospital wards (BELEI et al., 2006; COSTA et al., 2006; FOWLER et al., 1993; GARCIA et al., 2011; LISE et al., 2006; MOREIRA et al., 2005; PESQUERO et al., 2008; RANDO et al., 2009; SANTOS et al., 2009; TANAKA et al., 2007; TEIXEIRA et al., 2009).

The identification of pathogenic microorganisms in the ants analyzed, along with biological traits, can define ants as potential carriers of pathogens and the infestation in hospitals is a risk to public health (BEATSON, 1972). The association between enterobacteria and ants has been observed in some Brazilian hospitals, which always represents a risk to hospitalized patients. Enterobacteria are common among insects, and horizontally transmitted from one to another individual, but can be acquired from the environment, showing thus capacity of propagation and maintenance of the microorganism in the environment (MOREIRA et al., 2005; PEREIRA; UENO, 2008; PESQUERO et al., 2008).

Ants can move upon human material such as urine, feces, sputum, and carry the microorganisms to the utensils and surfaces in general. Among the microorganisms identified in the ants is the group of fecal coliforms, including *Escherichia*, *Enterobacter* and *Klebsiella*, present in human feces by being part of the intestinal flora; and *Staphylococcus aureus*, on the human skin and nasopharynx, is the responsible for the most of nosocomial infections (TRABULSI, 1991).

Regarding the *Escherichia coli*, even being part of the gastrointestinal tract of humans it has been reported as one of the most important agents for extraintestinal infections, such as diarrhea in adults and children, and experimental keratoconjunctivitis, an infection similar to shigellosis (JAWETZ et al., 1998). Thus, the problem becomes relevant in the studied hospitals, once this bacterium was identified in ants *Odontomachus* sp and *Pheidole* sp1 and sp2 found in areas with the presence of children, such as nursery and maternity (SANTOS et al., 2009).

Ants have great capacity to maintain the association between bacteria and fungi, including some antibiotic-resistant pathogenic species, characterizing a condition of risk of nosocomial infection (PANTOJA et al., 2009; PEREIRA; UENO, 2008). Among the multiresistant strains, stand out the genera *Acinetobacter*, *Streptococcus*, *Gemella*, *Enterococcus faecalis* and *Klebsiella* (MOREIRA et al., 2005; TRABULSI, 1991) and *K. Pneumoniae* (TANAKA et al., 2007). Thus, it is evident the importance of the problem in the public health area, once these bacteria are becoming increasingly resistant to a higher number of antibiotics.

**Table 1.** Species of ants, microorganisms carried by them, and hospital units examined by the 14 Brazilian studies.

Title	Authors	Ant species	Microorganisms	Hospital unit
Ants as carriers of resistant bacteria in hospitals	Moreira et al. (2005)	<i>Tapinoma melanocephalum</i> ; <i>Paratrechina longicornis</i> ; <i>Monomorium pharaonis</i> ; <i>Solenopsis saevissima</i> ;	<i>Bacillus</i> spp.; <i>Enterobacter amnigenus</i> ; <i>Enterobacter cloacae</i> ; <i>Enterococcus faecalis</i> ; <i>Klebsiella pneumoniae</i> ; <i>Staphylococcus saprophyticus</i> ; <i>Acinetobacter baumannii</i> ; <i>Gemella morbillorum</i> ; <i>Staphylococcus epidermidis</i> ; <i>Staphylococcus equorum</i> ; <i>Klebsiella oxytoca</i> ; <i>Staphylococcus aureus</i> ; <i>Staphylococcus simulans</i> ; <i>Staphylococcus warneri</i> ; <i>Serratia rubidae</i> <i>Staphylococcus cohnii</i> <i>Bacillus</i> spp. <i>Enterobacter agglomerans</i> ; <i>Gemellaha emolysans</i> ; <i>Enterococcus faecium</i> ; <i>Gemellaha emolysans</i> ; <i>Streptococcus acidominimus</i> ; <i>Staphylococcus lugdunensis</i> .	Adult and pediatric wards, Cardiology, gynecology, trauma, postoperative; Adult and pediatric ICU.
Occurrence, behavior, and vectoring of fungi by ants in the hospital of the Federal University of Juiz de Fora, Minas Gerais State	Barros et al. (2006)	<i>Camponotus</i> sp., <i>Camponotus crassus</i> ; <i>C. atriceps</i> ; <i>C. renggeri</i> ; <i>Wasmannia auropunctata</i> ; <i>Paratrechina longicornis</i> ; <i>M. floricola</i> ; <i>Pheidole</i> sp., <i>Tapinoma melanocephalum</i> ; <i>Odontomachus</i> sp.	<i>Aspergillus</i> sp.; <i>Cladosporium</i> sp.; <i>Penicillium</i> sp.; <i>Candida</i> sp. and <i>Aspergillus niger</i> .	Not specified
Ants: analysis on microorganisms carried in the hospital environment	Belei et al. (2006)	Not specified	<i>Filamentous fungi</i> , <i>Coagulase negative staphylococci</i> , <i>acinetobacter</i> sp.; <i>micrococcus</i> sp.; and <i>bacillus</i> sp..	NICU and other hospital units.
Ants as mechanical vectors of microorganisms in the school hospital of the Federal University of triângulo mineiro	Costa et al. (2006)	<i>Tapinoma melanocephalum</i> ; <i>Pheidole</i> sp. and <i>Paratrechina longicornis</i> .	<i>Staphylococcus</i> sp.; Gram-positive bacilli, <i>Pseudomonas</i> sp.; and <i>Micrococcus</i> sp..	Wards, service of nutrition and dietetics, and pediatric ICU
Association between ants (Hymenoptera: Formicidae) and bacteria in hospitals of Santa Catarina State	Lise et al. (2006)	<i>Monomorium pharaonis</i> ; <i>Solenopsis saevissima</i> ; <i>Paratrechina longicornis</i> ; <i>Brachymyrmex</i> sp.; <i>Solenopsis</i> sp.; <i>Camponotus</i> sp.; <i>Tapinoma melanocephalu</i> .	<i>Acinetobacter</i> sp.; <i>Acinetobacter haemolyticus</i> ; <i>Oerskovia</i> sp.; <i>Corynebacterium</i> sp.; <i>Corynebacterium diphtheriae</i> ; <i>Corynebacterium jeikeium</i> ; <i>Enterococcus</i> sp.; <i>Listeria monocytogenes</i> ; <i>Neisseria</i> sp.; <i>Planococcus</i> sp.; <i>Pseudomonas luteola</i> ; <i>Sphingobacterium</i> sp.; <i>Sphingomonas paucimobilis</i> ; <i>Staphylococcus</i> sp.; <i>Staphylococcus intermedius</i> ; <i>Staphylococcus saprophyticus</i> ; <i>Stenotrophomonas maltophilia</i> ; <i>Streptococcus agalactiae</i> ; <i>Streptococcus bovis</i> ; <i>Neisseria</i> sp.; <i>Planococcus</i> sp. and <i>S. agalactiae</i> .	Adult inpatient unit, surgical, oncology and pediatrics.
Urban ants and the transport of nosocomial bacteria	Rodvalho et al. (2007)	<i>Tapinoma melanocephalum</i> and <i>C. vittatus</i>	Coagulase positive staphylococci, Coagulase negative staphylococci and Gram-negative bacilli.	Unit of infectious diseases, emergency room, and burn unit
Bacteria carried by ants in hospital environment	Tanaka et al. (2007)	<i>Monomorium pharaonis</i> and <i>Tapinoma melanocephalum</i> .	<i>Corinebacterium</i> sp.; <i>Klebsiella pneumoniae</i> ; <i>K. ozaenae</i> ; <i>Escherichia coli</i> .	Nursery, surgical center, nursing station, sterilizing room, blood donation room, and ICU
Ants as microorganism carriers in hospitals	Pereira e Ueno (2008)	125 ants of the same not-specified species	<i>Hafnia alvei</i> ; <i>Klebsiella pneumoniae</i> ; <i>Enterobacter agglomerans</i> ; <i>Enterobacter cloacae</i> ; <i>Enterobacter sakazakii</i> ; <i>Serratia liquefaciens</i> ; <i>Serratia marcescens</i> . <i>Epidermopyton floccosum</i> ; <i>Trichophyton rubrum</i> ; <i>Trichophyton verrucosum</i> ; <i>Cladosporium carruoni</i> ; <i>Aurobasidium pullulans</i> ; <i>Wangiella dermatitidis</i> ; <i>Conidiobolus coronatus</i> ; <i>Fonsecaea pedrosoi</i> ; <i>Aspergillus niger</i> ; <i>Aspergillus flavus</i> ; <i>Aspergillus fumigatus</i> ; <i>Monilia sitophita</i> .	Medical clinic, blood unit, laundry, and orthopedics.

continue...

...continuation

Title	Authors	Ant species	Microorganisms	Hospital unit
Ants in hospital environment and their potential to transmit bacteria	Pesquero et al. (2008)	<i>Pheidole</i> sp1; <i>Hipoponera</i> sp1; <i>Dorymyrmex pyramicus</i> ; <i>Linepithema humile</i> ; <i>Camponotus</i> sp1; <i>Brachymyrmex</i> sp1; <i>Brachymyrmex</i> sp2; <i>Paratrechina fulva</i> ; <i>Cardiocondyla</i> sp1;	<i>Escherichia coli</i> ; <i>Staphylococcus</i> sp.; <i>Enterococcus</i> sp.; <i>Klebsiella</i> sp.; <i>Salmonella</i> sp. and <i>Aeromonas</i> sp.	Restrooms, nursery, medical offices, kitchen, sterilizing room, hematology, laboratory, milk kitchen unit, nursing station, suture room, plaster room, Adult and pediatric ICU
Ants (Hymenoptera: Formicidae) as bacterial vectors in two hospitals of the municipality of Divinópolis, Minas Gerais State	Santos et al. (2009)	<i>Pheidole</i> sp1 and sp2; <i>Linepithema humile</i> ; <i>Wasmannia auropunctata</i> ; <i>Camponotus</i> sp.; <i>Odontomachus</i> sp; <i>Solenopsis</i> sp.; <i>Acromyrmex</i> sp.; and <i>Tapinoma melanocephalum</i> .	<i>Escherichia coli</i> ; <i>Pseudomonas aeruginosa</i> ; pathogenic and non-pathogenic <i>Staphylococcus</i> spp; <i>Streptococcus faecalis</i> and other <i>Enterococcus</i> sp.	Warehouse, nursery, chapel, sterilizing room, surgical center, medical clinic, drugstore, kitchen, bedrooms, laboratory, maternity, ICU, radiology.
Characterization of ant fauna in establishments of health area in the municipality of Bandeirantes, Paraná State.	Rando et al. (2009)	<i>Tapinoma melanocephalum</i> ; <i>Paratrechina fulva</i> ; <i>Monomorium pharaonis</i> ; <i>C. atriceps</i> ; <i>Brachymyrmex</i> sp.; <i>Pheidole</i> sp.4; <i>Pheidole</i> sp.3; <i>Pheidole</i> sp.2; <i>Pheidole</i> sp.1; <i>Dorymyrmex</i> sp.	<i>Staphylococcus</i> sp.; <i>Serratia</i> sp.; <i>Klebsiella</i> sp.; <i>Escherichia coli</i> ; <i>Salmonella</i> sp.; and <i>Pseudomonas</i> sp..	Surgical center, hallways, drugstore, laundry, nursing stations, reception, adult and pediatric inpatient units, and ICU
Microbiota associated with urban ants in a Brazilian university hospital.	Teixeira et al. (2009)	<i>Tapinoma melanocephalum</i> .	Gram-positive bacilli, Gram-negative bacilli, Gram-positive cocci, <i>Filamentous fungi</i> , <i>Pseudomonas</i> ; <i>Staphylococcus</i> ; and <i>Streptococcus</i> .	Surgical center, and ICU.
Ants (Hymenoptera: Formicidae) as carriers of fungi in hospitals: emphasis on genera <i>Tapinoma</i> and <i>Pheidole</i>	Pantoja et al. (2009)	<i>Tapinoma melanocephalum</i> ; <i>Paratrechina longicornis</i> ; <i>Camponotus</i> ; <i>Solenopsis</i> ; <i>Pheidole</i> .	<i>Absidia</i> sp.; <i>Acremonium</i> sp.; <i>Acremonium hyalinulum</i> ; <i>Aspergillus flavus</i> ; <i>Aspergillus niger</i> ; <i>Aspergillus chraceus</i> ; <i>Aspergillus oryzae</i> ; <i>Aspergillus sydowii</i> ; <i>Aspergillus versicolor</i> ; <i>Chrysosporium inops</i> ; <i>Cladosporium</i> <i>Sphaerospermum</i> ; <i>Cokeromyces</i> sp.; <i>Cunninghamella</i> <i>Bertholletiae</i> ; <i>Cyphellophora</i> sp.; <i>Fusarium</i> sp.; <i>Fusarium proliferatum</i> ; <i>Fusarium solani</i> ; <i>Mortierella polycephala</i> ; <i>Mucor</i> sp.; <i>Mycocentrospora acerina</i> ; <i>Ochroconis gallopava</i> ; <i>Paecilomyces marquandii</i> ; <i>Paecilomyces variotii</i> ; <i>Penicillium</i> sp. <i>Rhinochlamydia aquaspersa</i> ; <i>Scopulariopsis koningii</i> ; <i>Scytalidium</i> sp.; <i>Tritirachium oryzae</i> ; <i>Candida albicans</i> ; <i>Candida glabrata</i> ; <i>Candida guilliermondii</i> ; <i>Candida parapsilosis</i> ; <i>Rhodotorula</i> sp.; <i>Rhodotorula mucilaginosa</i> ; <i>Saccharomyces</i> sp.	NICU, adult ICU, surgical center, transplant units, pediatrics, hematology, surgery, gynecology.
Association between ants (Hymenoptera: Formicidae) and pathogenic bacteria in five hospitals in the city of Pelotas	Gonçalves et al. (2011)	<i>Cardiocondyla emeryi</i> <i>Pheidole nubila</i> , <i>Pheidole spininodis</i> , <i>Pheidole</i> sp.1, <i>S. saevissima</i> , <i>Tetramorium bicarinatum</i> ,	Enterobacteria, <i>Pseudomonas aeruginosa</i> , <i>Pseudomonas fluorescens</i> , <i>Pseudomonas putida</i> , <i>Staphylococcus epidermidis</i> and <i>Staphylococcus saprophyticus</i>	Surgical center, hallways, drugstore, laundry, nursing stations, reception, adult and pediatric inpatient units, and ICU

The inpatient units for adults and children, Intensive Care Units (ICU) for adults and children, as well as nursery, surgical centers, nursing stations,

sterilizing rooms, kitchens and laboratories are among the wards with high infestation of ants (COSTA et al., 2006; LISE et al., 2006; MOREIRA

et al., 2005; PESQUERO et al., 2008; RANDO et al., 2009; SANTOS et al., 2009; TANAKA et al., 2007; TEIXEIRA et al., 2009). Among them, some units with critically ill patients such as ICU, surgical centers and nurseries, receive a lower flow of people and materials, and have a higher frequency of environmental hygiene, then it was expected a lower index of infestation of ants than observed (BRAGANÇA; LIMA, 2010). However, their presence was not related to the lack of cleaning, some species are attracted to sterilized materials, serum and medications used.

Multiresistant bacterial strains isolated from ants in places such as nurseries indicate a direct effect on disease transmission and consequently on increased rates of infection and severity of nosocomial infections. Undoubtedly, this needs to be discussed with sectors related to prevention of hospital infection (TANAKA et al., 2007).

According to these studies, ants in hospitals should be considered a threat to human health, because they are vectors of pathogenic bacteria, but not associate the presence of ants with indices of nosocomial infections, nor neglect their presence, without controlling or monitoring their presence in these environments (CINTRA-SOCOŁOWSKI, 2007).

## Conclusion

The presence of ants in hospitals should receive attention since they can carry diverse bacterial species, including multiresistant strains. In this way, ants in this environment should warn the CCIH and all multidisciplinary team for the effective control of nosocomial infections, because only with the participation of the entire hospital community it will be possible to develop from basic measures, such as hand washing, and the maintenance of a clean working environment, to the control of ants, but the professional awareness comes to the fore as prevention.

This problem can be minimized by adopting some measures to control ants in hospitals, such as the ban on entry of food and flowers, cover small cracks on the walls, removal of tree branches close to the windows and outside walls, removal of debris in the outer area. Because several factors favor the occurrence of ants in hospitals, including the deficient structure, even with efforts of a committee on hospital infection control, given the circulation of a large amount of people (patients, relatives, employees, suppliers) and goods (clothes, foodstuffs, flowers, and other objects) that favor the entry of these insects, in addition to the drugs that attract them.

## References

- BARROS, R. A. M.; CAMPOS-FARINHA, A. E.; PREZOTO, F. Ocorrência, comportamento e vetoração de fungos por formigas no Hospital. **Revista Brasileira de Zoociências**, v. 8, n. 2, p. 217-220, 2006.
- BEATSON, S. H. Pharaoh's ants as pathogens vectores in hospitals. **The Lancet**, v. 1, n.1, p. 425-427, 1972.
- BELEI, R. A.; GUARIENTE, M. H. D. M.; CAVASSIN, E. D.; RIBEIRO, P. H. V. Formigas: análise dos microorganismos carreados no ambiente hospitalar **Nursing**, v. 9, n. 95, p. 777-781, 2006.
- BRAGANÇA, M. A. L.; LIMA, J. D. Composição, abundância e índice de infestação de espécies de formigas em um hospital materno-infantil de Palmas, TO. **Revista Neotropical Entomology**, v. 39, n. 1, p. 124-130, 2010.
- BUENO, O. C.; FOWLER, H. G. Exotic ants and native ant fauna of Brazilian hospitals. In: WILLIAMS, D. F. (Ed.). **Exotic ants: biology, impact and control of introduced species**. Boulder: Westview Press, 1994. p. 191-198.
- CHADE, D.; MAITRE, A. Ants: potential mechanical vectors of hospital infections in Trinidad. **Transactions of the Royal Society of Tropical Medicine and Hygiene**, v. 84, n. 2, p. 297-297, 1990.
- CINTRA-SOCOŁOWSKI, P. Histórico sobre as pesquisas com formigas em ambientes hospitalares no Brasil. **Biológico**, v. 69, p. 35-38, 2007. (Supl. 2).
- COSTA, S. B.; PELLI, G. P.; CARVALHO, P.; OLIVEIRA, A. G.; SILVA, P. R.; TEIXEIRA, M. M.; MARTINS, E.; TERRA, A. P. S.; RESENDE, E. M.; OLIVEIRA, C. C. H. B.; MORAIS, C. A. Formigas como vetores mecânicos de microorganismos no hospital escola da universidade federal do triângulo mineiro. **Revista Sociedade Brasileira de Medicina Tropical**, v. 39 n. 6, p. 527-529, 2006.
- EDWARD, J. P.; BAKER, L. F. Distribution and importance of Tha Pharaoh's *Monomorium pharaonis* (L) in national Health Service Hospitals in England. **Journal of Hospital Infection**, v. 2, n. 2, p. 245-254, 1981.
- EICHELER, W. Health aspects and control of *Monomorium pharaonis*. In: MEER, R. K. V.; JAFE, K.; CEDENO, A. (Ed.). **Applied Mirmecology: a world perspective**. Boulder: Westview Press, 1990. p. 671-675.
- ESPALDER, X.; ESPEJO, F. *Tapinoma melanocephalum* (Fabricius, 1793) a new exotic ant in Spain (Hymenoptera, Formicidae). **Orsis**, v. 17, n. 4, p. 101-104, 2002.
- FOWLER, H. G.; BUENO, O. C.; SADASTSUNE, T.; MONTELLI, A. Ants as potential vectors of pathogens in hospitals in the state of São Paulo, Brazil. **Insect Science and its Applications**, v. 14, n. 3, p. 367-370, 1993.
- GARCIA, F. R. M.; AHLERT, C. C.; FREITAS, B. R.; TRAUTMANN, M. M.; TANCREDI, S. P.; LUTINSKI, J. A. *Ants* (Hymenoptera: Formicidae) in five hospitals of Porto Alegre, Rio Grande do Sul State, Brazil. **Acta Scientiarum. Health Sciences**, v. 33, n. 2, p. 203-209, 2011.
- GONÇALVES, M. G.; LOECK, A. E.; SILVA, W. P.; ROSADO, J. L. O.; BASTOS, C. P.; BASSANI, M. T. Associação entre formigas (Hymenoptera: Formicidae) e

- bactérias patogênicas em cinco hospitais do município de Pelotas, RS. **Arquivos do Instituto Biológico**, v. 78, n. 2, p. 287-295, 2011.
- IPINZA-REGLA, J.; FIGUEROA, G.; OSORIO, J. *Iridomyrmex humilis* 'hormiga argentina como vector de infecciones intrahospitalarias: I Estudio bacteriológico. **Folia Entomológica Mexicana**, v. 50, n. 1, p. 81-96, 1981.
- JAWETZ, E.; MELNICK, J.; ADELBURG, E. A. **Microbiología médica**. Rio de Janeiro: Guanabara Koogan, 1998.
- KIM, C. W.; KIM, D. I.; CHOI, S. Y.; PARK, J. W.; HONG, C. S. Pharaoh ant (*Monomorium pharaonis*); Newly identified important inhalant allergens in bronchial asthma. **Journal Korean of Medical Science**, v. 20, n. 2, p. 390-396, 2005.
- KLOTZ, J. K.; MANGOLD, J. R.; VAIL, K. M.; DAVIS JUNIOR, L. R.; PATTERSON, R. S. A survey of urban pests ants (Hymenoptera: Formicidae) of Peninsular Florida. **Florida Entomologist**, v. 78, n. 1, p. 109-118, 1995.
- LISE, F.; GARCIA, F. R. M.; LUTINSKI, J. A. Association of ants (Hymenoptera: Formicidae) with bacteria in hospitals in State of Santa Catarina. **Revista da Sociedade Brasileira de Medicina Tropical**, v. 39, n. 6, p. 523-526, 2006.
- MOREIRA, D. D. O.; MORAES, V.; MOTTA, O. V.; CAMPOS, A. E. C.; TONHASCA JÚNIOR, R. A. Ants as potential bacteria carriers in three hospitals in Campos dos Goytacazes, Rio de Janeiro State, Brazil. **Neotropical Entomology**, v. 34, n. 6, p. 999-1006, 2005.
- NA, J. P. S.; LEE, C. Y. Identification key to commonurban pest ants in Malaysia. **Tropical Biomedicine**, v. 18, n. 1, p. 1-17, 2001.
- NELDER, M. P.; PAYSEN, E. S.; ZUNGOLI, P. A.; BENSON, E. P. Emergence of the introduced ant *Pachycondyla chinensis* (Formicidae: Ponerinae) as a public health threat in the southeastern United States. **Journal of Medical Entomology**, v. 43, n. 5, p. 1094-1098, 2006.
- OLAYA-MASMELA, L. A.; ULLOA, P. C.; PAYAN, A. Ants (Hymenoptera: Formicidae) in hospital centers of Valle del Cauca as vectors of nosocomial pathogens. **Revista Colombiana de Entomologia**, v. 31, n. 2, p. 183-187, 2005.
- PANTOJA, L. D. M.; MOREIRA FILHO, R. E.; BRITO, E. H. S.; ARAGÃO, T. B.; BRILHANTE, R. S. N.; CORDEIRO, R. A.; ROCHA, M. F. G.; MONTEIRO, A. J.; QUINET, Y. P.; SIDRIM, J. J. C. Formigas (Hymenoptera: Formicidae) como portadoras de fungos no Hospital Ambientes: ênfase sobre os gêneros *Tapinoma* e *Pheidole*. **Journal of Medical Entomology**, v. 46, n. 4, p. 895-899, 2009.
- PEREIRA, R. S.; UENO, M. Formigas como veiculadoras de microrganismos em ambiente hospitalar. **Revista da Sociedade Brasileira de Medicina Tropical**, v. 41, n. 5, p. 492-495, 2008.
- PESQUERO, M. A.; FILHO, J. E.; CARNEIRO, L. C.; FEITOSA, S. B.; OLIVEIRA, M. A. C.; QUINTANA, R. C. Formigas em ambiente hospitalar e seu potencial como transmissoras de bactérias. **Neotropical Entomology**, v. 37, n. 4, p. 472-477, 2008.
- RANDO, J. S. S.; MATSUMOTO, L. S.; SILVA, G. V.; QUIRINO, A. F. A.; HADDAD, R. E. Caracterização da mirmecofauna em estabelecimentos ligados a área da Saúde no Município de Bandeirantes, PR. **Arquivos do Instituto Biológico**, v. 76, n. 4, p. 665-671, 2009.
- RODOVALHO, C. M.; SANTOS, A. L.; MARCOLINO, M. T.; BONETTI, A. M.; BRANDEBURGO, M. A. M. Urban ants and transportation of nosocomial bacteria. **Neotropical Entomology**, v. 36, n. 3, p. 454-458, 2007.
- SANTOS, P. F.; FONSECA, A. R.; SANCHES, N. M. Formigas (Hymenoptera: Formicidae) como vetores de bactérias em dois hospitais do município de Divinópolis, Estado de Minas Gerais. **Revista da Sociedade Brasileira de Medicina Tropical**, v. 42, n. 5, p. 565-569, 2009.
- TANAKA, I. I.; VIGGIANI, A. M. F. S.; PERSON, O. C. Bactérias veiculadas por formigas em ambiente hospitalar. **Arquivos Médicos do ABC**, v. 32, n. 2, p. 60-63, 2007.
- TEIXEIRA, M. M.; PELLI, A.; SANTOS, V. M.; REIS, M. G. Microbiota associated with tramp ants in a Brazilian University Hospital. **Neotropical Entomology**, v. 38, n. 4, p. 537-541, 2009.
- TRABULSI, L. R. **Microbiologia**. Rio de Janeiro: Atheneu, 1991.
- ULLOA, P. C. Hormigas urbanas. In: FERNÁNDEZ, F. (Ed.). **Introducción a las hormigas de la región neotropical**. Bogotá: Instituto de Investigación de Recursos Biológicos Alexander von Humboldt, 2003. p. 337-349.
- ZARZUELA, M. F. M.; RIBEIRO, M. C. C.; CAMPOS-FARINHA, A. E. C. Distribuição de formigas urbanas em um hospital da região sudeste do Brasil. **Arquivos do Instituto Biológico**, v. 69, n. 1, p. 85-87, 2002.

Received on June 22, 2010.

Accepted on August 16, 2011.

License information: This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.