



Nutrición Hospitalaria

ISSN: 0212-1611

nutricion@grupoaran.com

Sociedad Española de Nutrición  
Parenteral y Enteral  
España

Iglesia, Iris; González-Gross, Marcela; Huybrechts, Inge; de-Miguel-Etayo, Pilar; Molnar, Denes; Manios, Yannis; Widhalm, Kurt; Gottrand, Frédéric; Kafatos, Anthony; Marcos, Ascensión; de la O. Puerta, Alejandro; Leclerc, Catherine; De Henauw, Stefaan; Stehle, Peter; Kersting, Mathilde; Mouratidou, Theodora; Moreno, Luis A.

Associations between insulin resistance and three B-vitamins in European adolescents:  
the HELENA study

Nutrición Hospitalaria, vol. 34, núm. 3, 2017, pp. 568-577

Sociedad Española de Nutrición Parenteral y Enteral  
Madrid, España

Available in: <http://www.redalyc.org/articulo.oa?id=309251456010>

- How to cite
- Complete issue
- More information about this article
- Journal's homepage in redalyc.org

redalyc.org

Scientific Information System

Network of Scientific Journals from Latin America, the Caribbean, Spain and Portugal

Non-profit academic project, developed under the open access initiative



## Trabajo Original

Pediatría

### Associations between insulin resistance and three B-vitamins in European adolescents: the HELENA study

*Asociaciones entre la resistencia a la insulina y tres vitaminas del grupo B en adolescentes europeos: el estudio HELENA*

Iris Iglesia<sup>1,2</sup>, Marcela González-Gross<sup>3,4</sup>, Inge Huybrechts<sup>5,6</sup>, Pilar de-Miguel-Etayo<sup>1,4</sup>, Denes Molnar<sup>7</sup>, Yannis Manios<sup>8</sup>, Kurt Widhalm<sup>9</sup>, Frédéric Gottrand<sup>10</sup>, Anthony Kafatos<sup>11</sup>, Ascensión Marcos<sup>12</sup>, Alejandro de la O. Puerta<sup>13</sup>, Catherine Leclerc<sup>14</sup>, Stefaan De Henauw<sup>5</sup>, Peter Stehle<sup>15</sup>, Mathilde Kersting<sup>16</sup>, Theodora Mouratidou<sup>1</sup> and Luis A. Moreno<sup>1,4</sup>; on behalf of the HELENA Study Group

<sup>1</sup>GENUD (Growth, Exercise, Nutrition and Development) Research Group. Instituto Agroalimentario de Aragón (IA2). Instituto de Investigación Sanitaria Aragón (IIS Aragón). Universidad de Zaragoza. Zaragoza, Spain. <sup>2</sup>Red de Salud Materno-infantil y del Desarrollo (SAMID). Hospital Universitario Cruces. Barakaldo, Bizkaia. Spain. <sup>3</sup>ImFINE Research Group. Department of Health and Human Performance. Universidad Politécnica de Madrid. Madrid, Spain. <sup>4</sup>Centro de Investigación Biomédica en Red de Fisiopatología de la Obesidad y Nutrición (CIBEROBN). Madrid, Spain. <sup>5</sup>Department of Public Health. Ghent University. Ghent, Belgium. <sup>6</sup>International Agency for Research on Cancer (IARC). Lyon, France. <sup>7</sup>Department of Pediatrics. University of Pécs. Pécs, Hungary. <sup>8</sup>Department of Nutrition and Dietetics. Harokopio University. Athens, Greece. <sup>9</sup>Division of Clinical Nutrition and Prevention. Department of Pediatrics. Medical University of Vienna. Vienna, Austria. <sup>10</sup>University Lille 2. Lille, France. <sup>11</sup>School of Medicine. University of Crete. Crete, Greece. <sup>12</sup>Immunonutrition Research Group. Department of Metabolism and Nutrition. Institute of Food Science, Technology and Nutrition (ICTAN). Spanish National Research Council (CSIC). Madrid, Spain. <sup>13</sup>Department of Physiology. School of Medicine. University of Granada. Granada, Spain. <sup>14</sup>CREA (Council for Agricultural Research and Economics). Research Center for Food and Nutrition. Rome, Italy. <sup>15</sup>Department of Nutrition and Food Science. University of Bonn. Bonn, Germany. <sup>16</sup>Research Institute of Child Nutrition. Pediatric University Clinic. Ruhr-University Bochum. Bochum, Germany

## Abstract

**Objective:** To assess whether adolescents with high body mass index (BMI), or fat mass index (FMI), in combination with insulin resistance (assessed with the Homeostatic Model Assessment [HOMA] index), had also lower blood vitamin B<sub>6</sub>, folate and vitamin B<sub>12</sub> concentrations.

**Methods and materials:** Six hundred and fifteen adolescents from the Healthy Lifestyle in Europe by Nutrition in Adolescence (HELENA) study, with data on B-vitamins (both intakes and status), and BMI, FMI, HOMA, were selected. Intakes were assessed by two non-consecutive 24-h recalls. B-vitamins biomarkers were measured by chromatography and immunoassay. Analysis of covariance was applied to elucidate the differences in B-vitamins between combinations of groups defined according to the median of the z-scores of markers of body composition and insulin sensitivity.

**Results:** When considering energy intakes and education of the mother in the model, in females, vitamin B<sub>6</sub> intakes were higher in the high BMI/high HOMA group than in the high BMI-low HOMA group. Similarly, vitamin B<sub>6</sub> intakes were higher in the high FMI/high HOMA group than in the low FMI/low HOMA group. Plasma vitamin B<sub>12</sub> was significantly lower in males in the high FMI/high HOMA group than in the low FMI/low HOMA group, keeping also significant their trends throughout the groups, a fact that can be observed also for females (p < 0.05).

**Conclusion:** Adolescents with combined higher adiposity and higher HOMA insulin sensitivity showed lower vitamin B<sub>12</sub> plasma concentrations. These differences do not seem to be explained by dietary vitamin B<sub>12</sub> intake.

## Resumen

**Objetivo:** evaluar si los adolescentes con mayor índice de masa corporal (IMC), o de masa grasa (IMG), en combinación con la resistencia a la insulina (medida con el Modelo de Valoración Homeostática [índice HOMA]), ingieren y tienen valores más bajos de vitamina B<sub>6</sub>, folato y vitamina B<sub>12</sub>.

**Métodos y materiales:** seiscientos quince adolescentes participantes en el estudio *Healthy Lifestyle in Europe by Nutrition in Adolescence* (HELENA), con valores de ingesta y concentraciones de las vitaminas B<sub>6</sub>, folato y B<sub>12</sub>, e IMC, IMG y HOMA, fueron seleccionados. Las ingestas se midieron mediante dos recuerdos de 24 horas no consecutivos. Los biomarcadores de las vitaminas fueron obtenidos mediante cromatografía e inmunoensayo. Se aplicó el análisis de la covarianza para evaluar las diferencias entre las vitaminas (ingesta y concentraciones) entre las combinaciones de los grupos definidos según las medianas de los valores z de los marcadores de la composición corporal y de la sensibilidad a la insulina.

**Resultados:** considerando la ingesta energética y la educación de la madre en el modelo en chicas, las ingestas de vitamina B<sub>6</sub> fueron mayores en el grupo de mayor IMC/mayor HOMA que en el grupo mayor IMC/menor HOMA. Del mismo modo, el grupo constituido por mayor IMG/mayor HOMA presentó mayores ingestas de esta vitamina que el grupo formado por la combinación entre menor IMG/menor HOMA. La vitamina B<sub>12</sub> plasmática en chicos fue significativamente menor en el grupo formado por mayor IMG/mayor HOMA que en el grupo menor IMG/menor HOMA, manteniendo también la tendencia significativa en los grupos, lo que también se puede observar en las chicas (p < 0.05).

**Conclusiones:** los adolescentes con mayor adiposidad en combinación con una mayor sensibilidad a la insulina mostraron menores valores de vitamina B<sub>12</sub> plasmática. Estas diferencias no parecen estar explicadas por diferencias en la ingesta de vitamina B<sub>12</sub>.

### Key words:

Vitamin B<sub>6</sub>. Folic acid. Vitamin B<sub>12</sub>. Adolescent. Body mass index. Insulin resistance.

### Palabras clave:

Vitaminas B<sub>6</sub>. Ácido fólico. Vitamina B<sub>12</sub>. Adolescentes. Índice de masa corporal. Resistencia a la insulina.

Received: 19/09/2016  
Accepted: 30/12/2016

Iglesia I, González-Gross M, Huybrechts I, de-Miguel-Etayo P, Molnar D, Manios Y, Widhalm K, Gottrand F, Kafatos A, Marcos A, Puerta AO, Leclerc C, De Henauw S, Stehle P, Kersting M, Mouratidou T, Moreno LA; on behalf of the HELENA Study Group. Associations between insulin resistance and three B-vitamins in European adolescents: the HELENA study. Nutr Hosp 2017;34:568-577

DOI: <http://dx.doi.org/10.20960/nh.559>

### Correspondence:

Iris Iglesia. Growth, Exercise, Nutrition and Development (GENUD) Research group. Facultad de Ciencias de la Salud. Universidad de Zaragoza. C/ Pedro Cerbuna, 12. Edif. SAI, 2ª planta. 50009 Zaragoza, Spain  
e-mail: [iglesia@unizar.es](mailto:iglesia@unizar.es)

## INTRODUCTION

Prevalence of overweight and obesity in European children ranges between 10 and 40 percent among European adolescents (1), while obesity is currently considered as the fifth leading risk for global deaths (2). Aside genetics, inadequate lifestyle factors like unhealthy dietary habits and/or insufficient physical activity are the main attributable causes of both overweight and obesity (3). Childhood obesity has been shown to be accompanied by low micronutrient intake and micronutrient deficiencies (4). For instance, obesity has been related to low iron intake in children and adolescents in a study developed in Israel (5), where obese children and adolescents showed a higher prevalence of iron deficiency or even iron deficiency anemia than non-obese.

The body mass index (BMI) is the most widely used height-normalized index for the screening of excess body fat, also in adolescents, but it can be criticized since it does not discriminate between lean- and fat-mass (6). Measurement of skinfolds thickness allows an estimation of subcutaneous adipose tissue deposition and, thus, the use of the fat mass index (FMI = kg fat mass/m<sup>2</sup>) instead of the BMI for classifying obesity status in children (6). Obesity is often associated with hyperinsulinism and insulin resistance, which over time can develop into glucose intolerance, impaired  $\beta$ -cell function and diabetes mellitus type 2 (7).

Vitamin B<sub>12</sub> is a crucial nutrient present in animal products (8). Its main roles are linked with the cognitive function, bone health, and deoxyribonucleic acid (DNA)-replication during periods of rapid growth and development like childhood and adolescence (9,10). An optimal vitamin B<sub>12</sub> status during early life stages is essential in preventing future health risks like anemia (9). Besides, vitamin B<sub>12</sub> deficiency contributes to hyperhomocysteinemia, which is an independent risk factor for atherosclerotic disease (11). Sub-clinical deficiencies of vitamin B<sub>12</sub> status are not uncommon during adolescence and in high risk population groups like vegans or vegetarians, elderly or low-resource people (12).

Low vitamin B<sub>12</sub> status can be due to gut malabsorption syndromes, pernicious anemia (13), or secondary malabsorption produced by metformin therapy, an insulin sensitizer used for the treatment of type 2 diabetes and insulin resistance in adolescents (14,15).

A recent paper based on the HELENA study reported levels of B-related vitamins, such as B<sub>6</sub>, folate and B<sub>12</sub> (16), in which 2% of studied adolescents had low plasma vitamin B<sub>12</sub> and 5% had low holotranscobalamin (HoloTC) concentrations. Besides, low concentrations of both plasma folate (PF) and red blood cells (RBC-folate) were identified in 10% of the HELENA adolescents, and low pyridoxal-phosphate (PLP) concentrations were also identified in 5% of them.

A recent Australian study with obese adolescents (13) called for investigation of the associations between vitamin B<sub>12</sub> status and insulin sensitivity, including also dietary intakes. Consequently, this study aims to assess whether adolescents with higher body mass index (BMI) or fat mass index (FMI), in combination with higher insulin sensitivity (high Homeostatic Model Assessment [HOMA] index) had also lower B-vitamins concentrations.

To our knowledge, this is the first study in European adolescents assessing the association between B-vitamins intake and concentrations and insulin sensitivity, considering indicators of body composition like BMI or FMI.

## METHODS AND MATERIALS

The multicenter and cross-sectional study Healthy Lifestyle in Europe by Nutrition in Adolescence (HELENA-CSS) recruited adolescents aged 12.5-17.5 years, from ten cities from nine European countries: Athens and Heraklion, in Greece; Dortmund, in Germany; Ghent, in Belgium; Lille, in France; Pécs, in Hungary; Rome, in Italy; Stockholm, in Sweden; Vienna, in Austria; and Zaragoza, in Spain. The purpose of the study was to provide complete and reliable information about the nutritional status of European adolescents (17). Inclusion criteria were: not participating simultaneously in another clinical trial and being free of any acute infection occurring < one week before inclusion (18). The total number of participants was 3,528, with an average participation rate of 67%, which can be considered as acceptable for such a demanding epidemiological study (19). In one third of the sample in each study center (1,076 adolescents), blood drawing was obtained. Participants from Heraklion and Pécs (7% of the total sample) did not provide comparable dietary data. For the purposes of this analysis, 615 adolescents were included, having complete data on BMI, skinfold thickness to calculate FMI, maternal education, vitamin B<sub>6</sub>, folate and vitamin B<sub>12</sub> intakes and biomarkers, and having the HOMA index for insulin sensitivity or resistance excluding also outliers (for biochemical measurements, outliers were considered when values were  $\pm$  four standard deviations from the mean). Further details on the HELENA sampling procedures, pilot study and reliability of the data have been published elsewhere (19). Informed consent was obtained from all participants and their parents, and the protocol was approved by the Human Research Review Committees of the corresponding centers (20).

## ASSESSMENT OF VITAMIN B<sub>6</sub>, FOLATE, VITAMIN B<sub>12</sub> AND ENERGY INTAKES

Vitamin and energy intakes were assessed using the computerized 24-hour recall, self-administered HELENA-Dietary Assessment Tool (HELENA-DIAT), adapted for European adolescents from the Young Adolescents' Nutrition Assessment on Computer (YANA-C) software (21). The adolescents completed the 24-hour recalls twice in a fortnight period. Trained staff were present during completion (21). Obtained data was linked to the German Food Code and Nutrient Data Base (BLS [Bundeslebensmittelschlüssel], version II.3.1, 2005), with 12,000 coded foods, and with up to 158 nutrient data points available for each food item (21). When traditional or local foods were not available in the BLS table, recipes were composed using foods from the BLS as ingredients. The Multiple Source Method (MSM) (22) was applied to calculate usual nutrient intakes removing the effect of day-to-

day within-person variation and random error in the two recalls. B-vitamins diet densities were calculated as follows: (amount of B-vitamin intake per 1,000 kcal of diet/recommendation of the corresponding B-vitamin intake based on the Institute of Medicine recommendations) \*100. Recommendations for vitamin B<sub>6</sub> intakes are 1,300 µg in males and 1,200 µg in females; for folate, 400 µg in males and females; and for vitamin B<sub>12</sub> 2.4 µg in both sexes (9).

## ASSESSMENT OF VITAMIN B<sub>6</sub>, FOLATE AND VITAMIN B<sub>12</sub> BIOMARKERS CONCENTRATIONS

In schools, early in the morning, and in fasting status, 30 ml of blood were drawn according to a standardized blood collection protocol by a certified phlebotomist. More details on sample transport and quality assurance can be found elsewhere (23). For the measurement of pyridoxal 5'phosphate (PLP), biomarker of vitamin B<sub>6</sub>, ethylene diamine tetraacetic acid (EDTA) whole blood was centrifuged at 3,500 g for 15 min. The supernatants were stored at -80 °C until analyzed. PLP was measured by high performance liquid chromatography (HPLC) (Varian Deutschland GmbH, Darmstadt, Germany; coefficient of variation [CV] = 1%) with a modified method of Kimura et al. (16,24).

For the measurement of plasma folate and plasma vitamin B<sub>12</sub>, heparinized tubes were collected, placed immediately on ice, and centrifuged within 30 min (3,500 g for 15 min). The supernatant fluid was transported at a stable temperature of 4-7 °C to the central laboratory at the University of Bonn (IEL-Institut fuer Ernährungs [und Lebensmittelwissenschaften], Germany) and stored at 80 °C until assayed. After measuring hematocrit *in situ*, EDTA whole blood was used for the red blood cell folate (RBC-folate) analysis. EDTA whole blood was diluted 1:5 with freshly prepared 0.1% ascorbic acid for cell lysis and incubated for 60 min in the dark before storage at 80 °C. Plasma and RBC-folate and plasma vitamin B<sub>12</sub> were measured by means of a competitive immunoassay using the Immunolite 2000 analyzer (DPC Biermann GmbH, Bad Nauheim, Germany) (CV for plasma folate = 5.4%, RBC folate = 10.7%, cobalamin = 5.0%) (23). Sera for measuring holotranscobalamin (HoloTC) were obtained by centrifuging blood collected in evacuated tubes without anticoagulant at 3,500 g for 15 min within one hour. Once sent to the central laboratory, sera were aliquoted and stored at 80 °C until transport in dry ice to the biochemical lab at the Universidad Politécnica de Madrid for analysis (laboratory number 242 of the Laboratory Network of the Region of Madrid). HoloTC was measured by microparticle enzyme immunoassay (Active B<sub>12</sub>, Axis-Shield Ltd., Dundee, Scotland, UK) with the use of AxSym (Abbot Diagnostics, Abbott Park, IL, USA) (CV = 5.1%) (25).

Glucose was measured using enzymatic methods (Dade Behring, Schwalbach, Germany). Insulin levels were measured using an Immulite 200 analyzer (DPC Bierman GmbH, Bad Nauheim, Germany). The homeostasis model assessment (HOMA) calculation was used as a measurement of insulin resistance (glycaemia X insulin/22.5) (26).

## CONFOUNDERS

Maternal education was used as proxy of socioeconomic status, obtained via self-administered questionnaire completed by the adolescents and expressed as: elementary, lower secondary, higher secondary or tertiary education. This variable was one of the most related socioeconomic factors associated with the studied vitamins (27). Total energy intake in kcal/d assessed with the 24 hours' dietary recalls software HELENA-DIAT was also used as a covariate in the analyses.

## ANTHROPOMETRY

Anthropometry battery measurements were assessed following standardized and strictly controlled procedures previously described (28). Weight was measured in underwear and without shoes with an electronic scale (Type SECA 861) to the nearest 0.05 kg, and height was measured barefoot in the Frankfort plane with a telescopic height measuring instrument (Type SECA 225) to the nearest 0.1 cm. Body weight and height, together with subscapular and tricipital skinfold thicknesses, were measured in triplicate. BMI was calculated using the Quetelet formula (kg/m<sup>2</sup>). The body fat percentage was calculated using the Slaughter's equation (29), and thereafter the fat mass index (FMI) was calculated by dividing fat mass by height squared (m<sup>2</sup>).

## STATISTICAL ANALYSIS

The Statistical Package for Social Sciences version 20.0 (SPSS Inc., Chicago, IL, USA) was used to analyze the data. All analyses were sex-specific. Descriptive data are presented as means and standard deviations (SD). Z-scores of BMI and FMI, and HOMA considering age and sex were used in the analyses. The Mann-Whitney test for non-parametric variables was applied to examine differences between two predefined groups, i.e., above or under the median of the z-scores for BMI, FMI, and HOMA. Thereafter, obtained categories were combined to create four subsequent groups to analyze intakes and statuses of the vitamins: low BMI (or FMI) - low HOMA (the most favorable group in terms of body composition and insulin sensitivity); low BMI (or FMI) - high HOMA; high BMI (or FMI) - low HOMA; and, finally, high BMI (or FMI) - high HOMA (the less favorable group). The Kruskal-Wallis test was applied to look at the differences in B-vitamin intakes and biomarkers among groups, with a Mann-Whitney test approach to contrast values 2 x 2. Finally, maternal education and total energy intakes were included in the analysis of covariance (ANCOVA) as covariates. All statistical tests and corresponding p values were two-sided, and p < 0.05 was the cut-off to consider a result as statistically significant.

## RESULTS

In the whole HELENA sample males represented 45%, and this percentage is significantly lower than in our analysis, where males

represented the 47% ( $p = 0.004$ ). In terms of maternal education, included males and females showed significantly higher maternal education levels than those who were excluded (data not shown) ( $p < 0.001$ ). Males included in the dietary analysis had significantly lower energy intake ( $p = 0.001$ ) than those excluded; no significant differences were observed in females. There was no statistically significant difference in biomarkers of all three vitamins between the samples included and excluded (based on the inclusion criteria and required data availability).

Table I presents the main characteristics of the sample, by sex. FMI and HOMA were significantly higher in females than in males whilst total energy intake was lower ( $p < 0.05$ ). Table II presents adolescents' B-vitamins intakes and biomarkers by the corresponding groups based on the median z-scores of variables of interest (BMI, FMI, and HOMA). Adolescents with low BMI, HOMA and FMI z-scores consumed more B-vitamins than those with high BMI, HOMA and FMI z-scores, except in the case of males with low HOMA, who had lower folate intakes. Considering biomarkers, adolescents with lower BMI, FMI and HOMA showed higher B-vitamins levels, except in the case of males belonging to the low HOMA group, who had lower RBC-folate concentrations than their counterparts in the higher HOMA group ( $p < 0.05$ ).

Table III shows the differences in B-vitamins both in intakes and concentrations among the groups defined by the subsequent categories of  $<$  median and  $>$  median of BMI, FMI, and HOMA z-scores considered in combination, without any additional adjustment. All in all, results showed that adolescents belonging to the less favorable groups (high BMI-or FMI-/high HOMA) had lower B-vitamins intake and concentrations, and also lower total energy intakes. However, when an analysis of the covariance was performed to introduce in the model the covariates of education of the mother and total energy intake, most of the statistically significant differences among groups disappeared.

Table IV shows similarly, the adjusted results of the table II (by energy intakes, and maternal education). In females, vitamin B<sub>6</sub> intakes were higher in the high BMI/high HOMA group than in the high BMI/low HOMA group ( $p < 0.05$ ). Similarly, the high FMI/high HOMA group vitamin B<sub>6</sub> intakes were higher than in the low FMI/low HOMA group. Plasma vitamin B<sub>12</sub> in males were significantly lower in the high FMI/high HOMA than in the low FMI/low HOMA group, keeping also significant their trends throughout the groups, which can be observed also for females ( $p < 0.05$ ). These trends can be followed in figure 1.

## DISCUSSION

To the authors' knowledge, this is the first study to examine the association between B-vitamin intakes and the corresponding biomarker concentrations and insulin resistance according to markers of body composition such as body mass index and fat mass index in European adolescents. The results suggest an association between higher adiposity together with higher insulin sensitivity and plasma vitamin B<sub>12</sub> concentrations, showing the lowest vitamin B<sub>12</sub> plasma concentrations in those adolescents with higher levels of adiposity combined with higher HOMA insulin sensitivity. Three B-vitamins intake and status corresponding to vitamin B<sub>6</sub>, folate and vitamin B<sub>12</sub> have been investigated in this study, and we have obtained several differences in them in different groups constituted by the combinations between BMI, FMI and HOMA categories. However, intake of vitamin B<sub>6</sub> and plasma vitamin B<sub>12</sub> was the only for which a difference was found for categories of the mentioned groups when education of the mother and energy intake was taken into account.

The prevalence of insufficient vitamin B<sub>12</sub> biomarker levels in this sample of European adolescents is low (5% based on HoloTC

**Table I. Characteristics of the participants by sex**

Characteristics	Males (n = 281), mean $\pm$ SD	Females (n = 334), mean $\pm$ SD	p-values
Age*	14.8 $\pm$ 1.3	14.8 $\pm$ 1.2	0.71
BMI (kg/m <sup>2</sup> )	20.9 $\pm$ 3.6	21.2 $\pm$ 3.3	0.12
FMI (kg fat mass/m <sup>2</sup> )	12.6 $\pm$ 9.2	15.0 $\pm$ 6.8	0.03
HOMA	2.0 $\pm$ 1.2	2.2 $\pm$ 1.2	$< 0.00$
Energy intake (kcal/d)*	2,523 $\pm$ 816	1,892 $\pm$ 544	$< 0.00$
Vitamin B <sub>6</sub> diet density (%)*	57.2 $\pm$ 14.8	64.8 $\pm$ 16.3	$< 0.00$
Folate diet density (%)*	21.8 $\pm$ 5.8	24.4 $\pm$ 6.5	$< 0.00$
Vitamin B <sub>12</sub> diet density (%)*	104.4 $\pm$ 33.5	104.0 $\pm$ 38.0	0.43
ME: Low education (n, %)	17 (6)	29 (9)	0.17
ME: Medium-low education (n, %)	60 (21)	83 (25)	
ME: Medium-high education (n, %)	92 (33)	115 (34)	
ME: High education (n, %)	112 (40)	107 (32)	

BMI: Body mass index; FMI: Fat mass index; HOMA: Homeostasis Model Assessment; ME: Maternal education. B-vitamins diet densities were calculated as follows: (amount of B-vitamin intake per 1,000 kcal of diet/recommendation of the corresponding B-vitamin intake based on the Institute of Medicine recommendations) \*100. Recommendations for vitamin B<sub>6</sub> intakes are 1,300  $\mu$ g for males and 1,200  $\mu$ g for females; for folate, 400  $\mu$ g in males and females; and for vitamin B<sub>12</sub>, 2.4  $\mu$ g in both sexes. p-values in bold are the only significant ones, based either in Mann-Whitney test or t-test\* at 0.05 level two-sided.



**Table II.** B-vitamin intakes and concentrations and the groups established by the medians of the sex-specific z-scores of body mass index, fat mass index and Homeostasis Model Assessment index stratified by sex

B-vitamins	Males			Females		
	< median of BMI z-score <sup>^</sup> (n = 140)	> median of BMI z-score <sup>^</sup> (n = 141)		< median of BMI z-score <sup>^</sup> (n = 167)	> median of BMI z-score <sup>^</sup> (n = 167)	
	Mean ± SD	Mean ± SD	p-value	Mean ± SD	Mean ± SD	p-value
<i>Intakes</i>						
Vitamin B <sub>6</sub> (µg/d)	1,925 ± 608	1,725 ± 577	0.01	1,508 ± 511	1,385 ± 446	0.02
Vitamin B <sub>6</sub> diet density (%)	56.0 ± 13.5	58.4 ± 16.0	0.25	61.8 ± 15.3	67.9 ± 16.8	0.00
Folate (µg/d)	222 ± 73.2	206 ± 69.5	0.11	189 ± 57.2	172 ± 56.2	0.01
Folate diet density (%)	21.0 ± 4.8	22.7 ± 6.6	0.04	23.5 ± 5.6	25.4 ± 7.2	0.01
Vitamin B <sub>12</sub> (µg/d)	6.6 ± 2.4	5.8 ± 2.5	0.00	4.8 ± 1.6	4.4 ± 1.7	0.00
Vitamin B <sub>12</sub> diet density (%)	104.5 ± 32.6	104.4 ± 34.5	0.92	99.9 ± 30.9	108.1 ± 43.7	0.13
<i>Status</i>						
PLP (pmol/l)	67.3 ± 42.0	69.3 ± 47.4	0.75	63.7 ± 80.3	62.0 ± 44.4	0.58
PF (nmol/l)	18.6 ± 9.4	18.2 ± 11.4	0.29	18.8 ± 9.3	17.8 ± 10.1	0.09
RBC-folate (nmol/l)	818 ± 328	827 ± 433	0.47	772 ± 305	761 ± 311	0.67
Plasma B <sub>12</sub> (pmol/l)	358 ± 143	318 ± 115	0.04	403 ± 154	355 ± 159	0.00
HoloTC (pmol/l)	66.1 ± 36.9	64.5 ± 27.4	0.64	64.8 ± 36.1	65.7 ± 36.6	0.82
	< median of FMI z-score (n = 140)	> median of FMI z-score (n = 141)		< median of FMI z-score (n = 167)	> median of FMI z-score (n = 167)	
<i>Intakes</i>						
Vitamin B <sub>6</sub> (µg/d)	1,910 ± 606	1,740 ± 585	0.01	1,466 ± 458	1,426 ± 506	0.31
Vitamin B <sub>6</sub> diet density (%)	54.8 ± 12.9	59.6 ± 16.3	0.03	62.1 ± 15.7	67.6 ± 16.5	0.02
Folate (µg/d)	222 ± 72.8	206 ± 70.0	0.07	185 ± 56.0	176 ± 57.7	0.10
Folate diet density (%)	20.8 ± 4.9	22.9 ± 6.4	0.01	23.6 ± 4.9	25.3 ± 7.1	0.02
Vitamin B <sub>12</sub> (µg/d)	6.7 ± 2.5	5.7 ± 2.4	< 0.00	4.7 ± 1.5	4.5 ± 1.8	0.15
Vitamin B <sub>12</sub> diet density (%)	104.4 ± 33.6	104.4 ± 33.6	0.95	104.4 ± 33.6	104.4 ± 33.6	0.13
<i>Status</i>						
PLP (pmol/l)	65.5 ± 41.7	71.1 ± 47.3	0.24	67.1 ± 81.4	58.4 ± 41.0	0.60
PF (nmol/l)	18.5 ± 9.5	18.3 ± 11.2	0.49	19.1 ± 10.1	17.5 ± 9.3	0.07
RBC-folate (nmol/l)	825 ± 335	820 ± 428	0.40	787 ± 311	746 ± 304	0.14
Plasma B <sub>12</sub> (pmol/l)	362 ± 144	313 ± 113	0.01	403 ± 165	355 ± 147	0.01
HoloTC (pmol/l)	67.6 ± 38.4	63.0 ± 24.7	0.76	66.2 ± 38.3	64.4 ± 34.3	0.99
	< median of HOMA z-score (n = 140)	> median of HOMA z-score (n = 141)		< median of HOMA z-score (n = 167)	> median of HOMA z-score (n = 167)	
<i>Intakes</i>						
Vitamin B <sub>6</sub> (µg/d)	1,868 ± 620	1,781 ± 579	0.20	1,431 ± 459	1,461 ± 506	0.88
Vitamin B <sub>6</sub> diet density (%)	56.2 ± 14.2	58.1 ± 15.4	0.29	62.5 ± 15.7	67.1 ± 16.6	0.01
Folate (µg/d)	212 ± 70.4	216 ± 73.2	0.84	184 ± 57.4	178 ± 57.0	0.41
Folate diet density (%)	20.8 ± 5.1	22.9 ± 6.3	0.00	24.2 ± 6.6	24.7 ± 6.5	0.42

(Continuation in the next page)

**Table II (Cont.).** B-vitamin intakes and concentrations and the groups established by the medians of the sex-specific z-scores of body mass index, fat mass index and Homeostasis Model Assessment index stratified by sex

B-vitamins	Males			Females		
	< median of HOMA z-score (n = 140)	> median of HOMA z-score (n = 141)		< median of HOMA z-score (n = 167)	> median of HOMA z-score (n = 167)	
<i>Intakes</i>						
Vitamin B <sub>12</sub> (µg/d)	6.5 ± 2.7	5.9 ± 2.2	0.04	4.6 ± 1.7	4.5 ± 1.7	0.65
Vitamin B <sub>12</sub> diet density (%)	105.5 ± 35.0	103.3 ± 32.1	0.64	101.5 ± 37.4	106.4 ± 38.6	0.18
<i>Status</i>						
PLP (pmol/l)	67.4 ± 42.0	69.1 ± 47.3	1.0	67.5 ± 79.1	58.2 ± 46.3	0.24
PF (nmol/l)	18.9 ± 10.7	17.9 ± 10.1	0.43	19.6 ± 10.4	17.0 ± 8.9	0.00
RBC-folate (nmol/l)	815 ± 395	829 ± 373	0.74	766 ± 285	766 ± 329	0.70
Plasma B <sub>12</sub> (pmol/l)	354 ± 132	322 ± 129	0.03	397 ± 160	361 ± 154	0.03
HoloTC (pmol/l)	70.1 ± 40.4	60.5 ± 20.6	0.17	67.8 ± 40.1	62.8 ± 32.1	0.33

BMI: Body mass index; FMI: Fat mass index; HOMA: Homeostasis Model Assessment; PLP: Pyridoxal phosphate; PF: Plasma folate; RBC-folate: Red blood cell folate; HoloTC: Holotranscobalamin; SD: Standard deviation. B-vitamins diet densities were calculated as follows: (amount of B-vitamin intake per 100 kcal of diet/ recommendation of the corresponding B-vitamin intake based on the Institute of Medicine recommendations) \*100. Recommendations for vitamin B<sub>6</sub> intakes are 1,300 µg for males and 1,200 µg for females; for folate, 400 µg in males and females; and for vitamin B12, 2.4 µg in both sexes. p-value based on Mann-Whitney non-parametric test. Statistical significance at 0.05 two-sided level. ^Sex-specific z-scores of BMI, FMI and HOMA were adjusted by age.

**Table III.** Differences of B-vitamin intakes and biomarkers concentrations by resulting combination groups between z-scores of body mass index and fat mass index against z-scores of insulin resistance (HOMA) index

Males	Energy intake (mean, SD)	Vitamin B <sub>6</sub> (mean, SD)	Folate (mean, SD)	Vitamin B <sub>12</sub> (mean, SD)	PLP (mean, SD)	PF (mean, SD)	RBC- folate (mean, SD)	Plasma B <sub>12</sub> (mean, SD)	HoloTC (mean, SD)
<i>BMI by HOMA (n)</i>									
Low BMI-low HOMA (91)	2,659 ± 783 <sup>a</sup>	1,910 ± 618 <sup>a</sup>	215 ± 71	6.7 ± 2.4 <sup>ab</sup>	65.7 ± 40.8	18.3 ± 8.7	782 ± 300	366 ± 142 <sup>a</sup>	70.3 ± 43.5
Low BMI- high HOMA (49)	2,763 ± 893	1,953 ± 596	236 ± 77.0	6.3 ± 2.4	70.3 ± 44.4	19.1 ± 10.5	882 ± 370	342 ± 147	58.5 ± 17.6
High BMI-low HOMA (49)	2,502 ± 800	1,791 ± 552	208 ± 70.4	6.1 ± 3.1 <sup>a</sup>	70.8 ± 44.6	20.0 ± 13.7	877 ± 525	330 ± 111	69.7 ± 34.6
High BMI-high HOMA (92)	2,273 ± 757 <sup>a</sup>	1,689 ± 552 <sup>a</sup>	206 ± 69.5	5.6 ± 2.1 <sup>b</sup>	68.5 ± 49.0	17.3 ± 9.9	801 ± 373	311 ± 117 <sup>a</sup>	61.6 ± 22.1
p-value	< 0.00	0.03	0.19	0.01	0.97	0.64	0.55	0.08	0.44
<i>FMI by HOMA (n)</i>									
Low-FMI-low HOMA (86)	2,660 ± 770 <sup>a</sup>	1,903 ± 628 <sup>a</sup>	215 ± 69.1	6.8 ± 2.5 <sup>ab</sup>	65.5 ± 41.1	18.5 ± 9.1	778 ± 296	366 ± 141 <sup>a</sup>	71.5 ± 44.7
Low FMI-high HOMA (54)	2,843 ± 952	1,921 ± 569	235 ± 78.1	6.4 ± 2.4	65.6 ± 43.5	18.6 ± 10.4	908 ± 385	357 ± 149.0	60.4 ± 20.7
High FMI-low HOMA (54)	2,504 ± 822	1,805 ± 605	208 ± 73.1	6.0 ± 2.9 <sup>a</sup>	71.1 ± 44.0	19.7 ± 13.2	882 ± 525	332 ± 113	67.5 ± 30.7
High FMI-high HOMA (87)	2,223 ± 677 <sup>a</sup>	1,704 ± 574 <sup>a</sup>	206 ± 68.6	5.6 ± 2.0 <sup>b</sup>	71.0 ± 49.3	17.5 ± 9.9	785 ± 361	302 ± 112 <sup>a</sup>	60.6 ± 20.7
p-value	< 0.00	0.06	0.17	0.00	0.68	0.83	0.21	0.02	0.57

(Continuation in the next page)

**Table III (Cont.).** Differences of B-vitamin intakes and biomarkers concentrations by resulting combination groups between z-scores of body mass index and fat mass index against z-scores of insulin resistance (HOMA) index

Females	Energy intake (mean, SD)	Vitamin B <sub>6</sub> (mean, SD)	Folate (mean, SD)	Vitamin B <sub>12</sub> (mean, SD)	PLP (mean, SD)	PF (mean, SD)	RBC-folate (mean, SD)	Plasma B <sub>12</sub> (mean, SD)	HoloTC (mean, SD)
<i>BMI by HOMA (n)</i>									
Low BMI-low HOMA (100)	2,045 ± 483 <sup>ab</sup>	1,511 ± 488 <sup>a</sup>	194 ± 52.4 <sup>ab</sup>	4.8 ± 1.6 <sup>ab</sup>	64.7 ± 94.4 <sup>a</sup>	19.7 ± 9.3 <sup>ab</sup>	779 ± 290	418 ± 149 <sup>ab</sup>	67.2 ± 33.7
Low BMI-high HOMA (67)	2,043 ± 544	1,502 ± 546	181 ± 63.4	4.8 ± 1.5	62.2 ± 54.4	17.5 ± 9.2 <sup>a</sup>	760 ± 329	380 ± 159	61.3 ± 39.5
High BMI-low HOMA (67)	1,762 ± 449 <sup>a</sup>	1,312 ± 386 <sup>a</sup>	168 ± 61.5 <sup>a</sup>	4.3 ± 1.7 <sup>a</sup>	71.6 ± 49.1 <sup>a</sup>	19.4 ± 11.9	747 ± 280	366 ± 172 <sup>a</sup>	68.8 ± 48.3
High BMI-high HOMA (100)	1,727 ± 595 <sup>b</sup>	1,433 ± 478	175 ± 52.6 <sup>b</sup>	4.4 ± 1.7 <sup>b</sup>	55.5 ± 39.9	16.7 ± 8.7 <sup>b</sup>	770 ± 331	348 ± 151 <sup>b</sup>	63.7 ± 26.6
p-value	< 0.00	0.05	0.01	0.02	0.04	0.02	0.95	0.00	0.47
<i>FMI by HOMA (n)</i>									
Low FMI-low HOMA (102)	1,995 ± 437 <sup>ab</sup>	1,466 ± 417	189 ± 51.1	4.7 ± 1.5	70.0 ± 94.7	19.9 ± 10.4 <sup>a</sup>	790.7 ± 294	421 ± 159 <sup>abc</sup>	67.6 ± 35.4
Low FMI-high HOMA (65)	1,995 ± 554	1,467 ± 520	180 ± 64.1	4.7 ± 1.5	63.4 ± 56.1	17.8 ± 9.7	780 ± 339	374 ± 157 <sup>a</sup>	63.9 ± 42.6
High FMI-low HOMA (65)	1,832 ± 550 <sup>a</sup>	1,377 ± 517	175.4 ± 65.7	4.5 ± 1.9	64.2 ± 44.6	19.0 ± 10.4	728 ± 268	359 ± 156 <sup>b</sup>	68.1 ± 46.8
High FMI-high HOMA (102)	1,764 ± 604 <sup>b</sup>	1,457 ± 499	176 ± 52.3	4.5 ± 1.7	54.8 ± 38.3	16.6 ± 8.4 <sup>a</sup>	757 ± 324	353 ± 142 <sup>c</sup>	62.0 ± 23.4
p-value	< 0.00	0.64	0.24	0.54	0.35	0.02	0.51	0.01	0.57

BMI: Body mass index; HOMA: Homeostasis Model Assessment; FMI: Fat mass index; PLP: Pyridoxal phosphate; PF: Plasma folate; RBC-folate: Red blood cell folate; HoloTC: Holotranscobalamin; SD: Standard deviation. p-value based on Kruskal-Wallis non-parametric test. Statistical significance was established at 0.05 two-sided level. Superscripts letters represent those groups with statistical significant differences. Post hoc comparisons between groups have been established based on Mann-Whitney test. Statistical significance critical value established at 0.0167 two-sided level (resulting from dividing 0.05/3 comparisons with the reference category of "low BMI (or FMI)-low HOMA" which represents the more favorable option).

and 2% based on serum B<sub>12</sub>) (30), and corresponds approximately to the prevalence of inadequate vitamin B<sub>12</sub> intakes reported previously (2.9% in males and 6.0% in females) (8) in a larger sample of the same adolescents. In other studies performed in Australia and Canada, the percentage of adolescents identified with low or borderline B<sub>12</sub> status was higher than the one reported in our European adolescents (respectively, 32.1% in obese adolescents in the study from Australia (13), or 13.7% in all children and adolescents and 20.4% only in obese children and adolescents reported in the Canadian survey (31) using the same cut-points as in the Australian study). However, these percentages of low vitamin B<sub>12</sub> status refer exclusively to overweight/obese adolescents, and this could be the cause of such big differences with ours.

In our sample, 23.3% (32) of the adolescents were classified as overweight or obese. Even with these differences in rates of overweight/obesity and deficiency or not of vitamin B<sub>12</sub> between our study and the Australian study (13), our results confirm the association between lower levels of vitamin B<sub>12</sub> and higher HOMA

and FMI. Besides, this negative association between vitamin B<sub>12</sub> plasma levels and values of insulin sensitivity and body composition markers are likely reproducible for other micronutrients (total carotenoids, alpha-carotene, beta-carotene, beta-cryptoxanthin, lutein/zeaxanthin, lycopene, vitamin E, vitamin C, selenium, vitamin A, vitamin D, folate, vitamin B<sub>12</sub>, and RBC-folate), as reported in a study of US adults (33).

There are no previous studies investigating the same associations in a similar population group. Nevertheless, previous studies (4,13,31) focused on obese adolescents or wide general population groups have also found that higher BMI z-scores were associated with lower vitamin B<sub>12</sub> concentrations, with no significant gender effect.


There are several mechanisms which might explain this observation. For instance, adolescents have increased nutrient requirements secondary to increased growth and body size (34). Another reason is that adolescents with high BMI are thought to have diets with low micronutrient density. In our study, we found



**Table IV.** Adjusted estimates of B-vitamin intakes and biomarker status by combination of body mass index, fat mass index and Homeostasis Model Assessment index

Indicators	Vitamin B <sub>12</sub> intakes (µg/d)				Plasma B <sub>12</sub> (pmol/l)				Holotranscobalamin (pmol/l)													
	Males		Females		Males		Females		Males		Females											
	Mean	SE	Mean	SE	Mean	SE	Mean	SE	Mean	SE	Mean	SE										
BMI by HOMA (n)																						
Low BMI-low HOMA (90, 100)	6.5	0.2	4.5	0.1	364	13.7	416	15.7	70.1	3.5	65.1	3.8										
Low BMI-high HOMA (50, 67)	5.9	0.3	4.6	0.2	339	18.6	373	19.2	57.9	4.8	60.2	4.6										
High BMI-low HOMA (50, 67)	6.1	0.3	4.4	0.2	330	18.6	352	19.1	69.2	4.7	69.0	4.5										
High BMI-high HOMA (91, 100)	6.1	0.2	4.7	0.1	315	13.7	362	15.9	62.5	3.6	66.3	3.7										
F and p-values	1.01	0.39	0.66	0.58	2.17	0.09	1.78	0.15	1.85	0.14	0.64	0.59										
FMI by HOMA (n)																						
Low-FMI-low HOMA (86, 101)	6.5	0.2	4.5	0.1	361 <sup>a</sup>	13.8	426	15.3	70.5	3.5	66.5	3.7										
Low FMI-high HOMA (54, 66)	5.9	0.3	4.5	0.2	355	8.5	352	19.0	59.3	4.8	63.5	4.6										
High FMI-low HOMA (54, 66)	6.1	0.3	4.5	0.2	338	18.4	361	19.1	68.4	4.9	67.2	4.6										
High FMI-high HOMA (87, 101)	6.1	0.2	4.8	0.1	306 <sup>a</sup>	13.8	360	15.5	61.7	3.6	64.0	3.7										
F and p-values	1.18	0.32	0.71	0.55	2.89	0.04	2.94	0.03	1.73	0.16	0.18	0.91										
Indicators	Vitamin B <sub>6</sub> intakes (µg/d)				Pyridoxal phosphate (pmol/l)				Folate intakes (µg/d)				Plasma folate (pmol/l)				RBC-folate (pmol/l)					
	Males		Females		Males		Females		Males		Females		Males		Females		Males		Females			
	Mean	SE	Mean	SE	Mean	SE	Mean	SE	Mean	SE	Mean	SE	Mean	SE	Mean	SE	Mean	SE	Mean	SE		
BMI by HOMA (n)																						
Low BMI-low HOMA (90, 100)	1,832	46.1	1,409	36.5	65.9	4.9	63.6	6.9	207	5.5	183	4.6	18.7	1.1	19.6	1.0	792	41.0	776	31.7		
Low BMI-high HOMA (50, 67)	1,834	62.6	1,416	44.3	69.7	6.8	60.8	8.2	221	7.4	171	5.5	19.2	1.5	17.3	1.2	882	55.2	753	38.4		
High BMI-low HOMA (50, 67)	1,803	62.4	1,385 <sup>a</sup>	44.0	71.3	7.0	72.3	8.3	208	7.4	176	5.5	19.7	1.5	19.4	1.2	867	55.1	750	38.3		
High BMI-high HOMA (91, 100)	1,823	46.1	1,544 <sup>a</sup>	36.8	68.3	5.1	57.1	6.9	222	5.5	188	4.6	17.0	1.1	16.9	1.0	796	41.2	777	31.9		
F and p-values	0.06	0.98	3.57	0.02	0.16	0.93	0.70	0.55	1.79	0.15	2.18	0.09	0.88	0.45	1.69	0.17	0.93	0.43	0.17	0.92		
FMI by HOMA (n)																						
Low-FMI-low HOMA (86, 101)	1,817	46.4	1,393 <sup>a</sup>	35.8	65.8	5.0	68.9	6.7	205	5.5	181	4.5	18.7	1.1	19.8	1.0	786	41.2	788	30.9		
Low FMI-high HOMA (54, 66)	1,759	62.2	1,408	44.9	64.7	7.0	62.4	8.3	215	7.4	173	5.6	18.6	1.5	17.8	1.2	904	54.9	780	39.7		
High FMI-low HOMA (54, 66)	1,827	61.8	1,409	44.7	71.2	6.9	64.8	8.5	211	7.3	178	5.6	19.5	1.5	19.0	1.2	878	54.5	730	39.0		
High FMI-high HOMA (87, 101)	1,866	46.5	1,547 <sup>a</sup>	36.3	71.1	5.1	55.9	6.9	225	5.5	186	4.5	17.3	1.1	16.6	1.0	782	41.5	759	31.1		
F and p-values	0.61	0.61	3.68	0.01	0.31	0.82	0.59	0.62	2.30	0.08	1.22	0.30	0.53	0.66	1.91	0.13	1.63	0.18	0.50	0.68		

RBC-folate: Red blood cell folate; SE: Standard error; n.s.: Not significant. Significant differences ( $p < 0.05$ ) between groups are indicated by the same superscripts letters. F tests the effect of the model.



**Figure 1.**

Plasma vitamin B<sub>12</sub> concentrations (pmol/L) according to the combination between categories of FMI and HOMA by sex.

lower B-vitamins intake in adolescents with higher BMI while they have higher B-vitamins density diets. This can be explained by the fact that in the HELENA study, adolescents with higher BMI reported lower total energy diets likely affected by underreporting (35). Another reason could be that obese adolescents may have repeated short-term restrictive diets (34); this could be the case in our study (35), because adolescents with high BMI and high FMI were those with lower total energy intakes.

There were several differences between groups constituted by the combination of body composition markers and HOMA insulin sensitivity in terms of B-vitamins biomarkers concentrations. However, only adolescents belonging to the most favorable group in terms of combination between FMI and HOMA (lower FMI combined with lower HOMA) have higher plasma vitamin B<sub>12</sub> concentrations, as it had been already shown by another study (13). However, the previous study did not include the analyses of B-vitamins dietary intake, as it has been shown in our study. This fact cannot be explained by higher vitamin B<sub>12</sub> intakes but it might be due to lower vitamin B<sub>12</sub> density diets in comparison to those adolescents belonging to the less favorable combination groups, as shown in table II, likely resulting from the restrictive diets or underreporting behaviors previously mentioned. Besides, while we have obtained statistically significant differences for vitamin B<sub>6</sub> intake regarding BMI and FMI based on the median categories combined with the categories of HOMA, this was not the case for vitamin B<sub>12</sub>. This is of lots of interest owing to the fact that plasma vitamin B<sub>12</sub> is supposed to reflect changes in day-to-day diet rather than HoloTC, which is more efficient in predicting long-term diet changes (9), for which no difference among categories has been found.

Given that there seems not to be a plausible effect of adiposity and insulin resistance on plasma vitamin B<sub>12</sub> concen-

trations, linked to vitamin B<sub>12</sub> intake, we should consider the hypothesis that low vitamin B<sub>12</sub> concentrations could influence insulin resistance. A recent genome-wide analysis (36) suggested that increased DNA methylation is associated with increased BMI in adults, and vitamin B<sub>12</sub> is a determinant for DNA methylation.

Also, in a recent review about the transmission of obesity-adiposity and related disorders from the mother to the newborns (37), it is mentioned that in rural areas from India, with mothers consuming mainly vegetarian diets, the most insulin-resistant children were born to mothers who had low vitamin B<sub>12</sub> but high folate levels, suggesting that a balance between these two vitamins is essential. This might be a consequence from the fact that folate together with vitamins B<sub>12</sub> and B<sub>6</sub>, among others, regulate maternal 1-carbon metabolism, which influences cellular growth and differentiation by helping synthesis of nucleic acids. However, this explanation is not helpful to understand our results, owing to the differences in the characteristics of the population group and because our sample showed higher prevalence of folate deficiency than the vitamin B<sub>12</sub> one (30).

Other study (38) showed the importance of levels of homocysteine in insulin resistance with an improvement of it with a lowering homocysteine by folate + vitamin B<sub>12</sub> treatment and a correlation of 0.60 between homocysteine levels and insulin resistance, as was also shown by similar studies.

Our findings might be of particular concern in case of adolescents diagnosed as obese patients because they will be prone to a further decrease in vitamin B<sub>12</sub> concentrations if metformin therapy is recommended for the treatment of type 2 diabetes (15). For instance, in adults, vitamin B<sub>12</sub> malabsorption was observed in approximately 20% of patients using metformin, and this was associated with a 4-24% decrease of vitamin B<sub>12</sub> concentrations (39).

## LIMITATIONS AND STRENGTHS

The cross-sectional design of the study represents a limitation owing to the fact that causality cannot be established. The clinical interpretation of our findings is unclear because a very low percentage of adolescents presented inadequate intakes or very low serum concentrations of vitamin B<sub>12</sub>. However, the observed association between vitamin B<sub>12</sub> plasma level and body composition and insulin resistance warns us about the importance of having a healthy nutritional status and elucidates the fact that overweight or obese people cannot get nutritional deficiencies. The use of harmonized and standardized procedures in a large sample of adolescents from Europe (18) should be considered as the main strength of the study, as well as the use of previously validated questionnaires and procedures (18,19). Besides, the calculation of the usual intake values based on the MSM method to prevent limitations of the 24-h recalls (22), together with the use of widely accepted micronutrient biomarkers, strengthen the reliability of the observations with any other marker or symptom (40).

## CONCLUSION

Obesity is not only associated with cardiovascular conditions, cancer, dyslipidemias, etc., but also with micronutrient deficiencies which can lead to serious health problems. Among vitamins B<sub>6</sub>, folate, and B<sub>12</sub>, it seems that vitamin B<sub>12</sub> is the one most consistently and negatively associated with BMI, FMI, and insulin sensitivity or resistance (HOMA) without discriminating between sexes of European adolescents.

In male and female adolescents with combined higher adiposity measured with fat mass index and higher HOMA insulin sensitivity, low vitamin B<sub>12</sub> plasma concentrations were observed. These differences do not seem explained by dietary vitamin B<sub>12</sub> intake. Further studies are necessary to elucidate the potential role of low vitamin B<sub>12</sub> concentrations in the development of insulin resistance in adolescents in order to identify a plausible biological mechanism.

## REFERENCES

- Lien N, Henriksen HB, Nymoel LL, Wind M, Klepp KI. Availability of data assessing the prevalence and trends of overweight and obesity among European adolescents. *Public Health Nutr* 2010;13(10A):1680-7.
- Global Health Risk. Mortality and burden disease attributable to selected major risks. Geneva, Switzerland: World Health Organization; 2009.
- Division of Nutrition, Physical Activity, and Obesity, National Center for Chronic Disease Prevention and Health Promotion. Updated June 19, 2015/07/2016. Available from: <https://www.cdc.gov/obesity/childhood/causes.html>.
- Pinhas-Hamiel O, Doron-Panush N, Reichman B, Nitzan-Kaluski D, Shalitin S, Geva-Lerner L. Obese children and adolescents: A risk group for low vitamin B12 concentration. *Arch Pediatr Adolesc Med* 2006;160(9):933-6.
- Pinhas-Hamiel O, Newfield RS, Koren I, Agmon A, Lilos P, Phillip M. Greater prevalence of iron deficiency in overweight and obese children and adolescents. *Int J Obes Relat Metab Disord* 2003;27(3):416-8.
- Frankenfield DC, Rowe WA, Cooney RN, Smith JS, Becker D. Limits of body mass index to detect obesity and predict body composition. *Nutrition* 2001;17(1):26-30.
- Tresaco B, Bueno G, Moreno LA, Garagorri JM, Bueno M. Insulin resistance and impaired glucose tolerance in obese children and adolescents. *J Physiol Biochem* 2003;59(3):217-23.
- Iglesia I, Mouratidou T, González-Gross M, Huybrechts I, Breidenassel C, Santabarbara J, et al. Foods contributing to vitamin B6, folate, and vitamin B12 intakes and biomarkers status in European adolescents: The HELENA study. *Eur J Nutr* 2016. DOI: 10.1007/s00394-016-1221-1.
- Institute of Medicine Dietary Reference Intakes for Thiamin, Riboflavin, Niacin, Vitamin B6, Folate, Vitamin B12, Pantothenic Acid, Biotin, and Choline, a report of the Standing Committee on the Scientific Evaluation of Dietary Reference Intakes and its Panel on Folate, Other B Vitamins, and Choline and Subcommittee on Upper Reference Levels of Nutrients. Food and Nutrition Board. Washington DC: National Academy Press; 1998.
- Clemens TL. Vitamin B12 deficiency and bone health. *N Engl J Med* 2014;371(10):963-4.
- Kerr MA, Livingstone B, Bates CJ, Bradbury I, Scott JM, Ward M, et al. Folate, related B vitamins, and homocysteine in childhood and adolescence: potential implications for disease risk in later life. *Pediatrics* 2009;123(2):627-35.
- Hunt A, Harrington D, Robinson S. Vitamin B12 deficiency. *BMJ* 2014;349:g5226.
- Ho M, Halim JH, Gow ML, El-Haddad N, Marzulli T, Baur LA, et al. Vitamin B12 in obese adolescents with clinical features of insulin resistance. *Nutrients* 2014;6(12):5611-8.
- Laing SP, Swerdlow AJ, Slater SD, Burden AC, Morris A, Waugh NR, et al. Mortality from heart disease in a cohort of 23,000 patients with insulin-treated diabetes. *Diabetologia* 2003;46(6):760-5.
- Reinstatler L, Qi YP, Williamson RS, Garn JV, Oakley GP Jr. Association of biochemical B12 deficiency with metformin therapy and vitamin B12 supplements: The National Health and Nutrition Examination Survey, 1999-2006. *Diabetes Care* 2012;35(2):327-33.
- González-Gross M, Benser J, Breidenassel C, Albers U, Huybrechts I, Valtuena J, et al. Gender and age influence blood folate, vitamin B12, vitamin B6, and homocysteine levels in European adolescents: The Helena Study. *Nutr Res* 2012;32(11):817-26.
- Healthy Lifestyle in Europe by Nutrition in Adolescence. Cited 1/07/2016. Available from: <http://www.helenastudy.com/abstract.php>.
- Moreno LA, González-Gross M, Kersting M, Molnar D, De Henauw S, Beghin L, et al. Assessing, understanding and modifying nutritional status, eating habits and physical activity in European adolescents: The HELENA (Healthy Lifestyle in Europe by Nutrition in Adolescence) Study. *Public Health Nutr* 2008;11(3):288-99.
- Moreno LA, De Henauw S, González-Gross M, Kersting M, Molnar D, Gottrand F, et al. Design and implementation of the Healthy Lifestyle in Europe by Nutrition in Adolescence Cross-Sectional Study. *Int J Obes (Lond)* 2008;32(Suppl 5):S4-11.
- Beghin L, Castera M, Manios Y, Gilbert CC, Kersting M, De Henauw S, et al. Quality assurance of ethical issues and regulatory aspects relating to good clinical practices in the HELENA Cross-Sectional Study. *Int J Obes (Lond)* 2008;32(Suppl 5):S12-8.
- Vereecken CA, Covents M, Sichert-Hellert W, Alvira JM, Le Donne C, De Henauw S, et al. Development and evaluation of a self-administered computerized 24-h dietary recall method for adolescents in Europe. *Int J Obes (Lond)* 2008;32(Suppl 5):S26-34.
- Harttig U, Haubrock J, Knuppel S, Boeing H. The MSM program: Web-based statistics package for estimating usual dietary intake using the Multiple Source Method. *Eur J Clin Nutr* 2011;65(Suppl 1):S87-91.
- González-Gross M, Breidenassel C, Gómez-Martínez S, Ferrari M, Beghin L, Spinneker A, et al. Sampling and processing of fresh blood samples within a European multicenter nutritional study: Evaluation of biomarker stability during transport and storage. *Int J Obes (Lond)* 2008;32(Suppl 5):S66-75.
- Kimura M, Kanehira K, Yokoi K. Highly sensitive and simple liquid chromatographic determination in plasma of B6 vitamins, especially pyridoxal 5'-phosphate. *J Chromatogr A* 1996;722(1-2):295-301.
- Ulleland M, Eilertsen I, Quadros EV, Rothenberg SP, Fedosov SN, Sundrehagen E, et al. Direct assay for cobalamin bound to transcobalamin (holo-transcobalamin) in serum. *Clin Chem* 2002;48(3):526-32.
- Matthews DR, Hosker JP, Rudenski AS, Naylor BA, Treacher DF, Turner RC. Homeostasis model assessment: Insulin resistance and beta-cell function from fasting plasma glucose and insulin concentrations in man. *Diabetologia* 1985;28(7):412-9.
- Iglesia I, Mouratidou T, González-Gross M, Novakovic R, Breidenassel C, Jiménez-Pavón D, et al. Socioeconomic factors are associated with folate and vitamin B12 intakes and related biomarkers concentrations in European adolescents: The Healthy Lifestyle in Europe by Nutrition in Adolescence study. *Nutr Res* 2014;34(3):199-209.
- Nagy E, Vicente-Rodríguez G, Manios Y, Beghin L, Iliescu C, Censi L, et al. Harmonization process and reliability assessment of anthropometric measurements in a multicenter study in adolescents. *Int J Obes (Lond)* 2008;32(Suppl 5):S58-65.
- Slaughter MH, Lohman TG, Boileau RA, Horswill CA, Stillman RJ, Van Loan MD, et al. Skinfold equations for estimation of body fatness in children and youth. *Hum Biol* 1988;60(5):709-23.
- González-Gross M, Benser J, Breidenassel C, Albers U, Huybrechts I, Valtuena J, et al. Gender and age influence blood folate, vitamin B12, vitamin B6, and homocysteine levels in European adolescents: The Helena Study. *Nutr Res* 2012;32(11):817-26.
- MacFarlane AJ, Greene-Finestone LS, Shi Y. Vitamin B12 and homocysteine status in a folate-replete population: Results from the Canadian Health Measures Survey. *Am J Clin Nutr* 2011;94(4):1079-87.
- Beghin L, Huybrechts I, Vicente-Rodríguez G, De Henauw S, Gottrand F, González-Gross M, et al. Main characteristics and participation rate of European adolescents included in the HELENA study. *Arch Public Health* 2012;70(1):14.
- Kimmons JE, Blanck HM, Tohill BC, Zhang J, Khan LK. Associations between body mass index and the prevalence of low micronutrient levels among US adults. *Med Gen Med* 2006;8(4):59.
- Damms-Machado A, Weser G, Bischoff SC. Micronutrient deficiency in obese subjects undergoing low calorie diet. *Nutr J* 2012;11:34.
- Huybrechts I. Dietary under-reporting by overweight and obese adolescents: Results from the HELENA Study. *Public Health Nutrition* 2012;15(Special Issue 8A):1555-.
- Dick KJ, Nelson CP, Tsaprouni L, Sandling JK, Aissi D, Wahl S, et al. DNA methylation and body-mass index: A genome-wide analysis. *Lancet* 2014;383(9933):1990-8.
- Jajnik CS. Transmission of obesity-adiposity and related disorders from the mother to the baby. *Ann Nutr Metab* 2014;64(Suppl 1):8-17.
- Setola E, Monti LD, Galluccio E, Palloschi A, Fraggaso G, Paroni R, et al. Insulin resistance and endothelial function are improved after folate and vitamin B12 therapy in patients with metabolic syndrome: Relationship between homocysteine levels and hyperinsulinemia. *Eur J Endocrinol* 2004;151(4):483-9.
- Mazokopakis EE, Starakis IK. Recommendations for diagnosis and management of metformin-induced vitamin B12 (Cbl) deficiency. *Diabetes Res Clin Pract* 2012;97(3):359-67.
- Henríquez-Sánchez P, Sánchez-Villegas A, Doreste-Alonso J, Ortiz-Andrellucchi A, Pfrimer K, Serra-Majem L. Dietary assessment methods for micronutrient intake: A systematic review on vitamins. *Br J Nutr* 2009;102(Suppl 1):S10-37.