



Anais da Academia Brasileira de Ciências

ISSN: 0001-3765

aabc@abc.org.br

Academia Brasileira de Ciências

Brasil

Nanni, Arthur; Roisenberg, Ari; Fachel, Jandyra M.G.; Mesquita, Gilberto; Danieli, Cristiano
Fluoride characterization by principal component analysis in the hydrochemical facies of Serra Geral
Aquifer System in Southern Brazil

Anais da Academia Brasileira de Ciências, vol. 80, núm. 4, diciembre, 2008, pp. 693-701

Academia Brasileira de Ciências

Rio de Janeiro, Brasil

Available in: <http://www.redalyc.org/articulo.oa?id=32713465010>


- How to cite
- Complete issue
- More information about this article
- Journal's homepage in redalyc.org

redalyc.org

Scientific Information System

Network of Scientific Journals from Latin America, the Caribbean, Spain and Portugal

Non-profit academic project, developed under the open access initiative



Anais da Academia Brasileira de Ciências (2008) 80(4): 693-701
(Annals of the Brazilian Academy of Sciences)
ISSN 0001-3765
www.scielo.br/aabc

Fluoride characterization by principal component analysis in the hydrochemical facies of Serra Geral Aquifer System in Southern Brazil

ARTHUR NANNI¹, ARI ROISENBERG², JANDYRA M.G. FACHEL³, GILBERTO MESQUITA⁴
and CRISTIANO DANIELI⁵

¹Departamento de Geoquímica, Instituto de Geociências, UFRGS, Av. Bento Gonçalves, 9.500, prédio 43126/105
91501-970 Porto Alegre, RS, Brasil

²Departamento de Geoquímica, Instituto de Geociências, UFRGS, Av. Bento Gonçalves, 9.500, prédio 43126/103
91501-970 Porto Alegre, RS, Brasil

³Departamento de Estatística, Instituto de Matemática, UFRGS, Av. Bento Gonçalves 9.500
91501-970 Porto Alegre, RS, Brasil

⁴Núcleo de Assessoria Estatística, Instituto de Matemática, UFRGS, Av. Bento Gonçalves 9.500
91501-970 Porto Alegre, RS, Brasil

⁵Instituto de Geociências, UFRGS, Av. Bento Gonçalves, 9.500, prédio 43126/105, 91501-970 Porto Alegre, RS, Brasil

*Manuscript received on May 7, 2007; accepted for publication on May 14, 2008;
presented by ALCIDES N. SIAL*

ABSTRACT

Principal component analysis is applied to 309 groundwater chemical data information from wells in the Serra Geral Aquifer System. Correlations among seven hydrochemical parameters are statistically examined. A four-component model is suggested and explains 81% of total variance. Component 1 represents calcium-magnesium bicarbonate groundwaters with long time of residence. Component 2 represents sulfated and chlorinated calcium and sodium groundwaters; Component 3 represents sodium bicarbonated groundwaters; and Component 4 is characterized by sodium sulfated with high fluoride facies. The components' spatial distribution shows high fluoride concentration along analyzed tectonic fault system and aligned on northeast direction in other areas, suggesting other hydrogeological fault systems. High fluoride concentration increases according to groundwater pumping depth. The Principal Component Analysis reveals features of the groundwater mixture and individualizes water facies. In this scenery, it can be determined hydrogeological blocks associated with tectonic fault system here introduced.


Key words: fractured aquifer, geostatistics, GIS, groundwater, hydrogeology.

INTRODUCTION

Water scarcity and increasing human consumption requires new sources of water with adequate potability. Therefore, an emergent necessity to understand groundwater resources is evident. Low cost treatment and technical advances in exploration turns groundwater into a vital and precious natural resource.

Fluoride content in water, like other chemical spe-

in excess. High fluoride contents in groundwater are responsible for human and animal health problems, including dental and skeletal fluorosis, which is detected worldwide, like in China (Lin et al. 2004, Genxu and Guo 2001), India (Kumar et al. 2001), Kenya (Motu 2002), and Israel (Kafri et al. 1989), among other countries. In the Rio Grande do Sul State, southern Brazil, the endemic fluorosis has been detected in several districts (e.g. Venancio Aires, Santa Cruz do Sul).



pogenic contamination (fertilizer application, brick and aluminum smelters, sewage piles and other sources) or to a natural origin, such as prolonged water-rock interactions (Nordstrom et al. 1989). The drinking water limit recommended by the World Health Organization for fluoride is 1.5 mg/L (WHO 2002), and in a moderate concentration (0.7-1.2 mg/L) prevents dental cavities.

The main aim of the current study is to understand the distribution of high-fluoride waters in the Serra Geral Aquifer System (SGAS) and to identify the hydrochemical types and their spatial distribution, using the Principal Component Analysis (PCA). PCA is an important tool for understanding the large quantity of data involved in extended aquifer studies (Invernizzi and Oliveira 2004). This aids to define geological sources and pathways for high fluoride, consequently, assisting future well locations and management of SGAS waters. Investigations about the source of high fluoride concentrations in previous studies had shown a relationship with tectonic structures and with pumping of deep groundwater.

GEOLOGY AND HYDROGEOLOGY OF THE STUDY AREA

The study area is located in the Rio Grande do Sul State, southern Brazil and extends from 27°S to 31°S and from 50°W to 57°W, corresponding approximately to 164.207 km² in the Parana Basin. The region is covered by a basaltic to rhyolitic Mesozoic volcanic sequence belonging to the Serra Geral Formation (SGF), whose thickness varies from 50 to 1000 meters (average of 550 meters), constituting a fractured aquifer that provides public water supply to more than 80% of the cities in the area.

SGAS hydrochemical characteristics indicate the influence of water mixing with other sedimentary aquifer belonging to the Parana Basin (Szikszay et al. 1981, Fraga 1992, Portela Filho et al. 2002, 2004). This aquifer is directly superimposed by the Guarany Aquifer System/GAS (Campos 2000) that has been the focus of several studies in the last few years, due to its spatial extent and storage potentiality as a transnational aquifer.

Tectonic structures cut the SGF controlling terrain surface and hydrochemical characteristics of the related

that may be considered as hydrogeological blocks (Lisboa 1996, Lisboa and Menegotto 1997, Machado 2005). The tectonic block limited by the Terra de Areia-Posadas Fault System and Mata-Jaguari Fault System is uplifted in the south-central area. The adjacent block to the north shows a gradual terrain lowering from east to west, conditioned by NE normal faults, parallel to the Leao and Perimpo fault systems.

The identification of different water facies for SGAS focused on tectonic block separations is important to define structures where high fluoride groundwater can ascend from deep levels to SGAS. This tectonic control for the high fluoride groundwaters has been reported in other regions (Licht 2001). In the study area the SGAS fluoride average concentrations are around 0.24 mg/L, with a minimum value of 0.02 mg/L, and the highest at 3.03 mg/L. Machado (2005) describes the influence of Ca²⁺HCO₃⁻ meteoric recharge on the SGAS and mixture mechanisms between the SGAS and the GAS inputting Na⁺HCO₃⁻ with SO₄²⁻ and Cl⁻ water to SGAS.

MATERIALS AND METHODS

HYDROCHEMICAL DATA

Hydrogeological data represent a network of 309 deep wells (Fig. 1) that exposes only SGAS groundwater used in public and private water supply. All data information was provided by governmental groundwater management agencies.

This study regarded only seven major parameters to evaluate hydrochemical facies in the piper diagram, namely fluoride, Na⁺, Ca²⁺, Mg²⁺, HCO₃⁻, SO₄²⁻ and Cl⁻.

STATISTICAL ANALYSIS

The PCA was performed using SPSS version 8 software (Nie et al. 1975) and tested for two, three and four principal components using the eigenvalue equal to 1 criterion (Kaiser 1958). Before this procedure, outlier suppression was carried out through boxplots and dispersion charts for all variables. The Varimax orthogonal rotation was used (Kaiser 1958), in order to facilitate Factor Analysis interpretation (Invernizzi and Oliveira 2004). Princi-



FLUORIDE IN FRACTURED AQUIFER BY GEOSTATISTIC TOOLS




Fig. 1 – Location map of SGAS(gray) in Rio Grande do Sul State, sampling wells and main tectonic fault systems: (1) Mata-Jaguari, (2) Areia-Posadas, (3) Perimpo, and (4) Leao.

SPATIAL ANALYSIS

Spatial distribution and interpretation of principal components defined by PCA was performed on Geographical Information System (GIS) with Quantum GIS version 0.7.4 software (Sherman et al. 2005). Cluster results were classified, considering the most important chemical parameter as an identification name.

The relationship between depth and fluoride concentration considers the maximum absolute groundwater pumping depth in each well. This procedure was carried out with Labplot version 1.5 software (Gerlach 2004) in order to investigate whether fluoride concentration in SGAS increases with depth.

RESULTS AND DISCUSSIONS

PRINCIPAL COMPONENT ANALYSIS

Table I presents the correlation coefficient matrix for hydrochemical data in SGAS

Component 1 is mainly influenced by Ca^{2+} , and Mg^{2+} , and explains 37.43% of the total variance, representing calcium-magnesium bicarbonated groundwaters with long residence time, evidenced by Mg^{2+} . Component 2 is defined by SO_4^{2-} and Cl^- , represents sulfated and chlorinated, calcium and magnesium groundwaters.

Component 3 is explained by HCO_3^- and Na^+ , defining sodium bicarbonated groundwaters.

Component 4 is defined by F, followed by SO_4^{2-} and Cl^- , corresponds to sodium sulfate with high fluoride facies.

Fluoride participates with similar intensity in Components 2 and 3, suggesting two groundwater sources that contain relatively high fluoride concentration. Therefore, fluoride was associated with sodium and sodium bicarbonated groundwaters.

Based on the above, cluster analysis was performed and selected a solution with nine clusters, since it




TABLE I
Correlation coefficient.

Parameter	Na ⁺	Mg ²⁺	Ca ²⁺	Cl ⁻	SO ₄ ²⁻	HCO ₃ ⁻	F ⁻
Na ⁺	1.0	0.0069	-0.043	0.235	0.263	0.46	0.339
Mg ²⁺		1.0	0.549	0.351	0.093	0.457	0.096
Ca ²⁺			1.0	0.273	0.207	0.4	-0.012
Cl ⁻				1.0	0.457	0.352	0.198
SO ₄ ²⁻					1.0	0.196	0.385
HCO ₃ ⁻						1.0	0.094
F ⁻							1.0

TABLE II
Rotated matrix with four components and characteristic parameters.

Parameters	Components			
	1	2	3	4
Mg ²⁺	.874	.063	.072	.139
Ca ²⁺	. 811	. 231	-.045	-.043
Na ⁺	-.143	.188	. 870	.271
Cl ⁻	.353	. 707	. 206	-.051
HCO ₃ ⁻	. 572	.095	. 699	-.091
SO ₄ ²⁻	.030	. 872	.074	.237
F ⁻	.067	.145	.150	.956
Explained variance	37.433	20.260	12.694	10.770
Cumulative % of variance	37.433	57.693	70.386	81.156

The cluster interpretation was based on final center scores. Thus, each cluster was renamed using related ion predominance (Table III). In order to facilitate the reading it was decided to designate Component 1 as **CaMg**, Component 2 as **SO4**, Component 3 as **Na**, and Component 4 as **F**.

It can be defined that clusters 3 and 4 are composed predominantly by **F** component groundwaters. Clusters 1 and 7 comprise the **CaMg** water facies. Cluster 8 is composed by **Na** component. The remaining groups show more than one component in high concentration.

In the water facies, components **SO4**, **Na**, and **F** were added to the main water facies. Therefore, these clusters were called **SO4Na**, **SO4F**, and **NaF**, due to

HYDROCHEMISTRY

The geochemical data were plotted in a piper diagram (Fig. 2), with the four renamed cluster defined by the PCA. The majority of wells (212 out of 309 wells) have a composition related to **meteoric waters** with Ca²⁺HCO₃⁻ nature and do not include high fluoride waters. In this case, 212 samples are not represented in the diagrams, in order to obtain a better fit by the other groups.

The **CaMg** facies represents a predominant HCO₃⁻ water type where Mg⁺² appears in more than 50% of the wells (Fig. 2).

The **Na** group shows their typical distribution in the piper diagram, representing Na⁺HCO₃⁻ waters.



FLUORIDE IN FRACTURED AQUIFER BY GEOSTATISTIC TOOLS

TABLE III
Components final cluster centers and clusters name reclassification.

Component	Cluster								
	1	2	3	4	5	6	7	8	9
CaMg	1.6143	−0.6137	−0.228	0.511	−0.152	−0.724	2.042	−0.722	−1.720
SO4	0.987	4.411	−0.377	−1.198	−0.113	9.303	−0.236	−0.260	−0.348
Na	−0.892	3.032	−0.881	−0.667	−0.305	−0.845	1.092	1.653	2.691
F	−0.330	−1.357	1.244	6.986	−0.250	2.849	0.285	−0.034	4.786
Cases	22	2	19	3	212	2	11	36	2
Water Facies	CaMg	SO4Na	F	F	Meteoric	SO4F	CaMg	Na	NaF

different groundwater sources or point out to mixture features during groundwater ascending recharge.

SO4Na, **SO4F**, and **NaF** facies appear only in two wells each. Both wells compose distinct groups as showed in Figure 2.

The spatial distribution of water facies confirms that fluoride has a relationship with the analyzed fault systems Mata-Jaguari, Terra de Areia-Posadas, Perimpo, and Leao (Fig. 3).

Meteoric Waters occurs in all SGAS, but more significantly in the tectonic block limited by Terra de Areia-Posadas, Jaguari-Mata and Perimpo Fault Systems, due to the uplift condition of this block.

The **CaMg** facies is spatially related to discharge zones of SGAS in the west, northwest, and southeast regions, representing waters at the final percolation stage under long time residence.

The **Na** facies is dominant in the northwest region, where SGF achieves the maximum thickness in the study area, suggesting a hydrogeological tectonic block. This characteristic indicates that ascendant waters in this region present only Na^+ enrichment, as a result of water-rock interaction in SGF. Thus, the non-interference of GAS in this scenery is evident, probably due to the maximum thickness of SGF.

SO4Na facies appears only in the northeast alignment direction of the Uruguay River, showing a close relationship to this regional alignment. **SO4F** facies presents a similar behavior, but occurs only in the east-west section of the Uruguay River.

NaF facies is spatially linked to the Na hydrogeo-

hydrogeological condition.

The **F** group related that high fluoride groundwaters are distributed in the central-northeast portion of the study area. Aligned wells associated with the main tectonic systems (Terra de Areia-Posadas, Leao and Perimpo) suggest a water ascension recharge. New tectonic systems are introduced in the present study, corresponding to the linear distribution of wells belonging to the SGAS facies in the central portion of SGAS, which is the nomination Fontoura Xavier-Parai and Graeff-Barracao hydrogeological alignments (Fig. 3). It must be stressed that the introduced structures are parallel to Leao and Perimpo Fault Systems.

The distribution of **NaF** facies is similar to the **Na** facies (Fig. 3). Both are chemically very similar, distinguished by a more intensive fluoride participation in the **NaF** facies.

The **SO4F** and **SO4Na** facies are scarce and associated with major regional structures that define the Uruguay River (Fig. 3). The **SO4F** facies occurs in the north sector while the **SO4Na** facies occurs essentially in the west sector of the study area.

The composition of the volcanic rocks belonging to the SGF does not comprise fluorine rich minerals, which provide anomalous fluoride content in the SGAS facies. This enrichment could be better explained by the ascending groundwater recharge crossing older Parana Basin sedimentary sequences combined with long residence time and extreme confinement conditions.

The A-D geological cross-section (Figs. 1 and 2) demonstrates the east-west gradual terrain lowering




Fig. 2 – Water facies distribution in the Piper Diagram, showing two fluoride hydrochemical types (I – sulfated; II – bicarbonated).

The depth variation shows fluoride concentration increase according to groundwater pumping depth (Fig. 5). It can be demonstrated that beyond F facies, all the other facies increase the fluoride content with the depth. This behavior probably is related to the influence of confined GAS and the proximity of other older aquifer systems.

CONCLUSIONS

The PCA methodology was efficient to discriminate hydrochemical water facies in a collection of more than

was achieved, using seven chemical components that validate six hydrochemical facies were distinguished in the SGAS, three of them containing substantial fluoride contents in combination with Na^+ and SO_4^{2-} . The spatial distribution of these facies confirms the presence of hydrogeological blocks limited by tectonic alignments.

Two tectonic structures are apparently present in the central area, indicated by linear distribution of high fluoride groundwater wells. An aquifer system status for SGAS is reaffirmed, considering complex chemical mixtures, recharge processes and percolation through the





Fig. 3 – Water facies distribution and tectonic fault systems: (1) Mata-Jaguari, (2) Terra de Areia-Posadas, (3) Perimpo, and (4) Leao. Introduced hydrogeological alignments are (5) Fontoura Xavier-Parai, (6) Victor Graeff-Barracão and (7) that indicates a tectonic block limit.




Fig. 4 – Geological cross-section showing the main tectonic structures responsible for high fluoride groundwater ascension. Modified from Machado, 2005.

The results were able to point out the influence of deeper confined aquifers on fluoride contents of the SGAS and add elements to minimize the costs for well location with better water quality.

(CNPq) and by the Fundação Estadual de Proteção Ambiental Henrique Luis (FEPAM) Rio Grande

RESUMO

A Análise de Componentes Principais foi aplicada




Fig. 5 – Relationship between fluoride concentration and groundwater pumping depth.

de quatro componentes foi utilizado por explicar 81% da variância total. A Componente 1 é representada por águas cálcio-magnesianas com longo tempo de residência, a Componente 2 representa águas bicarbonatadas sulfatadas e cloretadas, a Componente 3 representa águas bicarbonatadas sódicas e a Componente 4 é caracterizada por águas de fácies sódica e sulfatada com alto fluoreto. A distribuição espacial das componentes mostra águas com concentrações anômalas ao longo dos sistemas tectônicos de falhas, analisados e alinhados a NE em algumas áreas, sugerindo outros sistemas de falhas hidrogeológicos. As concentrações de fluoreto aumentam de acordo com a profundidade de bombeamento das águas. A Análise de Componentes Principais revelou feições de mistura e individualizou diferentes fácies de águas subterrâneas. Neste cenário, é possível determinar blocos hidrogeológicos associados com os sistemas tectônicos de falhas introduzidos no presente trabalho.

Palavras-chave: aquífero fraturado, geoestatística, SIG, águas subterrâneas, hidrogeologia.

REFERENCES

CAMPOS HCN. 2000. Modelación conceptual y matemática del Acuífero Guaraní, Cono Sur. Mapa hidrogeológico do aquífero Guaraní escala 1:250000. Acta Geol Leopold 4: 3–50.

em Recursos Minerais e Hidrogeologia. Universidade de São Paulo. 178 p.

GENXU W AND GUODONG C. 2001. Fluoride distribution in water and the governing factors of environment in arid north-west China. J Arid Environ 49: 601–614.

GERLACH S. 2004. Labplot 1.5.0 – An application for plotting and analysis of 2D and 3D functions and data. <http://labplot.sourceforge.net>.


INVERNIZZI A AND OLIVEIRA SMB. 2004. Hydrochemical characterization of a watershed through factor analysis. Águas Subterrâneas 18: 67–77.

KAFRI U, ARAD A AND HALICZ L. 1989. Fluorine Occurrence in Groundwater in Israel and its significance. J Hydrol 106: 109–129.

KAISER HF. 1958. The varimax criterion for analytic rotation factor analysis. Psychometrika 23: 187–200.

KUMAR VV AND SAI CST, RAO PLKM AND RAO CS. 2001. Studies on the distribution of fluoride in drinking water sources in Medchal Block, Ranga Reddy District, Andhra Pradesh, India. J Fluorine Chem 55: 229–236.

LICHT OAB. 2001. A geoquímica multielementar na gestão ambiental – identificação e caracterização de províncias geoquímicas naturais, alterações antrópicas da paisagem, áreas favoráveis à prospecção mineral e regiões de risco



FLUORIDE IN FRACTURED AQUIFER BY GEOSTATISTIC TOOLS

- LIN NF, TANG J AND BIAN JM. 2004. Geochemical environment and health problems in China. *Environ Geochem Health* 26: 81–88.
- LISBOA NA. 1996. Fácies, estratificações hidrogeoquímicas e seus controladores geológicos, em unidades hidrogeológicas do Sistema Aquífero Serra Geral, na Bacia do Paraná, Rio Grande do Sul. Tese de Doutorado. Curso de Pós-Graduação em Geociências. Universidade Federal do Rio Grande do Sul. 135 p.
- LISBOA NA AND MENEGOTTO E. 1997. Diferenciações hidrogeoquímicas no sistema aquífero Serra Geral no Rio Grande do Sul. In: XII SIMPÓSIO BRASILEIRO DE RECURSOS HÍDRICOS, p. 489–496.
- MACHADO JLF. 2005. Compartimentação espacial e arcabouço hidroestratigráfico do Sistema Aquífero Guarani no Rio Grande do Sul. Tese de Doutorado. Programa de pós-graduação em geologia – área de concentração em geologia sedimentar. Universidade do vale do rio dos Sinos. 237 p.
- MOTURI WKN, TOLE MP AND DAVIES TC. 2002. The contribution of drinking water towards dental fluorosis: a case study of Njoro Division, Nakuru District, Kenya. *Environ Geochem Health* 24: 123–130.
- NIE NJ, HULL CH, JENKINS JG, STEINBRENNER K AND BRENT DH. 1975. Statistical package for the socialsciences. 2nd ed. New York: 675 p.
- NORDSTROM DK, BALL JW, DONAHOE RJ AND WHITEMORE D. 1989. Groundwater chemistry and water-rock interaction at Stripa. *Geochim Cosmochim Acta* 53: 1727–1740.
- PORTELA FILHO CV, FERREIRA FJF, ROSA FILHO CV, BUCHMANN AC AND ROSTIROLLA SP. 2002. preliminar da conexão entre os Aquíferos Serra Geral e Guarani com base em dados aeromagnetométricos e hidroquímicos. In: CONGRESSO BRASILEIRO DE SUBTERRÂNEAS, 12. Florianópolis, SC, Brasil.
- PORTELA FILHO CV, FERREIRA FJF, ROSA FILHO CV, ROSTIROLLA SP. 2004. Compartimentação macroestrutural do Sistema Aquífero Serra Geral e sua relação com o Sistema Aquífero Guarani na região do arco de Ponta Grossa (Bacia do Paraná). *Rev Bras Geocienc* 3: 369–381.
- SHERMAN GE, SUTTON T, BLAZEK R AND LUTHI M. 2005. Quantum GIS User Guide – Version 0.7.4. <http://qgis.sourceforge.net/docs/userguide.html>.
- SZIKSZAY M, TEISSEDRE JM, BARNER U AND MACHADO JLF. 1981. Geochemical and isotopic characteristics of groundwater in the state of São Paulo, Brazil. *J Hydro* 54: 23–32.
- WHO (WORLD HEALTH ORGANIZATION). 2002. Guidelines for drinking-water quality. Environmental Health Criteria 227, 274 pp. World Health Organization, World Health Organization, Geneva.