



Anais da Academia Brasileira de Ciências

ISSN: 0001-3765

aabc@abc.org.br

Academia Brasileira de Ciências

Brasil

BOUROTTE, CHRISTINE; BERTOLO, REGINALDO; ALMODOVAR, MARTA; HIRATA, RICARDO
Natural occurrence of hexavalent chromium in a sedimentary aquifer in Urânia, State of São Paulo,
Brazil

Anais da Academia Brasileira de Ciências, vol. 81, núm. 2, junio, 2009, pp. 227-242

Academia Brasileira de Ciências

Rio de Janeiro, Brasil

Available in: <http://www.redalyc.org/articulo.oa?id=32713477009>


- How to cite
- Complete issue
- More information about this article
- Journal's homepage in redalyc.org

redalyc.org

Scientific Information System

Network of Scientific Journals from Latin America, the Caribbean, Spain and Portugal

Non-profit academic project, developed under the open access initiative



Anais da Academia Brasileira de Ciências (2009) 81(2): 227-242
(Annals of the Brazilian Academy of Sciences)
ISSN 0001-3765
www.scielo.br/aabc

Natural occurrence of hexavalent chromium in a sedimentary aquifer in Urânia, State of São Paulo, Brazil

CHRISTINE BOUROTTE¹, REGINALDO BERTOLO², MARTA ALMODOVAR³ and RICARDO HIRATA⁴

¹Instituto de Geociências, Universidade de São Paulo

Rua do Lago, 562, Cidade Universitária, 05508-080 São Paulo, SP, Brasil

²Instituto de Geociências, Universidade de São Paulo

Rua do Lago, 562, Cidade Universitária, 05508-080 São Paulo, SP, Brasil

³CETESB, Companhia de Tecnologia de Saneamento Ambiental do Estado de São Paulo
Av. Professor Frederico Hermann Jr., 345, Alto de Pinheiro, 05459-900 São Paulo, SP, Brasil

⁴Instituto de Geociências, Universidade de São Paulo

Rua do Lago, 562, Cidade Universitária, 05508-080 São Paulo, SP, Brasil

*Manuscript received on August 6, 2008; accepted for publication on December 19, 2008;
presented by ALCIDES N. SIAL*

ABSTRACT

Anomalous concentrations of hexavalent chromium have been detected in groundwater of the Adamantina Aquifer at least 54 municipalities located in the northwestern region of the State of São Paulo, southeast Brazil, occasionally exceeding the permitted limit for human consumption (0.05 mg.L^{-1}). An investigation was conducted in the municipality of Urânia, where the highest concentrations of chromium were detected regionally. It was defined that the origin of this contamination is natural, since high concentrations of chromium were detected in aquifer sandstones (average of 221 ppm) and also in pyroxenes (6000 ppm), one of the main heavy minerals found in the sediments. Besides, other possible diffuse or point sources of contamination were observed in the study area. Stratification of groundwater quality was observed and the highest concentrations of Cr^{6+} were detected at the base of the aquifer (0.12 mg.L^{-1}) where groundwater shows elevated values for redox potential (472.5 mV) and pH (8.61). The origin of Cr^{6+} in water may be associated with the weathering of pyroxene (augite), followed by the oxidation of Cr^{3+} by manganese oxides. The highest concentrations of Cr^{6+} are probably related to desorption reactions, due to the anomalous alkaline pH found in groundwater at the base of the aquifer.


Key words: chromium, groundwater, aquifer, natural contamination.

INTRODUCTION

Chromium has been identified as a public health problem due to its toxic effects even at low exposure levels (ATSDR 2000). In the environment, chromium exists in two main oxidation states, Cr^{3+} and Cr^{6+} , which develop different geochemical and biological activities, since Cr^{3+} is an essential metal nutrient and Cr^{6+} is carcinogenic. Cr^{6+} is more mobile, labile and toxic than

tions in natural waters. Under acidic and reducing conditions, Cr^{3+} species will predominate in water. Cr^{6+} species will prevail under alkaline and oxidizing conditions.

Groundwater contamination by chromium is generally associated with such anthropic activities as mining, electroplating, leather tanning and chemical manufacturing. High levels of chromium are



centrations of naturally occurring hexavalent chromium have been reported in groundwater in the Aroma Red Sands Aquifer, California (Gonzalez et al. 2005), at León Valley, Mexico (Robles-Camacho and Armienta 2000), in the western Mojave Desert, California (Ball and Izbicki 2004) and Arizona (Robertson 1995).

Although the natural presence of chromium in aquifers is scarce and little studied, concentrations of chromium higher than the recommended value for drinking water (0.05 mg.L^{-1}) have been known since 1977 in the Adamantina Aquifer (Almodovar and Pacheco 1995), located in the northwestern portion of the State of São Paulo. Water supplies in this region are primarily obtained via deep groundwater wells, most of which are managed by public water supply companies.

The main objective of this paper is that of characterizing the distribution of chromium species in groundwater of the Adamantina Aquifer and present the results of an investigation performed in the municipality of Urânia (Fig. 1), where the highest concentrations were detected. The concentration of major ions and water quality parameters were also measured and their relationship with chromium occurrence was considered, as well as the relationship between chemical species and well depth. The results shown in this paper are part of a wider study and some interpretations of the occurrence and distribution of chromium are also discussed.

STUDY AREA

GEOLOGICAL SETTING

The study area is located in the Paraná Basin, which is a volcanic-sedimentary basin that covers virtually 1 million km^2 in Brazil. More specifically, the study site is located in the western part of the Paulista Plateau, which covers 50% of the State of São Paulo. The parent rock consists of sedimentary rocks of the Bauru Basin (Upper Cretaceous age) and overlays basalts of the Serra Geral Formation (Jurassic-Cretaceous age).

The Bauru Basin covers an area of 104,000 km^2 and 42% of the area of the State of São Paulo. Based on the conception of Fernandes and Coimbra (2000), the Bauru Basin has two chronocorrelated geological units,


portions of the basin. The Caiuá Group is subdivided into the Rio Paraná, Goio-Erê and Santo Anastácio Formations, which are present in the western portion of the State of São Paulo. The Bauru Group, which is predominant in the State of São Paulo, outcrops in oriental domains of the Basin and is subdivided into the Vale do Rio do Peixe, Araçatuba, São José do Rio Preto, Presidente Prudente and Marília Formations (Fig. 1).

In this context, sedimentary rocks of the Vale do Rio do Peixe Formation or, in other words, the Adamantina Formation conceived by Soares et al. (1980), outcrop in the municipality of Urânia. This stratigraphic unit is constituted of well-sorted fine to very fine sandstones, interbedded with siltstones. Carbonate cementation is found locally. The sandstones are disposed in submetric tabular layers of massive aspect, presenting zones of coarse tabular bedding and zones with tabular and through cross-bedding.

Fernandes and Coimbra (2000) suggest that deposition of the Vale do Rio do Peixe Formation occurred primarily by eolic action in extensive plain-like areas and, secondarily, by occasional torrents, forming *wadis* deposits. Finally, the authors suggest that the lower contact of the Vale do Rio do Peixe Formation is gradual with the Santo Anastácio Formation or, as occurs in the study area, discordant and directly on the basalts of the Serra Geral Formation.

Brandt Neto et al. (1985) indicate that the mineralogy of the sediments is mainly composed of quartz and, secondarily, feldspars, kaolinite, montmorillonite, opaque minerals and carbonates as cement. The origin of the sedimentary rocks is related to the deposition of sediments caused by the erosion and transport of other Phanerozoic sediments, metamorphic rocks of the Araxá and Canastra Groups, basalts of the underlying Serra Geral Formation and alkaline rocks of the Triângulo Mineiro (A.M. Coimbra, unpublished data).

The average thickness of the Bauru Group is around 100 m. Greater thickness generally occurs on the crests of hills between the main rivers of the western portion of the State of São Paulo. In the Urânia region, the sedimentary rocks were found to have a thickness of up to



NATURAL OCCURRENCE OF HEXAVALENT CHROMIUM




Fig. 1 – Location of the study area and stratigraphy of Bauru Basin (modified from Fernandes and Coimbra 2000)

gion, the Serra Geral Formation has a thickness of between 500 and 1500 m. In the Urânia region, deep supply wells exploiting groundwater from the underlying Guarani Aquifer show basaltic rocks with a thickness of around 900 m.

HYDROGEOLOGICAL SETTING


There are two main hydro-stratigraphic units in this region, namely: the Bauru Aquifer System, which constitutes the main source of water supply, and the Serra Geral Aquifer, represented by basalts of the underlying Serra Geral Formation. The municipality of Urânia is located in the context of the traditionally named Adamantina Aquifer, in concordance with designation of the Adamantina Formation by Soares et al. (1980).

The Adamantina Aquifer can be characterized as a

area, where groundwater divides coincide with the age basins and the local effluent rivers correspond to discharge areas of the aquifer (Hirata et al. 1997). The aquifer is a moderately permeable formation, with average values of hydraulic conductivity and transmissivity close to 1.0×10^{-5} m/sec and $40 \text{ m}^2/\text{day}$, respectively (DAEE 1976).

The Serra Geral Aquifer system is constituted by flood basalts and associated intrusive rocks. It is characterized as a fractured, unconfined to semi-confined, discontinued, anisotropic and heterogeneous aquifer.

The northwestern region of the State of São Paulo shows an average temperature of 20°C and average precipitation of 1271 mm per year, which is distributed in an uneven form, with most precipitation occurring during the rainy period from December to March.



SAMPLING AND ANALYTICAL METHODS

Table I presents a summary of the groundwater sampling events from 1977 to 1999. Data from 144 deep supply wells located in 54 cities of the study area were initially obtained at a public water supply company. These data include in-situ and laboratory parameters obtained between 1977 and 1993. Fieldwork activities were then conducted in 12 localities in the region during December 1994, in order to complement the geographical investigation of chromium distribution (Almodovar and Pacheco 1995).

From June 1998 to June 1999, groundwater samples from 31 wells located in the municipality of Urânia were analyzed in order to assess variation of chromium concentrations with depth and pumping time. During sampling activities, water from public and private supply wells was collected, together with water from hand-dug shallow supply wells. In order to evaluate variation of chromium concentrations over a pumping period of 24 hours, a sampling program was performed during several pumping intervals at two deep supply wells (PP02 and PP04).

Samples collected between 1977 and 1993 were obtained and analyzed by a public water supply company. Little information is available regarding sampling techniques, sample preservation and analytical methods.

Samples collected between 1994 and 1999 were obtained at the nearest tap, located prior to the water tank. All wells were equipped with electric pumps. These samples were split and an aliquot for determination of cations and metals was filtered through a $0.45\mu\text{m}$ -pore filter and acidified with ultrapure HNO_3 to $\text{pH} < 2.0$. The aliquot for anion determination was just stored at a temperature of 4°C until analysis.

Measurements of pH, Eh, EC, Cr^{6+} and alkalinity of water samples were performed in the field. Alkalinity analysis was performed by titration with H_2SO_4 using an end-point based on the Gran plot (Appelo and Postma 1993). Analytical methods for chemical species quantified in the sampled groundwater are summarized in Table II.

In performing chemical analyses, OA/OC proce-

analyses generally within $\pm 10\%$. Saturation indexes of calcite and dolomite were calculated using PhreeqC (Parkhurst and Appelo 1999).

RESULTS AND DISCUSSION

REGIONAL DISTRIBUTION OF TOTAL CHROMIUM AND Cr^{6+}

A general assessment of total and hexavalent chromium concentrations in groundwater was conducted in the Northwestern region of the State of São Paulo based on existing data gathered between 1977 and 1993 (Almodovar and Pacheco 1995). Total chromium is distributed in various municipalities and districts of the region. In some localities, concentrations are lower than or equal to the potability limit (0.05 mg.L^{-1}), whereas in many others concentrations vary from this limit up to 0.155 mg.L^{-1} .

Table III shows a summary of basic statistics regarding chromium concentrations detected in groundwater at various localities during this period. Figure 2 shows a regional map with indication of the concentrations of total and hexavalent chromium obtained in groundwater in the study region.

Around 39% of all available analyses showed concentrations of chromium higher than (or equal to) the potability limit. The values showed a great degree of variation between different localities in the region and a clear distribution pattern could not be obtained. However, comparing the concentrations found in wells installed in the Adamantina and Serra Geral aquifers, it was observed that the highest concentrations of total chromium occurred in groundwater from the Adamantina Aquifer. Concentrations varied between 0.005 and 0.155 mg.L^{-1} , with the highest concentrations in the municipality of Urânia, whereas concentrations varied from 0.006 to 0.018 mg.L^{-1} in basalts of the Serra Geral Aquifer.

NATURAL VERSUS ANTHROPIC SOURCES

Anthropic processes and natural sources can be considered as possibilities for assessing the origin of chromium concentrations in groundwater. The Urânia region, a small town of approximately 10 000 inhabitants, was

NATURAL OCCURRENCE OF HEXAVALENT CHROMIUM

TABLE I
Characteristics of groundwater sampling events from 1977 to 1999.

Sampling date/ Source	Locality	Type/Number	Parameters for <i>in situ</i> analysis	Parameters for laboratory analysis
1977-1993 (public water supply company)	Data from 144 wells in 54 cities	2 shallow wells 121 deep wells	Electrical Conductivity (EC)	Cr_{tot} , Cr^{6+} , NO_3^-
December 1994 (Almodovar and Pacheco 1995)	12 localities	1 shallow well 24 deep wells	Air and watPb ²⁺ , or temperatures, pH, EC, total dissolved solids	Cr^{6+}
March 1995 (Almodovar and Pacheco 1995)	Urânia Santa Salete	P2, P4 and P5 wells P1 well	Air and water temperatures, pH, EC, Eh, total dissolved solids	Na^+ , K^+ , SiO_2 , Cr_{tot} , Cr^{6+} , SO_4^{2-} , F^- , Fe_{tot} , Mn^{2+} , Al^{3+} , NO_3^- , HCO_3^-
January 1995 (Almodovar and Pacheco 1995)	Urânia	P9 well	Air and water temperatures, pH, EC, Eh, total dissolved solids	Cl^- , SO_4^{2-} , Ca^{2+} , Mg^{2+} , Na^+ , K^+ , Al^{3+} , Cr_{tot}
Four rounds of sampling (one in each season) from June 1998 to June 1999	Urânia	14 shallow hand-dug wells and 17 deep supply wells PP02 and PP04 for study of chromium concentrations with pumping time	Air and water temperatures, pH, EC, Eh, O ₂ , Cr ⁶⁺ , Alkalinity	Ag^+ , Al^{3+} , Ba^{2+} , Br^- , Ca^{2+} , Cd^{2+} , Cl^- , Cr_{tot} , Cu^{2+} , Fe^{3+} , F^- , K^+ , Mg^{2+} , Mn^{2+} , Na^+ , Ni^{2+} , NO_2^- , NO_3^- , Pb^{2+} , P_{tot} , SO_4^{2-} , Sr^{2+} , Zn^{2+}

TABLE II
Analytical methods used during sampling events
from 1994 to 1999.

Chemical species	Analytical method
Cr ⁶⁺	Colorimetry – Aquaquant 1.14402 MERCK
Al ³⁺ , Ag ⁺ , Ba ²⁺ , Ca ²⁺ , Cd ²⁺ , Cr _{tot} , Cu ²⁺ , Fe _{tot} , Mg ²⁺ , Mn ²⁺ , Ni ²⁺ , Pb ²⁺ , Sr ²⁺ , Zn ²⁺	Atomic Absorption Spectrophotometry
K ⁺ , Na ⁺	Flame photometry
Cl ⁻ , Br ⁻ , F ⁻ , SO ₄ ²⁻ , NO ₃ ⁻ , NO ₂ ⁻ , PO ₄ ³⁻	Ion Chromatography (DIONEX 2010i)

TABLE III
Concentrations of chromium detected in 144 wells from 1977 to 1993
(Almodovar and Pacheco 1995).

N Samples	Mean	Standard Error	Minimum	Maximum
	mg.L ⁻¹			






Fig. 2 – Distribution of chromium concentrations in space (modified from Almodovar and Pacheco 1995).

entail or agricultural areas. Agriculture is the main economic activity in the area, primarily fruit cultivation (grape and pineapple) and small-scale cattle farming. Thus, the application of fertilizers and pesticides could possibly be a significant diffuse source of the chromium found in the aquifer. The chemical composition of fertilizers and pesticides commonly used in the Urânia area was then checked and no significant concentration of chromium was present.

Point sources of contamination, like steelworks, electroplating, leather tanning and chemical manufacturing, were not found to be present in the study area. Only small industries are present, such as rice processing and furniture manufacturing, which do not show an associated contaminant load of chromium. Besides this, these types of point sources of contamination would generate contamination plumes with limited and defined geo-

of chromium contamination in the study area, which reinforced the hypothesis of natural origin in the case of this contamination. This hypothesis was confirmed by Marcolan and Bertolo (2007), through chemical and mineralogical analysis using X-ray fluorescence, X-ray diffraction and scanning electron microscopy (SEM-EDS) on borehole samples collected from the top to the base of the aquifer in Urânia. The results of this investigation indicated that:

- (i) the concentration of chromium in sandstones is anomalous, with average and maximum values of 221 and 336 ppm, respectively, which are significantly higher than the reported average concentration in a generic sandstone (35 mg/kg) and soils from another regions in the State of São Paulo (26.3 mg/kg);



NATURAL OCCURRENCE OF HEXAVALENT CHROMIUM

- (iii) chromium concentrations in clay minerals and iron hydroxides are also relatively high (up to 650 ppm).

HYDROGEOCHEMICAL STUDY IN TOWN OF URÂNIA

Conceptual model

The geological and hydrogeological features of the Adamantina Aquifer in Urânia are quite similar to those described for the aquifer regionally. The aquifer is unconfined to semi-confined, constituted of well-sorted fine to very fine sandstones with carbonate cementation, relatively homogeneous and isotropic in the scale of study, and with a thickness varying from 66 to 165 m. The urban area of Urânia is located on the crest of a smooth hill, which corresponds to the local aquifer groundwater divide.

Groundwater flow lines that originate in this area converge to the Comprido Creek, which is considered as the main discharge area of the aquifer locally (Figure 3). The hydraulic gradient varies from 0.011 to 0.025 and the advective flow velocity of groundwater ranges from 20 to 400 m/year. Pumping tests showed a hydraulic conductivity value of $8.54\text{E-}6$ m/sec (F.A. Cagnon, unpublished data).

Three classes of supply wells were registered in Urânia, each representing conditions at a specific depth in the aquifer: private hand-dug wells (14), private 4" wells (10) and public 6" wells (7). In general, the private hand-dug wells (shallow wells) have a diameter of 1.2 m and reach an average depth of 11 m, being characteristic of the shallow part of the aquifer, where a local groundwater flow system with a shorter transit time predominates. The private 4" wells (intermediate wells) pump groundwater from intermediate depths in the aquifer, show an average depth of 57 m and are commonly cased with 4" tubes to a depth of 20 m. The public 6" wells (deep wells) extend to depths varying from 75 to 160 m and usually reach the top of basalts that comprise the Serra Geral Formation. These wells are continuously cased with 6" tubes and filters that are installed from the intermediate to deep intervals of the aquifer. This situation might result in mixture of water from these layers. A regional system of groundwater

indicate the occurrence of downward flow potential in the urban area and upward flow potential in the rural area. The Comprido Creek, which is confirmed by near the PP05, where results showed conditions of artesian flow controlled by topography.

Three hydrochemical zones were identified in the Adamantina Aquifer in Urânia that are in accordance with the depths of the studied wells. The first hydrochemical zone is the Shallow Zone (up to 30 m), associated with the private hand-dug wells (shallow wells) exhibiting Na-Ca-Cl- NO_3 facies (Figure 3). The intermediate Zone (30–70 m) is associated with the private 4" wells (intermediate wells) and shows Ca- HCO_3 facies. The Deep Zone (> 70 m) is associated with the public 6" wells (deep wells) and shows Na-Ca- HCO_3 facies.

Table IV shows the results of a statistical analysis performed on the analytical results of water samples collected from shallow, intermediate and deep wells during four rounds of monitoring activities performed between June 1998 and June 1999. Elevated values and standard deviation were advective obtained for Cr^{6+} , Fe_{total} , Mn^{2+} and SO_4^{2-} , indicating that the related mean value is not representative for the aquifer. In general, an anomalous high concentration of these parameters detected in a sample resulted in a tendency for elevation of the mean value.

Table IV also shows the average chemical composition of water from well PP04, which may better represent the hydrochemical conditions existing in the deep aquifer below intermediate depths, as it is solely screened from the base of the aquifer. As mentioned previously, the public 6" wells are frequently screened from intermediate depths to the base of the aquifer. As a result, the concentrations shown for water from these wells may in fact be considered as representative of a mixture of water from these depths.

From shallow to deep groundwater zones, the results showed an increase in pH and HCO_3^- concentrations and a decrease in electrical conductivity (Eh) indicating a drop in salinity. Eh also decreased, but it indicates oxidizing conditions for the aquifer. Comparing the three groundwater categories, concentrations





Fig. 3 – Conceptual hydrogeological and hydrogeochemical model of the Adamantina Aquifer in Urânia.


centrations of HCO_3^- , Na^+ , F^- , and Cr^{6+} are higher in the deep aquifer.

Comparing the mean values obtained for water samples collected in the different aquifer zones, NO_3^- and Cl^- concentrations in shallow groundwater are 5 and 8 times higher than in deep groundwater, respectively. At shallow depths, nitrate concentrations exceed the drinking-water potability value of 45 mg.L^{-1} and gradually decrease with depth. The occurrence of nitrate in the proximity of the surface can be attributed to the organic charge of septic systems and/or fertilizer use. Its decrease, together with the decrease of EC (salinity), Cl^- and Na^+ , can be attributed to contaminant dispersion and dilution processes. The sample collected from well PP04, which better represents hydrochemical conditions

such parameters as pH, HCO_3^- , Na^+ , Ca^{2+} and Mg^{2+} . Increase in pH and HCO_3^- concentrations from the top to the base of the aquifer controls the solubility of calcite and dolomite minerals (reaction 1), which exhibit increasing saturation values, until their saturation at the base of the aquifer (Table IV).



In addition, there is substantial enrichment of Na^+ in the aquifer relative to Cl^- . The shallow and intermediate depths of the aquifer show a Na/Cl molar ratio of 1.2, which increases to 46.3 in the deep aquifer, thus indicating that the rock is providing the water with Na^+ . The increase in Na^+ , followed by a decrease in Ca^{2+} and Mg^{2+} concentrations and $\text{Ca}^{2+}/\text{HCO}_3^-$ molar ratios, sug-



NATURAL OCCURRENCE OF HEXAVALENT CHROMIUM

TABLE IV
Summary of analytical data of groundwater in shallow, intermediate and deep wells studied in the municipality of Urânia
(four rounds of sampling from June 1998 to June 1999).

N wells		Shallow wells				Intermediate wells				Deep wells			
		14				10				7			
Parameter	Unit	Mean	Std.	Min.	Max.	Mean	Std.	Min.	Max.	Mean	Std.	Min.	Max.
Depth	m	11.4	4.5	5.2	18.0	57.1	17.5	40.0	100.0	130.6	30.8	75.0	160.0
Temperature	°C	25.5	3.0	19.9	30.5	26.3	1.9	22.9	30.1	25.9	2.6	21.5	29.7
Conductivity	μS/cm	273.8	100.5	130.0	499.0	205.0	59.6	93.3	272.0	185.1	49.2	119.7	268.5
Ph		5.78	0.43	5.03	6.45	6.55	0.54	5.42	7.08	7.40	1.03	6.09	8.90
Eh	mV	525.3	29.9	460.7	569.0	494.6	12.3	474.0	513.7	474.0	26.7	443.0	525.3
O ₂	mg.L ⁻¹	3.45	1.12	1.54	5.60	4.52	1.11	2.01	6.31	3.79	0.54	3.19	4.59
Cr total	mg.L ⁻¹	<i>0.006</i>	<i>0.003</i>	<i><0.005</i>	<i>0.013</i>	0.013	0.006	0.005	0.025	<i>0.051</i>	<i>0.045</i>	<i>0.013</i>	<i>0.12</i>
Cr ⁶⁺	mg.L ⁻¹	<i>0.002</i>	<i>0.003</i>	<i><0.005</i>	<i>0.010</i>	<i>0.013</i>	<i>0.008</i>	<i>0.005</i>	<i>0.025</i>	<i>0.044</i>	<i>0.042</i>	<i>0.010</i>	<i>0.12</i>
Na ⁺	mg.L ⁻¹	18.29	10.20	3.60	31.00	5.35	3.130	2.60	13.30	12.42	12.61	1.98	32.00
K ⁺	mg.L ⁻¹	5.94	1.76	2.30	8.90	3.90	1.140	2.60	6.27	3.170	2.55	0.30	7.77
Ca ²⁺	mg.L ⁻¹	17.73	9.82	6.52	40.50	21.04	9.620	7.73	38.05	15.64	10.55	4.30	35.20
Mg ²⁺	mg.L ⁻¹	6.36	3.09	1.30	13.65	7.34	2.600	3.25	11.00	4.46	3.16	0.35	9.20
Ba ²⁺	mg.L ⁻¹	0.30	0.16	0.08	0.77	0.22	0.100	0.08	0.42	0.15	0.10	0.03	0.35
Sr ²⁺	mg.L ⁻¹	0.16	0.08	0.06	0.33	0.16	0.090	0.03	0.31	0.25	0.21	0.07	0.58
HCO ₃ ⁻	mg.L ⁻¹	23.31	14.90	6.10	51.85	85.30	58.90	15.25	183.0	90.52	40.99	40.41	154.0
CO ₃ ⁻	mg.L ⁻¹	—	—	—	—	—	—	—	—	—	—	9.00	15.00
Cl ⁻	mg.L ⁻¹	22.19	13.06	5.73	45.58	<i>6.65</i>	<i>6.470</i>	<i>0.36</i>	<i>22.12</i>	2.66	1.39	0.89	4.09
SO ₄ ²⁻	mg.L ⁻¹	<i>0.69</i>	<i>0.67</i>	<i>0.180</i>	<i>2.46</i>	<i>0.28</i>	<i>0.512</i>	<i>0.01</i>	<i>1.72</i>	0.16	0.075	0.08	0.26
NO ₃ ⁻	mg.L ⁻¹	78.79	31.80	37.29	161.8	34.73	24.54	1.46	69.18	15.14	12.74	0.08	35.15
PO ₄ ³⁻	mg.L ⁻¹	0.08	0.04	0.04	0.18	0.12	0.100	0.01	0.30	<i>0.11</i>	<i>0.11</i>	<i>0.03</i>	<i>0.36</i>
F ⁻	mg.L ⁻¹	0.08	0.04	0.03	0.16	0.12	0.060	0.03	0.20	0.14	0.035	0.08	0.19
Fe total	mg.L ⁻¹	<i>0.03</i>	<i>0.03</i>	<i><0.01</i>	<i>0.09</i>	<i>0.02</i>	<i>0.030</i>	<i><0.01</i>	<i>0.07</i>	<i>0.025</i>	<i>0.017</i>	<i>0.01</i>	<i>0.05</i>
Mn ²⁺	mg.L ⁻¹	<i>0.07</i>	<i>0.09</i>	<i>0.01</i>	<i>0.31</i>	<i>0.01</i>	<i>0.010</i>	<i><0.01</i>	<i>0.04</i>	<i>0.01</i>	<i>0.02</i>	<i><0.01</i>	<i>0.05</i>
Sat.Ind.** Calcite		-3.09				-1.65							
Sat.Ind. Dolomite		-6.09				-3.23							

(1) N wells = number of wells; Std.: standard deviation; Min.: minimum; Max.: maximum; Sat.Ind. = saturation index. (2) Concentrations in *italic* and lower case highlight elevated values of standard deviation, indicating that the mean value is not a representative figure.

Reaction 2 decreases Ca²⁺ concentration and seems to control the dissolution of carbonate minerals, driving reaction 1 to the right, increasing the pH and HCO₃⁻ concentration. Table V summarizes the hydrogeochemical processes identified in the aquifer.

Behavior of chromium concentrations

Chromium concentrations also showed distinct behavior at different depths in the aquifer (Fig. 4), with an increase from the shallow to deep zones (Table IV). Ten of the 14 shallow wells showed non-detected concentra-

at low concentrations, ranging from the detection limit to 0.025 mg.L⁻¹. All deep wells showed the detection limit of chromium at concentrations that were generally higher than those found at intermediate depths. Three of the 7 samples showed concentrations exceeding the pollution limit (0.05 mg.L⁻¹).

Cr_{total} and Cr⁶⁺ concentrations are usually low (Table IV), except in the case of the shallow wells, indicating that dissolved chromium in groundwater is essentially present in its hexavalent form at the intermediate and deep aquifer levels. This is in accordance with the oxidizing environment and elevated pH conditions.




TABLE V
Summary of hydrogeochemical processes identified in the 3 different hydraulic zones studied.

	Shallow aquifer	Intermediate aquifer	Deep aquifer
Environment	Acidic waters under oxidizing conditions	Acidic to neutral waters under oxidizing conditions	Alkaline waters under oxidizing conditions
Water type	Na-Ca-Cl-NO ₃	Ca-HCO ₃	Na-Ca-HCO ₃
Cations	Na ⁺ and Ca ²⁺ > Mg ²⁺ and K ⁺	↑ Ca ²⁺ and Mg ²⁺ ↓ Na ⁺ and K ⁺	↑ Na ⁺ ↓ K ⁺ , Ca ²⁺ e Mg ²⁺
Anions	NO ₃ ⁻ > HCO ₃ ⁻ and Cl ⁻	↑ HCO ₃ ⁻ ↓ NO ₃ ⁻ and Cl ⁻	↑ HCO ₃ ⁻ ↓↓ NO ₃ ⁻ and Cl ⁻
Molar ratios (meq.L ⁻¹ /meq.L ⁻¹)	Na ⁺ /Cl ⁻ = 1.2 Ca ²⁺ /HCO ₃ ⁻ = 2.3	Na ⁺ /Cl ⁻ = 1.2 Ca ²⁺ /HCO ₃ ⁻ = 0.7	Na ⁺ /Cl ⁻ = 46.3 Ca ²⁺ /HCO ₃ ⁻ = 0.4
Geochemical processes – Major constituents	Low pH and most of the Na ⁺ , NO ₃ ⁻ and Cl ⁻ concentrations are attributed to contamination by septic systems and/or fertilizer use. Conditions for dissolution of carbonate minerals.	Hydraulic dispersion of the contaminants. Dissolution of carbonate minerals, increasing pH and Ca ²⁺ and HCO ₃ ⁻ concentrations.	Cation exchange of Na ⁺ by Ca ²⁺ and Mg ²⁺ . Dissolution of carbonate minerals up to their solubility limits. Dissolution of carbonate minerals affected by cation exchange reaction, increasing pH and HCO ₃ ⁻ concentrations.
Chromium	Low to very low concentrations (usually non-detect); Cr ³⁺ predominant. Mn ²⁺ usually detected.	Low concentrations (max. 0.025 mg.L ⁻¹); Cr ⁶⁺ predominant.	Moderate to high concentrations (max. 0.120 mg.L ⁻¹); Cr ⁶⁺ predominant; (+) correlation with pH, Na ⁺ and total alkalinity; (-) correlation with NO ₃ ⁻ . Mn ²⁺ usually not detected.
Possible geochemical processes – Chromium	Pyroxene weathering; Cr ³⁺ cations predominate in the form of hydroxides and/or adsorbed onto clay/hydroxides minerals; Possible reduction of Cr ⁶⁺ into Cr ³⁺ by organic matter, NH ₄ ⁺ , NO ₂ ⁻ and/or Fe ²⁺ .	Possible pyroxene weathering; Oxidation of Cr ³⁺ to Cr ⁶⁺ by MnO ₂ ; Adsorption of Cr ⁶⁺ anions by Al and Fe hydroxides.	Oxidation of Cr ³⁺ to Cr ⁶⁺ by MnO ₂ ; Increased pH and presence of competing anions cause desorption and mobilization of Cr ⁶⁺ anions.

↑ or ↓: elevation or decrease of concentrations in respect to the previous shallower aquifer level.

Correlation coefficients between chromium and the other chemical components were obtained. Except for significant correlations of Cr with pH ($R = 0.80$) and NO₃⁻ ($R = -0.61$) (Fig. 5), no statistically linear or inverse relationships were found in comparing chromium with other chemical species or physical parameters. As suggested in Figure 4, pH, Cr⁶⁺ and NO₃⁻ box plots showed different patterns between shallow, intermediate and deep wells.

a correlation between Cr⁶⁺ and pH, NO₃⁻, Na⁺ and total alkalinity has been observed (Fig. 5). The highest concentrations of chromium were observed in three deep wells (PP04 included), where the highest values of pH, Na⁺ and total alkalinity were found, as well as the lowest concentrations of NO₃⁻.

Cr⁶⁺ and Mn²⁺ concentrations showed contrasting tendencies, although no statistical correlation was found between them. Mn²⁺ was detected in all 14 shallow wells.





Fig. 4 – Box plots showing a comparison of chromium, pH and nitrate distribution for groundwater samples from shallow, intermediate and deep wells.

Sampling throughout different pumping intervals was performed over a 24-hours period with a constant flow rate in the public 6" wells (deep wells) PP02 and PP04. Samples were collected at 15 min, and then at 1, 6, 12, 18 and 24 h after the start of pumping. The results indicate that pH values and Cr^{6+} and Na^+ concentrations increase in both deep wells over the pumping period, whereas NO_3^- concentrations decrease (Fig. 6).

Given that the characteristics of the aquifer are considered to be of the unconfined to semiconfined type, continuous pumping allows for the capture of deeper flows in the aquifer, represented by older waters that have been subject to a longer period of water-rock interaction. The observed relationships between Cr^{6+} , pH, Na^+ and NO_3^- during this experiment are in accordance with the correlations observed in Figure 5.

Possible geochemical reactions involving chromium

Aqueous concentrations of chromium are dependent

transformation of $\text{Cr}^{3+}/\text{Cr}^{6+}$ and adsorption/desorption of $\text{Cr}^{3+}/\text{Cr}^{6+}$ onto mineral surfaces.

According to Richard and Bourg (1991), Chromium minerals are very scarce in nature and amorphous $[\text{Cr}^{3+}]$ hydroxide is probably the main phase controlling chromium solubility in natural environments. However, the chemical and mineralogical analyses conducted during this investigation indicated high concentrations of chromium in sandstones (average of 221 ppm), and especially in pyroxene (augite) crystals (6000 ppm), which is one of the main heavy minerals found in the sandstone. Pyroxene is one of the most reactive minerals in weathering processes, according to the Goldich stability series (Kehew 2001), thus, this mineral may be considered as the main source of chromium found in water. However, chromium occurs as Cr^{3+} in pyroxenes and a redox reaction must take place to make Cr^{6+} available and mobile in groundwater.

The redox process for transformation of Cr^{3+} to Cr^{6+} can take place in the presence of another







Fig. 5 – Relationship between hexavalent chromium concentrations and pH, nitrate, alkalinity and sodium concentrations.

of Cr^{3+} into Cr^{6+} at pH values greater than 9 (Fendorf and Zasoski 1992), however, according to Richard and Bourg (1991), the rate of oxidation is very slow and enables Cr^{3+} to be involved in faster concurrent reactions, like sorption or precipitation. Considering that the residence time of water in aquifers may reach thousands of years and pH values are quite elevated in the deep aquifer, the possibility cannot be neglected that this reaction may occur.

the oxidation of Cr^{3+} to Cr^{6+} in aquifers (Apte et al. 2006, Sedlak and Chan 1997, Richard and Bourg 1991, Eary and Rai 1987, Fendorf and Zasoski 1992, Fendorf 1995, Bartlett and James 1979).



This reaction is more effective under acidic conditions. According to this reaction, one might expect that Cr^{6+}



NATURAL OCCURRENCE OF HEXAVALENT CHROMIUM





Fig. 6 – Variation of pH values and chromium, nitrate and sodium concentrations throughout pumping period.

the deep aquifer, where Cr^{6+} concentrations are higher. However, any Mn^{2+} formed during Cr^{3+} oxidation will be converted back to MnO_2 in the presence of dissolved oxygen (Apte et al. 2006), which is observed in all depths in the aquifer in Urânia. In fact, chromium and manganese form a pair of chemical elements of opposite tendencies, since, under oxidizing conditions, Cr^{6+} is soluble as CrO_4^{2-} and Mn^{4+} is present as MnO_2 ; under reducing conditions, Cr^{3+} is removed from solution as $\text{Cr}(\text{OH})_3$ and Mn^{2+} is soluble (Richard and Bourg 1991).

In the shallow aquifer, more acidified water will contribute to more intensive weathering of pyroxene. Cr^{3+} rapidly precipitates as hydroxides or is adsorbed by Fe and Mn hydroxides (Richard and Bourg 1991). Part of the possible Cr^{6+} produced in the shallow aquifer might be reduced back to Cr^{3+} by organic matter (humic acids and humic or fulvic acids) or by Fe^{2+} oxidation, which are fast reactions. Nitrification processes, which convert ammonium to nitrate, acidify waters and also contribute to the immobilization of Cr^{3+} (Richard and Bourg 1991).



immobilized by adsorption to clay minerals (low pH_{zpc} minerals), a process that increases with pH (Richard and Bourg 1991).

Due to its anionic nature, Cr^{6+} is preferentially retained on positively charged surfaces, for example, Al and Fe hydroxides (high pH_{zpc} minerals), principally in more acidic conditions. In addition, most of the Cr^{6+} produced by Cr^{3+} oxidation reactions may be adsorbed by these minerals in the intermediate and deep portions of the aquifer.

However, adsorption of Cr^{6+} to minerals strongly decreases in the alkaline pH range and in the presence of other competing anions, increasing its mobility (Rai et al. 1989, Zachara et al. 1987). This may be the most probable phenomenon occurring in the deep aquifer, since the anomalous increase in pH, probably driven by cation exchange reactions involving Na^+ , Ca^{2+} and Mg^{2+} and by the dissolution of carbonate minerals up to their solubility limits (reactions 1 and 2), is likely causing the desorption of Cr^{6+} anions to the aquifer. Table V summarizes possible geochemical processes involving chromium in the different parts of the aquifer.

CONCLUSIONS

Chromium is found in groundwater of the Adamantina Aquifer, at concentrations often exceeding the established potability limit (0.05 mg.L^{-1}), in an extensive area covering at least 54 municipalities located in the north-western region of the State of São Paulo. An investigation conducted in the municipality of Urânia, where the highest concentrations were found, indicated that the origin of this contamination is natural, since elevated concentrations of chromium were detected in sandstones (average of 221 ppm) and especially in the mineral augite (6000 ppm), which is the main heavy mineral found in the sandstone. Besides this, no possible diffuse or point sources of contamination were observed in the study area. The results obtained for Urânia are good indication that the chromium found in groundwater regionally may have the same origin.

The distribution of chromium concentrations in the different hydrochemical and hydraulic zones of the

and hydrochemical conditions allow chromium release into aqueous solution. In most of the groundwater samples collected from the shallow wells, chromium has not been detected or observed concentrations are very low. The distribution pattern of these concentrations also reinforces the hypothesis that the origin of chromium is natural. Dissolved chromium is essentially present in its hexavalent form, especially at the intermediate and deep levels of the aquifer.


The origin of Cr^{6+} seems to be associated with the weathering of Cr-minerals, such as the Cr-augite identified, followed by the redox transformation of Cr^{3+} into Cr^{6+} by another redox couple, for example, $\text{H}_2\text{O}/\text{O}_2(\text{aq.})$ or $\text{Mn}^{2+}/\text{Mn}^{4+}$. According to many authors, manganese oxides are most likely responsible for Cr^{3+} oxidation. The processes of adsorption/desorption of Cr^{6+} probably represent important reactions, since the highest Cr^{6+} concentrations are found in the deep aquifer, where groundwater pH is more alkaline, which is a condition necessary for the desorption of Cr^{6+} . This phenomenon may be occurring in the deep aquifer, where cation exchange reactions are driving the dissolution of carbonate minerals, causing an anomalous increase in pH. Complementary field and laboratory studies are now being conducted in order to confirm the occurrence of such geochemical processes responsible for the presence of chromium in groundwater.

ACKNOWLEDGMENTS

The authors are indebted to Fundação de Amparo à Pesquisa do Estado de São Paulo (FAPESP) for financial support of the project (processes 97/13879-6, 98/15340-0 and 05/60502-3). Bourotte has received a post-doctoral grant from Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq) (No. 154492/2006-0).

RESUMO

Concentrações anômalas de cromo hexavalente foram detectadas em águas subterrâneas do Aquífero Adamantina em pelo menos 54 municipalidades localizadas na região noroeste do Estado de São Paulo, sudeste do Brasil, algumas vezes ultrapassando o limite máximo permitido para consumo humano




NATURAL OCCURRENCE OF HEXAVALENT CHROMIUM

como natural, pois foram detectadas concentrações de cromo nos arenitos do aquífero (média 221 ppm) e em piroxênios (6000 ppm), um dos principais minerais pesados encontrados nos sedimentos. Além disso, não foram observadas outras possíveis fontes de contaminação difusas ou pontuais na área de estudo. Uma estratificação da qualidade da água no aquífero foi observada e as mais elevadas concentrações de Cr^{6+} foram detectadas na base do aquífero ($0,12 \text{ mg.L}^{-1}$), onde as águas subterrâneas apresentam valores elevados de potencial redox ($E_h = 472,5 \text{ mV}$) e de pH (8,61). A origem do Cr^{6+} na água deve estar associada com o intemperismo do piroxênio (augita), seguida da oxidação do Cr^{3+} pelos óxidos de manganês. As maiores concentrações de Cr^{6+} estão provavelmente ligadas com reações de desorção, devido ao pH anormalmente alcalino da água subterrânea na base do aquífero.

Palavras-chave: cromo, água subterrânea, aquífero, contaminação natural.

REFERENCES

- ALMODOVAR MLN. 2000. A origem natural da poluição por cromo no Aquífero Adamantina, município de Urânia, São Paulo, (In Portuguese – The origin of natural pollution by chromium in the Adamantina Aquifer, Urânia municipality, São Paulo), Ph.D. thesis, Institute of Geosciences, University of São Paulo, 199 p.
- ALMODOVAR MLN AND PACHECO A. 1995. Anomalia de Cromo nas Águas Subterrâneas de Urânia no Noroeste do Estado de São Paulo. Boletim IG-USP: Série Científica 26: 1–12.
- APPELO CAJ AND POSTMA D. 1993. Geochemistry, Ground-water and Pollution. Taylor and Francis Books, London, 649 p.
- APTE AD, TARE V AND BOSE P. 2006. Extent of oxidation of Cr(III) to Cr(VI) under various conditions pertaining to natural environment. J Hazard Mater B128: 164–174.
- ATSDR. 2000. Toxicological profile for Chromium. Agency for Toxic Substances and Disease Registry, U.S. Department of Health and Human Services, 461 p.
- BALL JW AND IZBICKI JA. 2004. Occurrence of hexavalent chromium in ground water in the wester Mojave Desert, California. Appl Geochem 19: 1123–1135.
- BARTLETT RJ AND JAMES B. 1979. Behavior of chromium in soils: III. Oxidation. J Environ Qual 8: 31–35.
- CHARLET L AND MANCEAU AA. 1992. X-Ray absorption spectroscopic study of the sorption of Cr^{3+} at the water interface II. Adsorption, coprecipitation and precipitation on hydrous ferric oxide. J Colloid Interface Sci 148: 443–458.
- DAEE – DEPARTAMENTO DE ÁGUAS E ENERGIA ELÉTRICA. 1976. Estudo de Águas Subterrâneas. Região Administrativa 7, 8 e 9, Bauru, São José do Rio Preto e Araraquã. 4v. São Paulo, Brasil.
- EARY LE AND RAI D. 1987. Kinetics of Chromium(III) oxidation to Chromium(VI) by reaction with manganese dioxide. Environ Sci Technol 21: 1187–1193.
- FENDORF SE. 1995. Surface reactions of chromium in soils and waters. Geoderma 67: 55–71.
- FENDORF SE AND ZASOSKI RJ. 1992. Chromium(III) oxidation by $\delta\text{-MnO}_2$ 1. Characterization. Environ Sci Technol 26: 79–85.
- FERNANDES LA AND COIMBRA AM. 2000. Revisão geológica da parte oriental da Bacia Bauru (Neoproterozoico). Rev bras Geociencia 30: 723–734.
- GONZALEZ AR, NDUNDU K AND FLEGAL AR. 2000. Natural occurrence of hexavalent chromium in the Red Sands Aquifer, California. Environ Sci Technol 34: 5505–5511.
- HIRATA R, BASTOS C. AND ROCHA G. 1997. Mapa de vulnerabilidade a poluição dos aquíferos do Estado de São Paulo. Secretaria do Meio Ambiente do Estado de São Paulo. São Paulo, 2v.
- KEHEW AE. 2001. Applied chemical hydrogeology. Prentice-Hall. 368 p.
- MARCOLAN L AND BERTOLO R. 2007. Anomalias geoquímicas de Cromo em Arenitos da Formação Adamantina em Urânia – SP. XI Congresso Brasileiro de Geociências, São Paulo, 4 p.
- PARKHURST DL AND APPELO AAJ. 1999. User's guide to PHREEQC (version 2) – A computer program for geochemical reaction, batch-reaction, one dimensional transport and inverse geochemical modeling. US Geol. Survey Water Resource Invest.
- RAI D, EARY LE AND ZACHARA JM. 1989. Environmental geochemistry of chromium. Sci Total Environ 86: 15–25.
- RICHARD FC AND BOURG ACM. 1991. Aqueous geochemistry of chromium: a review. Water Res 25: 807–815.



- ROBLES-CAMACHO J AND ARMIENTA MA. 2000. Natural chromium contamination of groundwater at León Valley, México. *J Geochem Explor* 68: 167–181.
- SEDLAK DL AND CHAN PG. 1997. Reduction of hexavalent chromium by ferrous iron. *Geochim Cosmochim Acta* 61: 2185–2192.
- SOARES PC, LANDIM PMB, FÚLFARO VJ AND SOBREIRO NETO AF. 1980. Ensaios de caracterização estratigráfica do Cretáceo no Estado de São Paulo: Grupo Bauru. *Rev bras Geocienc* 10: 177–185.
- ZACHARA JM, GIRVIN DC, SCHMIDT RL AND RESCH CT. 1987. Chromate adsorption on amorphous iron oxyhydroxide in the presence of major groundwater ions. *Environ Sci Technol* 21: 589–594.