



Anais da Academia Brasileira de Ciências

ISSN: 0001-3765

[aabc@abc.org.br](mailto:aabc@abc.org.br)

Academia Brasileira de Ciências

Brasil

MAIA, EMANUEL; SIQUEIRA, DALMO L.; SALOMÃO, LUIZ C.C.; PETERNELLI, LUIZ A.;  
VENTRELLA, MARÍLIA C.; CAVATTE, RITHIELY P.Q.

Development of the banana plants 'Prata Anã' and 'FHIA-01' under the effect of paclobutrazol applied  
on the soil

Anais da Academia Brasileira de Ciências, vol. 81, núm. 2, junio, 2009, pp. 257-263  
Academia Brasileira de Ciências  
Rio de Janeiro, Brasil

Available in: <http://www.redalyc.org/articulo.oa?id=32713477011>

- How to cite
- Complete issue
- More information about this article
- Journal's homepage in [redalyc.org](http://redalyc.org)

[redalyc.org](http://redalyc.org)

Scientific Information System

Network of Scientific Journals from Latin America, the Caribbean, Spain and Portugal

Non-profit academic project, developed under the open access initiative



Anais da Academia Brasileira de Ciências (2009) 81(2): 257-263  
(Annals of the Brazilian Academy of Sciences)  
ISSN 0001-3765  
www.scielo.br/aabc

## Development of the banana plants ‘Prata Anã’ and ‘FHIA-01’ under the effect of paclobutrazol applied on the soil

EMANUEL MAIA<sup>1</sup>, DALMO L. SIQUEIRA<sup>1</sup>, LUIZ C.C. SALOMÃO<sup>1</sup>, LUIZ A. PETERNELLI<sup>2</sup>,  
MARÍLIA C. VENTRELLA<sup>3</sup> and RITHIELY P.Q. CAVATTE<sup>3</sup>

<sup>1</sup>Departamento de Fitotecnia, Setor de Fruticultura, Universidade Federal de Viçosa (UFV)  
Av. P.H. Rolfs s/n, Centro, 36570-000 Viçosa, MG, Brasil

<sup>2</sup>Departamento de Informática, Setor de Estatística, UFV, Av. P.H. Rolfs s/n, Centro, 36570-000 Viçosa, MG, Brasil

<sup>3</sup>Departamento de Biologia Vegetal, UFV, Av. P.H. Rolfs s/n, Centro, 36570-000 Viçosa, MG, Brasil

*Manuscript received on April 3, 2008; accepted for publication on September 9, 2008;  
presented by ELIBIO L. RECH*

### ABSTRACT

Under some cultivation conditions, excessive growth of the pseudostem of banana plants can be considered a limiting factor, and thus, the use of growth regulators can constitute a valid alternative. This work aimed to evaluate the action of paclobutrazol on the growth of the pseudostem of banana plants and other characteristics of their development. An experiment was installed with five paclobutrazol doses (0; 0.5; 1.0; 1.5; and 2.0 g of a.i. plant<sup>-1</sup>) combined with two banana plant cultivars (‘Prata Anã’ and ‘FHIA-01’). Paclobutrazol did not affect the number of days from planting to flowering, total leaf area and leaf emission rate. However, it increased leaf permanence on the plants and sucker number, reduced the foliar area of the leaves emitted after its application and provided an average reduction of pseudostem height of 26%.

**Key words:** foliar area, growth analysis, growth regulator, *Musa* spp., suckers.

### INTRODUCTION

Brazilian banana plant cultivation has undergone changes in the past years mainly related to seedling production due to two strong reasons: the advance of pest and disease into free areas and the formation of large export-oriented orchards. This scenario led to the production of a large amount of seedlings with high phytosanitary standard within a short period of time, made possible by the use of micro-propagation techniques (Leonel et al. 2004). For these seedlings to be produced, the matrix plants must be healthy and maintained under conditions that minimize the possibility of con-

tamination, what prompted some propagation laboratories to opt for maintaining their matrices under protected environment.

A common problem of banana plants cultivated under protected environment is the excessive growth of their pseudostem, which can reach more than six meters for some cultivars, requiring more planting space and causing, in some cases, the rupture of the plastic film surrounding the roof (El Otmani et al. 1992, Jeyakumar et al. 2003). Thus, the use of growth regulators to reduce plant height may be a solution for plants cultivated under protected environment, as shown by promising studies on banana plant growth regulation (El Otmani et al. 1992, Jeyakumar et al. 2003).



Mouco and Albuquerque 2005). The main plant responses observed after paclobutrazol application are: smaller length of the new shoots, green color intensification, increased chlorophyll concentration and reduced foliar area (Khurshid et al. 1997, Yim et al. 1997, Carvalho et al. 2005, Mouco and Albuquerque 2005). ‘Grande Naine’ banana plants showed reduced pseudostem and foliar area growth, increased chlorophyll concentration and no change in cycle duration and plant productivity during the first production cycle (El Otmani et al. 1992). Thus, this work proposes to evaluate the effects of paclobutrazol on some development characteristics of the banana plants ‘Prata Anã’ and ‘FHIA-01’ during the first production cycle.

#### MATERIALS AND METHODS

The experiment was conducted in the Fruit Culture Sector of the Universidade Federal de Viçosa (UFV), Viçosa, MG. The municipality of Viçosa is located in Zona da Mata Mineira (20°45’20” S and 42°52’40” W), at 651 meters of altitude in relation to the sea level. The experiment was conducted from August 2005 to December 2006. The treatments used were five paclobutrazol doses (0; 0.5; 1.0; 1.5; and 2.0 g of a.i. plant<sup>-1</sup>) combined with two banana plant cultivars (‘Prata Anã’ and ‘FHIA-01’). Cv. Prata Anã (AAB) was selected because it ranks among the cultivars with the largest planted area, and are well accepted in the domestic market (Leonel et al. 2004), besides belonging to the type ‘Prata’, which accounts for approximately 60% of the area cultivated in Brazil (Silva et al. 2002). The tetraploid cv. FHIA-01 (AAAB) was selected for presenting characteristics similar to those of cv. Prata Anã and for being resistant to Black Sigatoka (*Mycosphaerella fijiensis*) and other important diseases, such as Panama disorder (*Fusarium oxysporum* f.sp. cubense) (Alves 1999). The seedlings used for the implantation of the experiment came from the Cell and Plant Tissue Culture Laboratory of the Department of Plant Sciences of the UFV.

The experiment was installed in the field on November 3, 2005 in a factorial scheme and a complete randomized block design (Steel et al. 1997) with eight repe-

classes. Paclobutrazol (Pachlobutrazol 100 CE, 10% i.a., Wiser) was applied when the plants reached an average height of 75 cm considering the distance between soil and the second leaf of the foliar rosette on March 6, 2006. The product was applied on the soil surface, with humidity close to field capacity, around the pseudostem at a mean distance of 25 cm. The plants received water complementation during the most severe hydric deficit periods in the region. The cultural practices followed the current recommendations (Alves 1999).

Periodic evaluations consisted of: pseudostem height (cm), measurement from the soil up to second leaf of the foliar rosette; pseudostem circumference (cm), measurement at 0.30 m height from the soil surface; number of totally expanded leaves in the plant at the moment of evaluation; leaf emission rate and foliar area, using the non-destructive method (Turner 2003). These evaluations occurred between 11/13/2005 and 09/28/2006. Every three months after paclobutrazol application, thinning (sucker removal) and dead leaf removal were performed, followed by sucker number and fresh mass evaluation. Mean sucker fresh mass per plant and total number of suckers produced by the plants during the period evaluated were considered as experimental unit. Other complementary evaluations were plant height measurement (cm), from soil surface to rachis insertion point and number of days from planting to rachis emission, with the latter being evaluated during inflorescence emission.

To carry out the analysis of variance, two decompositions of the variation sources were used. For the data submitted to periodical collection, the split plot in time was applied with a factorial scheme being used to install the experiment in the plots. For the remaining variables, variation decomposition was usually applied for factorial scheme (Steel et al. 1997). When necessary, regression models were applied to study the variables. The following criteria were taken into consideration for selecting the models: lower AIC (*Akaike information criterion*) value, model significance and greater determination coefficient (R<sup>2</sup>).

To describe plant growth along the months, the



## DEVELOPMENT OF THE BANANA PLANTS

was used where:

- $Y_i$  = represents the growth observed in time  $t_i$ ;
- $t_i$  = the time spent from transplanting until determination of pseudostem height, in months;
- $a$  = asymptotic height;
- $b$  = scale parameter;
- $k$  = mean growth rate.

All analyses were carried out with the help of the statistical software R (R Core Team 2006).

## RESULTS AND DISCUSSION


No significant effect was observed for the “dose  $\times$  cultivar” interaction for any of the characteristics evaluated (Table I). The cultivar ‘Prata Anã’ presented higher leaf number, leaf emission rate and sucker number, shorter pseudostem height, foliar area of the leaves emitted after PBZ application and lower sucker fresh mass than the cultivar FHIA-01. Paclobutrazol (PBZ) did not influence pseudostem circumference and number of days from planting to flowering, in agreement with the results obtained by El Otmani et al. (1992), who evaluated the effect of PBZ on the banana plants ‘Grande Naine’ cultivated in protected environment in Morocco. On the other hand, in cloves (*Dianthus caryophyllus* L.) increase in the number of days from planting to flowering was verified only in the plants that received doses above 0.9 mg of a.i. of PBZ plant<sup>-1</sup> (Bañón et al. 2002).

Pseudostem growth reduction could be observed 17 days after application of the product, with a mean reduction of 14% (Fig. 1a). Increased PBZ doses led to shorter pseudostem height at the end of the assessment period (Fig. 1b). The parameters estimated confirm that increased PBZ doses provided a reduction in pseudostem height ( $a$ ) and growth mean rate ( $k$ ), Table II. Percent difference in pseudostem height at the end of the assessment period, between the control dose and the other doses varied between 18 and 30%, with an approximate mean of 26%. Doses between 1.5 and 2.0 g of a.i. of PBZ plant<sup>-1</sup> provided similar effects on the growth mean rate and the final pseudostem height. PBZ was

responsible for pseudostem elongation (Fernández 1995, Winkler and Helentjaris 1995).

Since a relation was visually observed between pseudostem height at the moment of application and the PBZ effect on plant growth, this situation was carefully evaluated. Figure 1d presents mean pseudostem height of the plants in the blocs before application and at the end of the assessment period (after product application). Block 1 plants were found to present shorter pseudostem height before application and at the end of the assessment period while block 8 plants showed, on average, the highest height. A regression analysis of pseudostem heights at the moment of PBZ application in function of added pseudostem height after product application presented a positive and significant regression coefficient ( $p = 0.0011$ ). This result indicates that the PBZ inhibiting effect was lower when the plants had a higher height at the moment of application, what has been explained by an effect of dilution in the total mass of the plant or by some problem associated with the sites of action of the product (less activity or less sensitivity). The results found agree with those obtained by (Bandara et al. 1998), who, evaluating the effect of time of PBZ application on vase-cultivated banana plants, observed that the effect of PBZ applied at initial stages was higher than when applied at advanced culture stages.

No differences were observed in total foliar area of the PBZ-treated plants (Table I). These results agree with those found by El Otmani et al. (1992) who observed a reduction in the total foliar area of banana plants ‘Grande Naine’ after application of 1.5 g a.i. of PBZ plant<sup>-1</sup>. However, a reduction in the foliar area was verified in the leaves emitted after PBZ application, as measured by the foliar area of the expanded youngest leaf. This observation shows that additions in the PBZ doses reduced the foliar area of the youngest leaf (Fig. 1e). A likely explanation for this fact (reduction of the area of the youngest leaf without affecting the total foliar area) would be a tendency in increased leaf permanence on PBZ-treated plants (Table III), since the product did not cause significant differences in leaf emission rate (Table I). These results





DEVELOPMENT OF THE BANANA PLANTS

TABLE I

F test probability values of the analysis of variance (p), coefficient of variation (CV%) and mean per cultivar for the variables: pseudostem circumference (PSC), number of days from planting to flowering (NDF), pseudostem height (PSH), plant height (HEI), total foliar area (TFA), foliar area of the totally expanded youngest leaf (YLA), leaf number (LN), leaf emission rate (LER), sucker number (SN), sucker fresh mass (SFM).

Variables	p			CV (%)	Means <sup>(1)</sup>	
	Cultivars	Dose	Dose × cultivar interaction		Prata Anã	FHIA 01
PSC (cm)	0.409	0.475	0.728	7.66	64.18 a	63.32 a
NDF	0.939	0.090	0.452	9.58	354.48 a	342.44 a
PSH (cm)	0.006	< 0.001	0.300	13.27	74.39 b	79.80 a
HEI (cm)	0.014	< 0.001	0.915	12.97	157.67 b	169.91 a
TFA (cm <sup>2</sup> )	0.920	0.180	0.980	21.70	39792.00 a	39918.00 a
YLA (cm <sup>2</sup> )	0.001	< 0.001	0.380	18.13	5119.00 b	5969.00 a
LN	< 0.001	0.002	0.877	11.65	12.23 a	10.58 b
LER (month leaves <sup>-1</sup> )	< 0.001	0.265	0.558	15.19	1.80 a	1.54 b
SN	0.013	0.016	0.935	20.96	6.38 a	5.65 b
SFM (kg)	0.032	0.205	0.602	41.70	10.35 b	12.70 a

(1) Mean pairs followed by same letter, in the lines, do not differ by the F test at 5% de probability.

TABLE II

Parameters estimated for the growth curves,  $Y_i = a/(1 + e^{b-k \times t_i})$ , and coefficient of determination ( $R^2$ ) to describe the growth of banana plants ‘Prata Anã’ and ‘FHIA 01’ under five doses of paclobutrazol (PBZ, g of a.i. plant<sup>-1</sup>) applied on soil.


PBZ doses	Growth curve parameters			$R^2$
	$a$	$b$	$k$	
0.0	145.53	3.14	1.08	0.996
0.5	118.79	2.96	1.19	0.993
1.0	106.51	2.88	1.24	0.989
1.5	102.61	2.91	1.29	0.988
2.0	99.96	2.98	1.33	0.984

TABLE III

Means of the number of leaves of banana plants ‘Prata Anã’ and ‘FHIA 01’ in function of paclobutrazol doses (PBZ), along the experiment evaluation.

Months	PBZ doses (g of a.i. plant <sup>-1</sup> )					Ĉ
	0.0	0.5	1.0	1.5	2.0	
Number of leaves						
March	14.44	14.69	14.25	14.38	13.89	-0.56 <sup>ns</sup>
June	9.37	10.19	11.94	11.25	12.50	8.37*
September	10.31	10.25	12.31	12.06	12.07	5.43 <sup>+</sup>

$\hat{C}$ : contrast among doses 0.0 g of a.i. of PBZ (control) and the other levels



after PBZ application, but without any alterations in the total foliar area of the plant.

PBZ did not influence sucker fresh mass, indicating that the plants, regardless of the paclobutrazol dose applied, had a similar sucker production and that the occurrences of sucker number variations were due to the appearance of a larger number of lateral buds (Table I). Since it was not possible to adjust a model that could explain the result found for all the doses evaluated for sucker number, we opted for adjusting a simple linear regression model up to the dose of 1.5 g of a.i. plant<sup>-1</sup>, considering that at the dose of 2.0 of a.i. plant<sup>-1</sup> there was an expressive sucker emission reduction (Fig. 1f). (Hamid and Williams 1997) studying the behavior of *Swainsona formosa* under the effect of different growth regulators, verified that PBZ application was followed by an increase in the number of lateral shoots, that would be reduced at higher PBZ levels.

#### RESUMO

Em algumas condições de cultivo o crescimento excessivo do pseudocaule de bananeiras pode ser considerado um fator limitante, assim a utilização de reguladores de crescimento pode constituir uma alternativa. O presente trabalho teve por objetivo avaliar a ação do paclobutrazol sobre o crescimento do pseudocaule de bananeiras e outras características do seu desenvolvimento. Para isso foi instalado um experimento com cinco doses do paclobutrazol (0; 0,5; 1,0; 1,5; e 2,0 g de i.a. planta<sup>-1</sup>) combinadas com dois cultivares de bananeira ('Prata Anã' e 'FHIA-01'). O paclobutrazol não afetou o número de dias do plantio ao florescimento, a área foliar total e a taxa de emissão de folhas. Contudo, aumentou a permanência das folhas nas plantas e o número de perfilhos, reduziu a área foliar das folhas emitidas após a sua aplicação e proporcionou uma redução média de 26% na altura do pseudocaule.

**Palavras-chave:** área foliar, análise de crescimento, reguladores de crescimento, *Musa* spp., perfilhos.

#### REFERENCES

- ALVES EJ. 1999. A cultura da banana: aspectos técnicos, socioeconômicos e agroindustriais. 2. Ed. Brasília: Embrapa.
- BAÑÓN S, GONZÁLEZ A, CANO EA, FRANCO JA AND FERNÁNDEZ JA. 2002. Growth, development and colour response of potted *Dianthus caryophyllus* cv. Mondriaan to paclobutrazol treatment. *Sci Hortic* 94: 371–377.
- CARVALHO JABS, PEIXOTO CP, SILVA SO, LEDO CAS, PEIXOTO MFSP AND ALVES JS. 2005. Uso da gibberelina GA<sub>3</sub> na seleção do porte de bananeira das cultivares Prata e Prata-Anã. *Rev Bras Frutic* 27: 449–453.
- EL OTMANI M, JABRI K AND SEDKI M. 1992. Paclobutrazol effect on development of greenhouse-growth banana: 2-year assessments. *Acta Hort* 296: 89–96.
- FERNÁNDEZ JAS, DOUMAS P, TEISSON C AND CÔTE F. 1995. Identificación y cuantificación de giberelinas en plantas variantes somaclonales y normales de *Musa* (cv. 'Grande Naine' AAA) mediante HPLC y espectrometria de masa. Proceeding 11<sup>th</sup> Reunión de la Asociación para la Cooperación en Investigación de Banano en El Caribe y en América Tropical, 13-18 February, San José, Costa Rica. Available at < [http://musalit.inibap.org/byindex\\_result.php?lang=en&index=nomed&idselect=152&page=12](http://musalit.inibap.org/byindex_result.php?lang=en&index=nomed&idselect=152&page=12)>.
- GUBBUK H, PEKMEZC IM AND ERKAN M. 2004. Production potential of Cavendish cultivars (*Musa* spp. AAA) under greenhouse and field conditions in subtropical areas of Turkey. *Acta Agric Scand* 54: 249–253.
- HAMID MM AND WILLIAMS RR. 1997. Effect of different types and concentrations of plant growth retardants on Sturt's desert pea (*Swainsona formosa*). *Sci Hortic* 71: 79–85.
- JEYAKUMAR P, KUMAR N AND KAVINO M. 2003. Physiological response of banana cv. 'Robusta' (AAA) to foliar applied plant growth regulators on productivity. *Madras Agric J* 90: 702–706.
- KHURSHID T, MCNEIL DL, TROUGHT MCT AND HILL GD. 1997. The response of young 'Braeburn' and 'Oregon spur delicious' apple trees growing under an ultrahigh density planting system to soil-applied paclobutrazol: I. Effect on reproductive and vegetative growth. *Sci Hortic* 71: 11–24.
- LEONEL S, GOMES EM AND PEDROSO CJ. 2004. Desempenho agrônômico de bananeiras micropropagadas em Botucatu. *SP Rev Bras Frutic* 26: 245–248.



#### DEVELOPMENT OF THE BANANA PLANTS

- R DEVELOPMENT CORE TEAM. 2006. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria.
- RADEMACHER W. 2000. Growth retardants: effects on gibberellin biosynthesis and other metabolic pathways. *Annu Rev Plant Physiol Plant Mol Biol* 51: 01–531.
- SILVA SO, FLORES JCO AND LIMA NETO FP. 2002. Avaliação de cultivares e híbridos de bananeira em quatro ciclos de produção. *Pesq Agropec Bras* 37: 1567–1574.
- STEEL GD, TORRIE JH AND DICKEY DA. 1997. Principles and procedures of statistics: a biometrical approach. 3<sup>rd</sup> ed., New York: McGraw-Hill, 666 p.
- TURNER DW. 2003. An integral method for estimating leaf area in bananas. *Infomusa* 12: 15–17.
- WINKLER RG AND HELENTJARIS T. 1995. The *Dwarf3* gene encodes a cytochrome P<sub>450</sub>-mediated step in gibberellin biosynthesis. *Plant Cell* 7: 1301–1311.
- YIM KO, KWON YW AND BAYER DE. 1997. Growth responses and allocation of assimilates of rice seedlings to roots after paclobutrazol and gibberellin treatment. *J Plant Physiol* 16: 35–41.