



Anais da Academia Brasileira de Ciências

ISSN: 0001-3765

aabc@abc.org.br

Academia Brasileira de Ciências

Brasil

Marzinek, Juliana; Oliveira, Denise M.T.

Structure and ontogeny of the pericarp of six Eupatorieae (Asteraceae) with ecological and taxonomic considerations

Anais da Academia Brasileira de Ciências, vol. 82, núm. 2, junio, 2010, pp. 279-291

Academia Brasileira de Ciências

Rio de Janeiro, Brasil

Available in: <http://www.redalyc.org/articulo.oa?id=32713482004>


- How to cite
- Complete issue
- More information about this article
- Journal's homepage in redalyc.org

redalyc.org

Scientific Information System

Network of Scientific Journals from Latin America, the Caribbean, Spain and Portugal

Non-profit academic project, developed under the open access initiative



Anais da Academia Brasileira de Ciências (2010) 82(2): 279-291
(Annals of the Brazilian Academy of Sciences)
ISSN 0001-3765
www.scielo.br/aabc

Structure and ontogeny of the pericarp of six Eupatorieae (Asteraceae) with ecological and taxonomic considerations

JULIANA MARZINEK¹ and DENISE M.T. OLIVEIRA²

¹Instituto de Biologia, Universidade Federal de Uberlândia

Rua Ceará, s/n, Bloco 2D, sala 28, Umuarama, 38405-315, Uberlândia, MG, Brasil

²Departamento de Botânica, Instituto de Ciências Biológicas, Universidade Federal de Minas Gerais

Avenida Antonio Carlos, 6627, Pampulha, 31270-901 Belo Horizonte, MG, Brasil

Manuscript received on October 20, 2008; accepted for publication on June 4, 2009

ABSTRACT

The ontogeny of cypselae and their accessory parts were examined using light and scanning electron microscopy for the species *Campuloclinium macrocephalum*, *Chromolaena stachyophylla*, *Mikania micrantha*, *Praxelis pauciflora*, *Symphypappus reticulatus*, and *Vittetia orbiculata*, some of these being segregated from the genus *Eupatorium*. A layer of phytomelanin observed in the fruit appears to be secreted by the outer mesocarp into the schizogenous space between the outer and inner mesocarp; its thickness was observed to vary among the different species examined. The bristles of the pappus are vascularized, except in *M. micrantha*, and have cells that are superficially projected and arranged acropetally; in *S. reticulatus* some of the projections are retrorse and a fracture line on the floral disk that is only seen in this species may indicate a double dispersal process. Numerous differences observed among the cypselae examined here reinforce earlier segregations of the genus *Eupatorium sensu lato*.

Key words: : anatomy, Asteraceae, carpopodium, fruit, pappus, phytomelanin.

INTRODUCTION


The tribe Eupatorieae (Asteraceae) comprises 190 genera and 2,000 species (Anderberg et al. 2007) that are encountered primarily in Mexico and Central and South America, with some representatives from North America, but with few species in the Old World (King and Robinson 1987). *Eupatorium* is a very complex genus and has experienced numerous segregations, as compiled by King and Robinson (1987).

Wagenitz (1976) suggested that anatomical studies of the fruits of the Asteraceae might aid in elucidating its systematics. Characteristics of the indumentum (Ritter and Miotto 2006), pappus (Bean 2001), carpopodium (Haque and Godward 1984), and the anatomy

combination of all of the characters cited above (Leal et al. 1997), have been used to delimit tribes, genera, and even species of this family.

In spite of the global occurrence of the family, the fruits of the Asteraceae have not been intensively studied, as can be confirmed by the discordance seen in the names attributed to its fruits. Marzinek et al. (2009) took into consideration many anatomical and ecological aspects of Asteraceae fruits, especially their complex origin, and reaffirmed their true nature as cypselae.

One character that stands out in descriptions of the cypselae of the Heliantheae *s.l.* (which includes Eupatorieae) is their dark appearance (Anderberg et al. 1997). This aspect can be attributed to the deposition of a layer of phytomelanin, an organic material that



Dhakal 2001). Sometimes this phytomelanin layer is referred to as a “carbon layer” or black pigment layer (see Roth 1977). Panero (2007) indicates the occurrence of phytomelanin as a synapomorphy of the Asteraceae. The Phytomelanin Cypsela Clade (PCC) is composed of more than 5,000 species that produce this substance.

Ontogenetic studies of the reproductive organs of Eupatorieae are generally scarce, and are almost exclusively limited to the publications of Pandey and Singh (1983, 1994). In spite of the importance of the Eupatorieae within the Asteraceae, no single study has focused simultaneously on the ontogeny, indumentum, pappus, and carpodium of the species within this taxon.

The present work analyzed the development of the pericarp and the accessory parts of the cypselae of six species of Eupatorieae from Brazil (some of them segregated from *Eupatorium s.l.*), emphasizing the most relevant structures to taxonomic and ecological questions related to this tribe.

MATERIALS AND METHODS

Specimens of *Campuloclinium macrocephalum* (Less.) DC., *Chromolaena stachyophylla* (Spreng.) R. King and H. Robinson, *Mikania micrantha* H.B.K., *Praxelis pasciflora* (H.B.K.) R. King and H. Robinson, *Symphycarpus reticulatus* Baker, and *Vittetia orbiculata* (DC.) R. King and H. Robinson were collected in *cerrado* vegetation fragments in the municipality of Botucatu, São Paulo State, Brazil. Reference plant material was prepared and deposited in the BOTU Herbarium (Holmgren et al. 1990) as collections 25,552 to 25,557.

Micromorphological analyses of the surfaces of the cypselae were performed with material fixed in glutaraldehyde (2.5% in 0.1M phosphate buffer, pH 7.3, maintained at 4°C). Samples were processed and, then, mounted on aluminum stubs, gold coated, and subsequently examined using a scanning electron microscopy (Quanta 200, FEI Company) to generate digital images.

Anatomical studies were performed on ovaries and cypselae in different phases of development that had been fixed in FAA 50 for 48 hours (Johansen 1940) and, then, conserved in 70% ethanol (Jensen 1962). The fixed

gitudinal sections 6 to 10 μ m thick were prepared using a rotary microtome, stained with 0.05% toluidine blue at pH 4.7 (O'Brien et al. 1964), and subsequently mounted in synthetic resin. The permanent slides were observed under an Olympus BX41 optical microscope and the images captured digitally.

The following histochemical tests were performed: ruthenium red to indicate the presence of polysaccharides and pectins (Jensen 1962); phloroglucinol with hydrochloric acid to stain lignified cell walls (Sass 1951); Sudan IV to stain lipidic substances; Lugol solution for starch; ferric chloride with sodium carbonate to indicate the presence of phenolic compounds (Johansen 1940); Dragendorff reagent to detect alkaloids (Yoder and Mahlberg 1976); and bromophenol blue for proteins (Mazia et al. 1953).


The results are described in ontogenetic terms. Considering the inferior origin of the ovary, the *lato sensu* definition of the pericarp was adopted, in which the exocarp is produced by the outer epidermis of the inferior ovary, the endocarp by the inner epidermis, and the mesocarp by the ground region where the vascular tissues are inserted.

Trichomes distribution was analyzed using both scanning electron and optical microscopy. Ten cypselae ($n=10$) of each species mounted in glycerin were also observed. The terms scarce (when trichomes were present on up to 10% of the cypselae surface), frequent (between 11% and 50%), and abundant ($>50\%$) were used.

The terminology used to designate the outward projected portions of the pappus fibers was adapted from Hickey (1979), considering the angle of divergence between the distal portion of the projected cells and the apex of the bristles. These angles were classified as narrow ($>45^\circ$) or moderate (between 45° and 65°) acute. The nomenclature proposed by Barthlott et al. (1998) was used to describe the bristle surfaces.

RESULTS

The ovaries of the floral buds of the species examined were all inferior (Fig. 1), bicarpelar, and unilocular



PERICARP OF EUPATORIEAE (ASTERACEAE)




Fig. 1 – Ovary of the floral bud. Figs. 1, 5, 7, 9 – Longitudinal sections. Figs. 2-4, 6, 8, 10 – Transversal sections. Fig. 1 – *Campuloclinium macrocephalum*; note lignified basal epidermal cells (arrow). Fig. 2 – *Mikania micrantha*. Figs. 3-4 – Detail of the ovarian walls of *Campuloclinium macrocephalum* and *Symphyopappus reticulatus*, respectively. Fig. 5 – *Campuloclinium macrocephalum*; note fusiform nuclei of mesophyll cells (arrowhead). Fig. 6 – Detail of the ovarian wall of *Mikania micrantha* showing the transmission tissue (*). Fig. 7 – Detail of the floral disk of *Praxelis pauciflora*; note salient external cells of the pappus with evident nuclei (arrow). Fig. 8 – Detail of part of the floral disk of *Mikania micrantha* demonstrating secretory cavities near the vascular bundles. Figs. 9-10 – Carpopodium of *Symphyopappus reticulatus*. Abbreviations: ca – carpel; ie – inner epidermis; im – inner mesophyll; lo – locule; oe – outer epidermis; om – outer mesophyll; ov – ovule; ow – ovarian wall; pa – pappus; pe – petal; vb – vascular bundle. Bars – 250µm (1); 150µm (2, 10); 20µm (3, 5); 50µm (4, 6-7, 9); 100µm (8).

The outer ovarian epidermis is uniseriate (Fig. 3) with cubical cells, slightly longitudinally elongated (Fig. 5) and vacuolated, with visible nuclei; biseriate tectorial cells (Fig. 4), as well as uni- and biseriate glandular trichomes.

The ovarian mesophyll is composed of two regions. The outer mesophyll region demonstrates elongated cells composed of two layers of cells and longitudinally elongated cells; in transversal section, the subepidermal cells appear isodiametric.



Campuloclinum macrocephalum (Fig. 3), *Ch. stachyophylla*, *P. pauciflora*, and *V. orbiculata* have two layers in the outer mesophyll, while five to six cell layers are observed in *M. micrantha* (Fig. 6) and *S. reticulatus* (Fig. 4). Procambial strands vascularize the ovary in the outer mesophyll, but differentiated vascular elements are only rarely observed (Fig. 6).

The inner mesophyll demonstrates from four to five cylindrical layers of small-diameter cells arranged longitudinally, with evident fusiform nuclei (Fig. 5). The more internal layers have large intercellular spaces and irregular shapes in comparison with the external layers (Figs. 3-4, 6).

The inner epidermis in *Ch. stachyophylla*, *P. pauciflora*, *S. reticulatus* (Fig. 4), and *V. orbiculata* is uniseriate and little evident, while in *Ca. macrocephalum* (Fig. 3) and *M. micrantha* these cells are elongated longitudinally, juxtaposed, with evident nuclei, and of large diameters in comparison with the cells of the inner mesophyll. The transmission tissue in all of the species studied can be seen in the inner epidermis where the carpels are joined, and it is composed of a multiseriate region of small diameter cells with pectin-containing cell walls and evident nuclei (Fig. 6).

The floral disk has an uniseriate epidermis that covers varying layers of elongated parenchyma cells disposed obliquely along the periphery; these cells form a protuberance where the pappus is inserted (Fig. 7); there are vascular traces of other floral parts immersed in this same region. In *M. micrantha*, two lateral secretory cavities were observed near the bundles that irrigate the corolla (Fig. 8).

A pappus is present in all of the species studied and is connate at its base. Each bristle is composed of a multiseriate group of cells that are rounded in transversal section and longitudinally elongated. The distal extremities of the external cells are projected towards the exterior, and evident nuclei are seen in the cells in this region (Fig. 7).

A protuberance that constitutes the carpopodium and is composed of cubic parenchyma cells with thin walls can be observed at the basal region of the ovary.


metrical carpopodium (Fig. 9), with epidermal cells of different sizes that are larger than the internal cells and arranged in a single layer (Fig. 10). The other species have indistinct carpopodium at the median section of the ovary. The ovule trace is single and robust, reaching to the base of the locule (Fig. 1).

The exocarp remains unaltered during development (Figs. 11, 13, 17, 19, 21), except in *M. micrantha*, where it develops an ornamented cuticle (Fig. 15). The cellular density of the outer mesocarp distinctly increases in all of the species, and these cells have large nuclei (Figs. 13, 17).

The most significant events that occur in the young pericarp take place in the inner mesocarp, with the most external cells forming numerous anticlinal projections in the direction of the outer mesocarp, and producing schizogenous spaces that are evident between the two mesocarpic regions (Figs. 11-22). These spaces are most ample in *S. reticulatus* (compare Figs. 19 and 20 with the other species). The lobed contour of the internal periclinal wall of the outer mesocarp (which is in contact with the numerous anticlinal projections referred to above) produces large intercellular spaces in *M. micrantha* and *S. reticulatus*, (Figs. 16, 20). The external peripheral layer of the inner mesocarp becomes lignified during the later stages of development in all of the species examined, forming sclereids.

The deposition of phytomelanin is initiated in the intercellular spaces described above, forming a continuous, thick, dark layer that is evident in all of the species, but thickest in *M. micrantha* and *S. reticulatus* (Figs. 23-34). Phytomelanin occupies the whole space between the outer and inner mesocarp, but did not give a positive reaction in any of the histochemical tests applied.

Lignification of the walls of the external cells of the inner mesocarp occurs rather late in the development of *M. micrantha*. The schizogenous spaces between the two mesocarpic regions in this species are contiguous with the intercellular spaces of the outer mesocarp, resulting in an irregular deposition of phytomelanin so that it occupies only part of that space (Fig. 28). In *Ca. macrocephalum* and *S. reticulatus* (Fig. 35), as well as



PERICARP OF EUPATORIEAE (ASTERACEAE)




Fig. 2 – Pericarp during development. From left to right, first and third columns – Transversal sections. Second and fourth columns – Longitudinal sections. Figs. 11-22 – Immature pericarp initiating the formation of schizogenous spaces (arrowhead). Figs. 23-34 – Pericarp during maturation with phytomelanin deposits. Figs. 11-12, 23-24 – *Campuloclinium macrocephalum*. Figs. 13-14, 25-26 – *Chromolaena stachyophylla*. Figs. 15-16, 27-28 – *Mikania micrantha*; note phytomelanin deposits on the outer mesocarp (arrow). Figs. 17-18, 29-30 – *Praxelis pauciflora*. Figs. 19-20, 31-32 – *Symphyopappus reticulatus*. Figs. 21-22, 33-34 – *Vittetia orbiculata*. (fl – fibrous layer; im – inner mesocarp; om – outer mesocarp). Bars – 20 μ m (11-14, 17-18, 21-26, 29-30, 33-34); 50 μ m (15-16, 19-20, 27-28, 31-32).

stachyophylla (Fig. 36), *M. micrantha*, and *P. pauciflora*. The inner mesocarp collapses at this time and is partially reabsorbed (Fig. 24).

towards the apex. All the other species examined have small cypselae, making this analysis much more difficult.




Fig. 3 – Cypselae during maturation. Figs. 35-36, 42-43 – Transversal sections. Figs. 39, 41, 44 – Longitudinal sections. Fig. 35 – Phytomelanin deposit in the external intercellular spaces of the vascular bundle of *Symphyopappus reticulatus*. Fig. 36 – Interruption of phytomelanin layer in the proximity of a vascular bundle in *Chromolaena stachyophylla*. Fig. 37 – Mature cypsel of *Praxelis pauciflora*. Figs. 38-39 – Detail of the micromorphology and anatomy of the floral disk region of *Chromolaena stachyophylla* respectively. Figs. 40-41 – Detail of the micromorphology (arrowhead – start of the fracture line) and anatomy of the pappus crown of *Symphyopappus reticulatus*. Figs. 42-43 – Mature pericarp of *Campuloclinium macrocephalum*. Fig. 44 – Carpodium of *Vittetia orbiculata* showing lignified surface. (ca – carpodium; fd – floral disk; pa – pappus; pe – pericarp; se – seed). Bars – 75 μ m (35); 20 μ m (36); 50 μ m (37); 100 μ m (38-39, 44); 30 μ m (40); 150 μ m (41-43).

upper region of the fruit: after lignification of the most internal cell walls of the floral disk (Fig. 39), the peripheral parenchyma cells dehydrate and collapse, thereby reducing their volume in most species (Fig. 38) and increasing the distance among the pappus bristles. Only *S. reticulatus* demonstrated lignification of the cell walls

The bristles of the pappus also become lignified during fruit development, but remain alive throughout this phase and show conspicuous nuclei (Fig. 39).

Mature pericarps undergo intense dehydration mainly in their furrows, and most of the cell layers collapse so that only some exocarp cells, fibers, vascular



PERICARP OF EUPATORIEAE (ASTERACEAE)

The pappus is trichomic at maturity, with a persistent, bristly, and isomorphous aspect; the bristles are ordered into a whorl. The pappus has from 14 to 43 bristles, depending on the species: 14-16 in *M. micrantha*; 16-19 in *P. pauciflora*; 17-20 in *Ch. stachyophylla*; 22-25 in *V. orbiculata*; 25-27 in *S. reticulatus*; and 40-43 in *Ca. macrocephalum*. The distal extremities of the cells projecting from the surface of the bristles have a consistently acute divergence angle that ranges from narrow in *Ca. macrocephalum* (Fig. 45), *Ch. stachyophylla* (Fig. 46) and *M. micrantha* (Fig. 47), to moderate in *P. pauciflora* (Fig. 48), *S. reticulatus* (Fig. 49) and *V. orbiculata* (Fig. 50). These extremities have sharp apices, except in *M. micrantha* (Fig. 47) where it is rounded; the cell extremities in *S. reticulatus* are also sometimes rounded.

Micromorphological examination of the bristle surfaces reveals variable patterns: reticulated in *Ca. macrocephalum* (Fig. 51); short longitudinal threads in *Ch. stachyophylla* (Fig. 52); finely striated and with projections demonstrating slight apical compression in *M. micrantha* (Fig. 53); and more finely striated on the surface of *P. pauciflora* (Fig. 54), while smooth in *S. reticulatus* (Fig. 55) and *V. orbiculata* (Fig. 56). Some of the bristle projections on the pappus of *S. reticulatus* are retrorsely flexed (see detail in Fig. 49).

The pappus bristles of the species examined here demonstrate completely lignified cell walls (Figs. 57-62). Most bristles are vascularized, with the exception of those of *M. micrantha*, but the extension of the vascular bundles varied among all the species. Only 1.7% of the total length of the bristles are vascularized in *Ch. stachyophylla*; only 2% in *Ca. macrocephalum*; 4% in *P. pauciflora*; 10% in *S. reticulatus*; and in *V. orbiculata* 37% of the length of each bristle has a small vascular bundle.

The carpopodium is lignified and symmetrical at maturity in most of the species examined (Figs. 63-65, 69), being asymmetrical only in *P. pauciflora* (Fig. 66) and *S. reticulatus* (Figs. 67-68). The carpopodium surface ranged from: undifferentiated in *M. micrantha*, having a thickness of approximately two cell layers in *Ch. stachyophylla*; having from one to three differentiated

The indumentum was observed to be v among the cypselsae evaluated, as can be seen b
paring Figure 37 (*P. pauciflora*), 38 (*Ch. stachyop*
and 40 (*S. reticulatus*). Five types of trichome
observed: two biseriate tectors and three glandu
latter ranging from uni- to biseriate. Table I l
five trichome types observed among the species
ined, specifying the variations in their horizontal
ribs and furrows) and vertical (considering the
mal third, median, and distal regions) distributi
the cypsela.

DISCUSSION

The species examined here demonstrated ovarian
tures typical of the Eupatorieae, including an un
epidermis with trichomes, outer mesophyll with
juxtaposed cells, inner mesophyll composed of
chyma cells with ample intercellular spaces, and
ner epidermis with differentiated cells in two
that form the transmission tissue. This data corro
the only previously published work concerning th
(Pandey and Singh 1983, 1994).

The dark color of the cypselsae observed in
groups of Asteraceae is due to the presence of ph
lanin, and this substance is deposited in the scl
nous spaces between the outer and inner mesocar
species studied. According to Hanausek (1910)
Roth (1977), there are three possible origins of th
bon or phytomelan layer”: “intracellular secreti
the accumulation of a black pigment within the
especially the epidermis cells; by an “intercellula
tion” that is formed in lysigenous intercellular
or the disintegration of entire cells or cell layer
subsequent carbonization of their walls. The spec
amined in the present study did not demonstrate
lar disintegration, while the outer mesocarp cel
secretory characteristics, such as dense cytoplasm
evident nuclei. This layer apparently synthesizes
polymerizes phytomelanin, which is then depos
the space created by the cellular projections betw
inner and outer mesocarp. According to Pande
(1989), phytomelanin precursors are synthesized





Fig. 4 – Micromorphology (first and second rows) and transversal sections of the pappus (third row), and micromorphology of the carpodium (fourth row). Figs. 45, 51, 57, 63 – *Campuloclinium macrocephalum*. Figs. 46, 52, 58, 64 – *Chromolaena stachyophylla*. Figs. 47, 53, 59, 65 – *Mikania micrantha*. Figs. 48, 54, 60, 66 – *Praxelis pauciflora*. Figs. 49, 55, 61, 67, 68 – *Symphyopappus reticulatus*. Figs. 50, 56, 62, 69 – *Vittetia orbiculata*. Bars – 50 μ m (45, 47, 49); 20 μ m (46, 48, 50-62); 150 μ m (63-69).

There were distinct differences among the thicknesses of the phytomelanin layers among the various species studied. *Mikania micrantha* and *S. reticulatus* have thicker phytomelanin layers than other species have, which may be related to the greater numbers of secretory layers in the outer mesocarps of these species. According to Hanausek (1910, 1912) *apud* Roth (1977), phytomelanin may serve as a protective screen against excessive insolation or in mechanical protection of the pericarp. As the studied species occur in the *cerrado* biome, protection against excessive insolation could be very important. Pandey and Dhakal (2001) reviewed the available literature concerning phytomelanin






spp). However, J. Marzinek and D.M.T. Oliveira (unpublished data) analyzed the cypsels of *M. micrantha* and *S. reticulatus* and found insect larvae in 10% and 22% of the fruits, respectively. As these are the species with the thickest recorded phytomelanin layers, the deterrent effect of this substance cannot be confirmed, and the role of phytomelanin in the cypsels still needs to be elucidated.

Phytomelanin was observed here to be deposited in an acropetal direction in *Ca. macrocephalum*, contrary to the direction reported by Vries (1948) *apud* Hegnauer (1977) for *Tagetes patula*, a species of Helenieae (*sensu* Bremer 1994). No other published references were en-



PERICARP OF EUPATORIEAE (ASTERACEAE)

TABLE I
Types* and distribution of trichomes on the cypselae of the examined species.
(A: apex; M: median region; B: base; -: absent; +: rare; ++: frequent; +++: abundant).

Type of trichoma															
		Type I			Type II			Type III			Type IV			Type V	
Species	Distribution	A	M	B	A	M	B	A	M	B	A	M	B	A	M
<i>Campuloclinium macrocephalum</i>	ribs	+	+	+	-	-	-	-	-	-	-	-	-	-	-
	furrows	-	-	-	-	-	-	+	+	+	-	-	-	-	-
<i>Chromolaena stachyophylla</i>	ribs	++	+	+	-	-	-	++	-	-	-	-	-	-	-
	furrows	+	+	+	-	-	-	+	-	-	-	-	-	-	-
<i>Mikania micrantha</i>	ribs	-	-	-	-	-	-	-	-	-	-	-	-	-	-
	furrows	-	-	-	-	-	-	+	-	-	+	+	+	-	-
<i>Praxelis pauciflora</i>	ribs	+++	+++	++	-	-	-	-	-	-	-	-	-	-	-
	furrows	++	++	+	-	-	-	-	-	-	-	-	-	-	-
<i>Symphiopappus reticulatus</i>	ribs	-	-	-	-	-	+	-	-	-	-	-	-	-	-
	furrows	-	-	-	++	++	++	-	-	-	-	-	-	-	-
<i>Vittetia orbiculata</i>	ribs	++	++	++	-	-	-	-	-	-	-	-	-	-	-
	furrows	+	+	+	-	-	-	+++	++	+	-	-	-	++	-

* Type I: tector, biseriate, 4-cells; Type II: tector, biseriate, 4-6 cells of hyaline content; Type III: glandular, biseriate, peduncle with 4-6 hyaline cells, distal 2-4 cells globoid to spatulate with dense content; Type IV: glandular, uniseriate, peduncle with 6-7 cells, secretory portion unicellular; Type V: glandular, biseriate, peduncle with 2-4 hyaline cells, distal 8-10 cells with dense content.

in this study, and by the velocity with which deposition occurred.

The indumenta of the cypselae of the Asteraceae have been examined by numerous authors (e.g. Drury and Watson 1966, Wetter 1983, Blanca and Guardia 1997, Mukherjee and Sarkar 2001), and this character has a significant taxonomic value. Roth (1977) observed that biseriate trichomes are typically found on the pericarp in this family, and they apparently aid in fixing dispersed fruits to the soil. Types I and III trichomes referred to in the present study (biseriate non-glandular with four cells, and biseriate glandular with peduncle composed of from four to six hyaline cells, respectively) were common among the Eupatorieae studied. Among the Asteraceae of the *cerrado* vegetation studied by Castro et al. (1997), type IV trichomes (glandular and uniseriate with peduncle composed of from six to seven cells and unicellular secretory portion) occurred exclu-

that some simple, pluricellular and uniseriate trichomes as well as many glandular trichomes, were observed on the surface of the fruits of *M. micrantha*. However, type IV trichomes that are found in the present examination of this species were not reported by those authors.

An analysis of the occurrence of trichomes on the different species revealed some interesting results (see Table I): *P. pauciflora* demonstrated a relatively homogeneous distribution of those structures, but had no biseriate tector trichomes known as twin hairs, the characteristic type of trichome in the Asteraceae. *V. orbiculata* demonstrated the most heterogeneous distribution of trichomes (including the biseriate tector trichomes as well as two additional secretory types), all of them being relatively abundant in the furrows of the cypselae. Differences in trichomes between the ribs and furrows are clearly distinguishable in *Ca. macrocephalum*. Considerable differences in the distribution of trichomes among the apical, medi-



selsae of different Asteraceae were quite characteristic. Their internal structure reveals certain characteristics that are taxonomically useful, mainly the arrangement of the support tissues that are preserved even after dehydration of the cypselae. Studies of the ontogeny of dry pericarps are likewise very informative among the Asteraceae, for few defined layers persist at maturity. However, only phytomelanin and residues of cell walls and vascular bundles remain after drying in the Eupatorieae, making comparisons among the different *taxa* very difficult. Pandey and Singh (1983, 1994) studied ontogenetic aspects of Eupatorieae fruits, but these authors cited only a few characteristics related to development.


Haque and Godward (1984) observed that the carpopodium (the abscission zone of the cypselae) sometimes demonstrated structures that facilitated separation from the inflorescence axis and the subsequent dispersal of the fruit. According to King and Robinson (1987), carpopodium anatomy is very variable in Eupatorieae. Among the species studied, *M. micrantha* had an indistinct carpopodium, while this structure is clearly asymmetrical in both *P. pauciflora* and *S. reticulatus*. *Vittetia orbiculata* had the most distinct carpopodium of all the species studied, followed then by *Ca. macrocephalum*. Haque and Godward (1984) examined the carpopodium and its usefulness in the taxonomy of the Asteraceae. These authors reported that *Ca. macrocephalum* had the most differentiated carpopodium among the Asteroideae (*sensu* Bremer 1994) studied. This finding is not in conflict with the present work, however, because these authors did not examine any species of *Vittetia* that had a more conspicuous carpopodium than *Ca. macrocephalum*.

An analysis of the floral disk of *M. micrantha* demonstrated the existence of two cavities associated with each vascular bundle. Reports of internal secretory structures are rare for the cypselae of the Asteraceae, being more frequently observed in vegetative organs. Castro et al. (1997) reported the presence of ducts in the leaves of *M. officinalis*, and Oliveira (1972) observed secretory elements in the leaves, flowers, and fruits of *M. hirsutissima* var. *hirsutissima*. These authors did not specify, however, what type of secretions were produced, in much

of the most important characteristics of the capitulum. According to Mukherjee and Sarkar (2001), the pappi encountered in the cypselae of the Asteraceae represent a classic source of taxonomic information and, depending on the degree of specificity of these characteristics, these structures can be used to identify genera or even species.

It is important to note that the vascular system of the pappus may be differentiated or not. The vasculature of the pappus has been described for some representatives of *Chaenactidinae* belonging to the tribe Heliantheae (Robinson 1981), as well as for some species of *Helogyne* (Eupatorieae), where the vascular bundle may extend for up to half of the length of the bristle (King and Robinson 1987). The bristles examined in the present work were all vascularized, with the exception of *M. micrantha*. Variations in the lengths of the bristle vascular bundles clearly demonstrates structural reduction of the calyx among the Asteraceae, reinforcing the classical affirmation of Cronquist (1955, 1981) and Ramiah and Sayeeduddin (1958) that the pappus is in fact a modified calyx. King and Robinson (1987) affirmed that the pappus is formed by a series of capillary bristles in most species of Eupatorieae, a condition that is considered primitive for the tribe. The pappus may even be reduced to five to ten bristles within this group, as in *Hofmeisteria*, or to just a few bristles as in *Fleischmannia*, or they may be totally absent as in *Ageratum* and some other genera. The presence of reduced vascularized lengths in the bristles of Eupatorieae is an indication of the evolutionary tendency to reduce the number of bristles, even to the point of completely suppressing the pappus. It has been well established that the vascular system is very conservative in plants, and that its reduction commonly precedes a reduction of the organ being served (Eames and MacDaniels 1947). This phenomenon has been previously reported for some species of Eupatorieae.

The pappus has an important role in fruit dispersal, including the performance of hygroscopic movements that regulate the moment of dispersal (Pijl 1982). In an aerodynamic sense, the pappus increases the resistance between the air and the fruit, thus prolonging the time



PERICARP OF EUPATORIEAE (ASTERACEAE)

hydration of the fibers of the floral disk apparently results in a retraction of the more peripheral fibers and the opening of the pappus bristles, thus favoring anemochoric dispersal.

Stuessy and Garver (1996) suggested that the pappus has a double function over time: it first has a defensive role in the flower, mainly against insects, while it is essential for dispersal in the fruit. This double function of the pappus may be one of the factors responsible for the great evolutionary success of the Asteraceae. The suggestion that the pappus has a role in the defense of the flowers was not supported in the species examined in the present study, however, because field observations revealed the existence of capitula that had been completely destroyed by insect predation (J. Marzinek, unpublished data).

Although the Asteraceae typically have a persistent calyx, some cypselae have a rupture line near their apex that seems to direct all of the bristles of the pappus into a single crown-like structure. This structure was observed only in *S. reticulatus* in the present study, but had been reported for *Carduus arvensis* by Dandeno (1905), who noted that the rupture line facilitates germination because water absorption can occur through the apical region of the cypselae. Shmida (1985) reported that a non-deciduous pappus promotes the dispersal of cypselae far from the mother plant. Experiments are needed to more accurately evaluate water uptake by seeds enclosed in the cypselae, and will be important in defining the role of the rupture line in germination.

It is interesting that the pappus is associated with two very distinct dispersal modes in the Asteraceae: biotic dispersal by epizoochory (in which the pappus bristles take on the form of hooks or become reflexed and able to attach to passing animals); and abiotic dispersal by anemochory (in which the extremities of the bristles tend to be erect). Curiously, diaspores of *S. reticulatus* demonstrate both modes, as the cypselae first escapes from the inflorescence axis by anemochory and, then, after initial dispersal, the backward-reflexed bristle extremities favor biotic dispersal.

Capillary pappi composed of thin and flexible bris-

and D.M.T. Oliveira, unpublished data). In the *S. reticulatus*, if the thick and rigid bristles of the pappus did not separate from the cypselae during dispersal, contact with the substrate would be much more difficult. The separation of the pappus crown may also facilitate water uptake and germination of the seeds in some species, but this hypothesis still needs to be tested.

The taxonomic characteristics of the species of the Eupatorieae here examined (including the anatomy of the pappus, carpopodium, indumentum, and cypselae maturation) are all very variable and support the recognition of the genus *Eupatorium* as proposed by Robinson (1987).

ACKNOWLEDGMENTS

The authors thank Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq) for the research grants awarded to J. Marzinek and D.M.T. Oliveira; Fundação de Amparo à Pesquisa do Estado de São Paulo (FAPESP), Programa BIOTA (Proc. 00/12469-7); Fundação de Amparo à Pesquisa do Estado de Minas Gerais (FAPEMIG) for financial support; and Dr. Antônio Magalhães Teles for identifying the Eupatorieae species.

RESUMO

A ontogênese das cípselas e de suas partes acessórias e a dispersão de espécies de *Eupatorium* e *Carduus* foram estudadas em *Eupatorium puloclinium macrocephalum*, *Chromolaena stachyophylla*, *Carduus micrantha*, *Praxelis pauciflora*, *Symphphyopappus reticulatus* e *Vittetia orbiculata*, parte delas segregadas do gênero *Eupatorium*, foi estudada em microscopia de luz e eletrônica de varredura. A camada de fitomelanina presente no fruto recentemente é secretada pelo mesocarpo externo e possui espessura variável entre as espécies, depositando-se em espaços quízógeno entre o mesocarpo externo e interno. As cípselas são vascularizadas, exceto em *M. micrantha*, e as células projetadas superficialmente, dispostas acropetamente. Em *S. reticulatus*, algumas projeções são retrógradas e a presença de linha de fratura sob o disco floral, observada apenas nesta espécie, pode indicar processo duplo de dispersão. As numerosas diferenças registradas entre as cípselas estudadas refor-



REFERENCES

- ANDERBERG AA ET AL. 2007. Compositae. In: KADEREIT JW AND JEFFREY C (Eds), Families in genera of vascular plants. Vol. VIII. Flowering Plants, Eudicots, Asterales, Berlin, Springer-Verlag, p. 61–576.
- BARTHOLOTT W, NEINHUIS C, CUTLER D, DITSCH F, MEUSEL I, THEISEN I AND WILHELMI H. 1998. Classification and terminology of plant epicuticular waxes. *Bot J Linn Soc* 126: 237–260.
- BEAN AR. 2001. Pappus morphology and terminology in Australian and New Zealand thistles (Asteraceae, tribe Cardueae). *Austrobaileya* 6: 139–152.
- BLANCA G AND GUARDIA D. 1997. Fruit morphology in *Tragopogon* L. (Compositae: Lactuceae) from the Iberian Peninsula. *Bot J Linn Soc* 125: 319–329.
- BREMER K. 1994. Asteraceae: Cladistics and Classification. Portland, Timber Press, 752 p.
- BRUHL JJ AND QUINN CJ. 1990. Cypselae anatomy in the ‘Cotuleae’ (Asteraceae-Anthemideae). *Bot J Linn Soc* 102: 37–59.
- CASTRO MM, LEITÃO-FILHO HF AND MONTEIRO WR. 1997. Utilização de estruturas secretoras na identificação dos gêneros de Asteraceae de uma vegetação do cerrado. *Rev Bras Bot* 20: 163–174.
- CRONQUIST A. 1955. Phylogeny and taxonomy of the Compositae. *Am Midl Nat* 53: 478–511.
- CRONQUIST A. 1981. An integrated system of classification of flowering plants. New York: Columbia University Press, 1262 p.
- DANDENO JB. 1905. The parachute effect of thistle-down. *Science* 22: 568–572.
- DRURY DG AND WATSON L. 1966. Taxonomic implications of a comparative anatomical study of Inuloideae – Compositae. *Am J Bot* 53: 828–833.
- EAMES AJ AND MACDANIELS LH. 1947. An introduction to plant anatomy. 2nd ed., New York, McGraw-Hill, 427 p.
- HABERLANDT G. 1914. Physiological plant anatomy. London: Macmillan *apud* SHELDON JC AND BURROWS FM. 1973. The dispersal effectiveness of the achene-pappus units of selected Compositae in steady winds with convection. *New Phytol* 72: 665–675.
- HANAUSEK TF. 1910. Über die perikarphöker von *Dahlia pinnatifida* (W.) Desf. *Bot. Anz. Wien* 26: 25–27.
- HANAUSEK TF. 1912. Untersuchungen über die kohleähnliche Masse der Kompositen. *Denkschr d kais Akad Wiss Wien, Math-Nat Kl* 87: 93–142 *apud* ROTH I. 1977. Fruits of Angiosperms. In: LINSBAUER K (Ed), *Handbuch der Pflanzenanatomie* v. 10, Berlin, Gebrüder Borntraeger, 675 p.
- HAQUE MZ AND GODWARD MBE. 1984. New records of the carpel in Compositae and its taxonomic use. *Bot J Linn Soc* 89: 321–340.
- HICKEY LJ. 1979. A revised classification of the architecture of dicotyledonous leaves. In: METCALF CR AND CHALK L (Eds), *Anatomy of the dicotyledons*. Oxford: Clarendon Press, p. 25–39.
- HOLMGREN PK, HOLMGREN NH AND BARNETT LC. 1990. Index Herbariorum. Part I: The Herbaria of the world. New York, International Association for Plant Taxonomy, 693 p.
- JENSEN WA. 1962. Botanical histochemistry: principles and practice. San Francisco: W.H. Freeman, 408 p.
- JOHANSEN DA. 1940. Plant microtechnique. New York, McGraw-Hill, 523 p.
- KING RM AND ROBINSON H. 1987. The genera of the Eupatorieae (Asteraceae). Kansas, Missouri Bot Gard, 581 p.
- LESZEK P, VINCENT D AND WILSON SL. 1997. The systematic value of the surface micromorphology and anatomy of cypselae of some members of the Senecioideae, Liabeae and Vernoniaceae (Asteraceae). *S Afr J Bot* 63: 382–399.
- MARZINEK J, DE-PAULA OC AND OLIVEIRA DMT. 2008. Cypselae or achene? Refining terminology by considering anatomical and historical factors. *Rev Bras Bot* 31: 549–553.
- MAZIA JC, BREMER PA AND ALFERT M. 1953. The cytochemical staining and measurement of protein with mercuric bromophenol blue. *Biol Bull* 104: 57–67.
- MUKHERJEE SK AND SARKAR AK. 2001. Morphology and structure of cypselae in thirteen species of the tribe Asteraceae (Asteraceae). *Phytomorphology* 51: 17–26.
- O’BRIEN TP, FEDER N AND MCCULLY ME. 1964. Polychromatic staining of plant cell walls by toluidine blue O. *Protoplasma* 59: 368–373.
- OLIVEIRA F. 1972. Contribuição para o estudo botânico de *Mikania hirsutissima* D.C var. *hirsutissima*. II. Morfologia externa e anatomia da folha, flor, fruto e semente. *Rev Farm Bioquim Univ São Paulo* 10: 15–36.
- PANDEY AK AND DHAKAL MR. 2001. Phytomelanin in Compositae. *Curr Sci* 80: 933–940.



PERICARP OF EUPATORIEAE (ASTERACEAE)

- PANDEY AK AND SINGH RP. 1994. Development and structure of seed and fruit in Eupatorieae and Heliantheae (Compositae). *Proc Natl Acad Sci India* 64: 115–126.
- PANDEY AK, LEE WW, SACK FD AND STUESSY TF. 1989. Development of the phytomelanin layer in fruits of *Ageratum conyzoides* (Compositae). *Am J Bot* 75: 739–746.
- PANERO JL. 2007. Compositae: key to the tribes of the Heliantheae Alliance. In: KADEREIT JW AND JEFFREY C (Eds), *Families in genera of vascular plants*. Vol. VIII. Flowering Plants, Eudicots, Asterales, Berlin, Springer-Verlag, p. 391–395.
- PIJL L VAN DER. 1982. Principles of dispersal in higher plants. New York, Springer-Verlag, 214 p.
- RAMIAH N AND SAYEEDUDDIN M. 1958. Homology of the pappus in the light of trichome distribution. *Curr Sci* 10: 402–403.
- RITTER MR AND MIOTTO STS. 2006. Micromorfologia da superfície do fruto de espécies de *Mikania* Willd. (Asteraceae) ocorrentes no estado do Rio Grande do Sul, Brasil. *Acta Bot Bras* 20: 241–247.
- ROBINSON H. 1981. A revision of the tribal and subtribal limits of the Heliantheae (Asteraceae). *Smithson Contrib Bot* 51: 1–102.
- ROTH I. 1977. Fruits of Angiosperms. In: LINSBAUER K (Ed), *Handbuch der Pflanzenanatomie* v. 10, Berlin, Gebrüder Borntraeger, 675 p.
- SASS JE. 1951. Botanical microtechnique. 2nd ed. Iowa State University, 228 p.
- SHMIDA A. 1985. Why do some Compositae have a persistently deciduous pappus? *Ann Missouri Bot C* 184–186.
- STUESSY TF AND GARVER D. 1996. The defensive pappus in heads of Compositae. In: CALIGARI PL AND HIND DJN (Eds), *Compositae: biology and utilization*. Proceedings of the International Compositae Conference, Kew, R Bot Gard, p. 81–91.
- VRIES MA DE. 1948. Over de vorming van phytochloor bij *Tagetes patula* L. en enige andere Compositae. Thesis, University of Leiden *apud* HEGNAUER R. 1950. The chemistry of the Compositae. In: HEYWOOD VH AND HIND DJN (Eds), *The biology and chemistry of the Compositae*. London, Academic Press, p. 284–335.
- WAGENITZ G. 1976. Systematics and phylogeny of the Compositae (Asteraceae). *Plant Syst Evol* 125: 29–46.
- WETTER MA. 1983. Micromorphological characters for generic delimitation of some new world Senecioneae (Asteraceae). *Brittonia* 35: 1–22.
- YODER LR AND MAHLBERG PG. 1976. Reaction of histochemical and histochemical indicators in laticifers and specialized parenchyma cells of *Catharanthus roseus* (Apocynaceae). *Am J Bot* 63: 1167–1173.