



Anais da Academia Brasileira de Ciências

ISSN: 0001-3765

[aabc@abc.org.br](mailto:aabc@abc.org.br)

Academia Brasileira de Ciências

Brasil

Macluf, Cecilia; Meza Torres, Esteban I.; Solís, Stella M.  
Isoetes pedersenii, a new species from Southern South America  
Anais da Academia Brasileira de Ciências, vol. 82, núm. 2, junio, 2010, pp. 353-359  
Academia Brasileira de Ciências  
Rio de Janeiro, Brasil

Available in: <http://www.redalyc.org/articulo.oa?id=32713482011>


- How to cite
- Complete issue
- More information about this article
- Journal's homepage in [redalyc.org](http://redalyc.org)

[redalyc.org](http://redalyc.org)

Scientific Information System

Network of Scientific Journals from Latin America, the Caribbean, Spain and Portugal

Non-profit academic project, developed under the open access initiative



Anais da Academia Brasileira de Ciências (2010) 82(2): 353-359  
(Annals of the Brazilian Academy of Sciences)  
ISSN 0001-3765  
www.scielo.br/aabc

## *Isoetes pedersenii*, a new species from Southern South America

CECILIA MACLUF<sup>1</sup>, ESTEBAN I. MEZA TORRES<sup>2</sup> and STELLA M. SOLÍS<sup>2</sup>

<sup>1</sup>Cátedras de Morfología Vegetal y Palinología, Facultad de Ciencias Naturales y Museo  
UNLP, Paseo del Bosque s/n, 1900, La Plata, Argentina

<sup>2</sup>Cátedra de Morfología Vegetal, Facultad de Ciencias Exactas y Naturales y Agrimensura  
Instituto de Botánica del Nordeste (UNNE-CONICET), Av. Libertad 5470, Corrientes, Argentina

Manuscript received on November 26, 2008; accepted for publication on September 28, 2009

### ABSTRACT

The name *Isoetes pedersenii* H.P. Fuchs (Lycophyta), a species known only from the Mburucuyá National Park, Corrientes, Argentina, is validated. Observations were carried out on herbarium material with stereoscopic, light and scanning electron microscopes. The species is described and typified. A diagnosis and discussion about its distribution and its relationship with the morphology of other species of *Isoetes* are provided.

**Key words:** Argentina, Corrientes, *Isoetes*, Lycophyta, new species.

### INTRODUCTION

The Isoetaceae constitutes a family with a wide distribution. The representatives of this family live from subarctic to warm regions in all continents from sea level to approximately 4,500 high. *Isoetes* L. has about 350 species (Hickey et al. 2003), of which seven species were reported by Ponce (1996) as growing in Argentina. They occur in the provinces of Buenos Aires, Formosa, Córdoba, Tucumán, and in the Patagonian region. Later, MacLuf and Ponce (2008) updated the occurrence of *Isoetes* in Argentina.

The two farms that constitute the Mburucuyá National Park belonged to the botanist Dr. Honoris Causa Troels Myndel Pedersen, who donated these farms in order to create a national park. For 50 years Pedersen was an assiduous collector who formed a herbarium with approximately 30,000 specimens (Arbo 2004). The duplicates of his collections were distributed to the following herbaria: A, B, BR, C, CTES, K, L, LP, MBM, MO, NY, P, S, SI, U, US, and his collections are cited like this in

collections gave to this park an added botanical value since many of Pedersen's specimens were recognized as types for many new species (Arbo 2004).

During the development of the Project “Flora del Parque Nacional Mburucuyá”, subproject “Flora de Lycophyta”, several nomenclatural problems became evident for the treatment of taxa in that national park. One of the nomenclatural problems concerns *Isoetes pedersenii*, which was effectively published for the first time by H.P. Fuchs-Eckert (1982) and later accepted by MacLuf et al. (2006a, b). However, in none of the publications it was published with a latin diagnosis as specified in articles 32.1 of the International Code of Botanical Nomenclature of Vienna (McNeill et al. 2007). Fuchs-Eckert (1982) mentioned the existence of a holotype housed in the Herbarium of Paris (P), which was collected in Corrientes, Argentina. The data supplied in this publication is mentioned below:

“*Isoetes pedersenii* (H.P. Fuchs, in schedis) Fuchs-Eckert, Rev., in hb P (24 December 1980) ined.) A



ex) METTENIUS 1859, Fil. Lechl. (2): 36, in textu (Januar 1859), *nomen seminudum*, ex) A. Braun 1862, in Verh. Bot. Ver. Prov. Brandenbeg.”

R.J. Hickey (1985), in his unpublished thesis, updated the neotropical species of *Isoetes*, and provided an English description of *Isoetes pedersenii*. Later, Macluf et al. (2006a, b) performed an analysis with electron microscopy of the microspores of all the species of Southern South America, among which *I. pedersenii* was included. Additional background information on this name was documented in correspondence dated XI.30.1981 sent by Fuchs to Pedersen, which is kept in the Library of the Botany Institute of the Northeast (IBONE). A list of specimens collected by Pedersen and determined by Fuchs was provided in this letter. In the detail of a specimen, the following is specified:

“T.M. Pedersen, no. 8105: *Isoetes pedersenii* H.P. FUCHS, in sched., ined. (TYPUS).”

As part of the study of the Isoetoids that grow in Southern South America and within the boundaries of the Project “Florula of the National Park Mburucuyá”, the name used by Fuchs-Eckert (1982), R.J. Hickey, unpublished data and Macluf et al. (2006a, b), *Isoetes pedersenii*, is described, typified and validated in accordance with the standards imposed by the International Code of Botanical Nomenclature.

#### MATERIALS AND METHODS

Studies were performed on herbarium material borrowed from several herbaria, which are abbreviated in accordance with Holmgren et al. (1990). The material was recovered by warming it in distilled water with some drops of non-ionized detergent. Permanent preparations were made for the anatomical descriptions. This material was dehydrated in an ascending series of tertiary butylic alcohol and was introduced into paraffin (Johansen 1940); transverse sections were made with a rotary microtome at 12–15  $\mu\text{m}$ . These were dyed with saffranine-azure blue (Lique et al. 1996) and were

camera included an Olympus CO11 stereoscopic microscope dissecting scope. Illustrations of features were carried out with a Wild M5 stereoscopic microscope and an Olympus BX 50F binocular microscope. For studies with SEM, the spores were handled with moist brushes without any chemical treatment and placed on double-stick tape on bronze stubs. The samples were coated with gold and examined under a Jeol JSM-35 CF microscope at the SEM laboratory of the Facultad de Ciencias Naturales y Museo, Universidad Nacional de La Plata, La Plata, Argentina.

Megaspore morphology follows the terminology proposed in Tryon and Lugardon (1991).

#### RESULTS


*Isoetes pedersenii* H.P. Fuchs ex Meza T. and Macluf, spec. nov. (Figs. 1–4).

*Cormus erectus, bise trilobatus, 18 mm latus, 11 mm elatus; radices dichotomae. Folia 70–80, erecta, 12–30 cm longa, 4–6 mm basi lata, 1–6 mm lata; alae hyaline et membranaceae, 3 mm basi latae, 2 mm medio latae, 10–25 cm longae (50% per folia longitudinem ascendentes); fasciculi fibrosi praesentia; subula triquetra, atroviridis; stomata praesentia; squamae phyllopodia absentia. Sporangium ellipticum usque circulare, hyalinum, 5–6 mm longum, 3 mm latum, basale. Velum incompletum, revolutus versus is apex. Ligula deltata, auriculata, 1.3 mm longa, 1.85 mm lata. Labium inconspicuum usque absens. Megaspores, 395–411  $\mu\text{m}$  diametro, verrucosae, verrucae 21.3–43.4  $\mu\text{m}$  diametro; zona non dissimilis. Microspores, 30–35  $\mu\text{m}$  longae, 20–24  $\mu\text{m}$  latae, echinatae.*

**Holotypus:** Argentina, Corrientes province: Dep. Mburucuyá, Estancia Santa María, low generally flooded land, 18 March 1967, *Pedersen 8105* (CTES); *Isotypi:* A, C, LP, P, S.

**Etymology:** The species was dedicated by Fuchs to Dr. T.M. Pedersen, collector and expert of the flora from Corrientes who collected the type specimen.

Corm erect, 2 to 3 lobed, 18 mm wide, 11 mm high; roots dichotomously branched. Leaves 70–80, erect, 12–



A NEW SPECIES OF *ISOETES* FROM SOUTH AMERICA




Fig. 1 – *Isoetes pedersenii*. A – Plant habit. B – Cross section of the subula (terminal region of a microphyll), with 4 air chambers, fiber bundles (black circles) and vascular tissue (cut line). C – Detail of the labium folded in order to let the ligule remnant be seen. D – Basal portion of a microphyll with megasporangium. E – Cross section of microsporangium, with trabeculae arranged transversely. F – Detail of the subula apex. (Pedersen 8105, CTES).

to 25 cm long (extending until 85% of the leaf length); fibrous bundles present; subula triquetre, dark green (Fig. 1. F); stomata present; scales and phyllonodia ab-

wide. Labium conspicuous, 3-3.5 × 1-1.5 mm (C). Megaspores subtriangular, trilete, 395 to 411 equatorial diameter, verrucate; verrucae 21.3 to






Fig. 2 – *Isoetes pedersenii*. A-C. Megaspores observed with SEM. A – Proximal view. The laesurae join in the equatorial zone or cingulum. Microspores are observed on the megaspore surface. B – Detail of the proximal structure showing tubercles of different diameter. C – Distal View. (Pedersen 8105, LP).

*Isoetes pedersenii* perhaps is endemic of Argentina, and is known only from the type collection made in Mburucuyá National Park, Corrientes. It grows in flooded land; the presence of stomates and fibrous bundles and the absence of scale leaves suggest an amphibious habit. *Isoetes pedersenii* is probably to be also found in the adjacent regions of Argentina, Paraguay

of subula (microphyl tip). Morphologically, it is similar to *Isoetes panamensis* Maxon and C.V. Morton and to *Isoetes gardneriana* A. Braun, from which it is differentiated mainly by the megaspore ornamentation. *Isoetes panamensis* has cones on the megaspore surface, whereas in *I. gardneriana* the megaspores have tubercles.



# A NEW SPECIES OF *ISOETES* FROM SOUTH AMERICA





Fig. 3 – *Isoetes pedersenii*. A-D. Microspores observed with SEM. A – Proximal view (top) and equatorial view (bottom). B – Major equatorial view showing the echinate-tuberculate ornamentation. The echinae are higher and more densely distributed distally. C – Lower equatorial view showing the supra-laesural expansion (Macluf et al. 2006a). D – Detail of the equatorial surface. The ornamentation consists of cones distributed in the entire surface. Some cones show their apex broken. (*Pedersen 8105*, LP).

eral, and one abaxially, with strongly lignified thick walls and reduced lumen, immersed in a subepidermal (tissue) parenchyma (Fig. 1, B, Fig. 4, C). The presence or absence of peripheral fibers, number and location of *Isoetes* have been considered to be a diagnos-

ton (1985). The presence of this tissue would s be related to the habitat (Pfeiffer 1922, Takamiy 1997). The aquatic species lack, mechanical s tissue and, in terrestrial and amphibious species, it developed. This cannot be completely confirme



## REFERENCES

ARBO MM. 2004. Flórula del Parque Nacional Mburucuyá. In: ACEÑOLAZA FG (Ed), Temas de la biodiversidad del Litoral Fluvial Argentino. INSUGEO. Tucumán, Miscelánea 12.

FUCHS-ECKERT HP. 1982. Zur heutigen Kenntnis von Vorkommen und Verbreitung der südamerikanischen *Isoetes*-Arten. Proceedings of the Koninklike Nederlandse Akademie van Wetenschappen, Series C 85: 205–260.

HALL JB. 1971. Observations on *Isoetes* in Ghana. Bot J Linn Soc 64: 117–139.

HICKEY RJ, MACLUF CC AND TAYLOR WC. 2003. A Re-evaluation of *Isoetes savatieri* Franchet in Argentina and Chile. Amer Fern J 93: 126–136.

HOLMGREN PK, HOLMGREN NH AND BARNETT LC. 1990. Index Herbariorum, 8<sup>th</sup> ed., NYBG, New York USA, 693 p.

JOHANSEN DA. 1940. Plant microtechnique. McGraw-Hill Book Company, Inc., New York.

O nome *Isoetes pedersenii* H.P. Fuchs (Lycophyta) foi validado para a espécie identificada apenas no Parque Nacional de Mburucuyá, em Corrientes na Argentina. Material preservado em herbário foi avaliado com microscópios estereoscópico, de luz branca e eletrônico de varredura. A espécie foi descrita e tipificada. Um diagnóstico e uma discussão sobre a distribuição e relação com a morfologia de outras espécies de



A NEW SPECIES OF *ISOETES* FROM SOUTH AMERICA

- KRAPOVICKAS A. 2000. Troels Myndel Pedersen (26.IX. 1916–5.II.2000). *Bonplandia* 10: 193–196.
- LUQUE R, SOUSA HC AND KRAUS JE. 1996. Métodos de coloração de Roeser (1972) – modificado – E. Kropp (1972), visando a substituição do azul de astra por azul de alcião 8GS ou 8GX. *Acta Bot Bras* 10: 199–212.
- MACLUF C AND PONCE M. 2008. Isoetaceae. In: ZULOAGA ET AL. (Eds), Catálogo de las plantas vasculares del Cono Sur (Argentina, Sur de Brasil, Chile, Paraguay y Uruguay) I: Pteridophyta, Gymnospermae y Monocotyledoneae. *Monographs in Systematic Botany from the Missouri Botanical Garden* 107: 984.
- MACLUF C, MORBELLI MA AND GIUDICE GE. 2006a. Microspore Morphology of *Isoetes* (Lycophyta) species from Southern South America. *Bot Rev* 72: 121–134.
- MACLUF C, MORBELLI MA AND GIUDICE GE. 2006b. Microspore Morphology of *Isoetes* species (Lycophyta) from Southern South America. Part II. TEM analysis of some selected types. *Bot Rev (Lancaster)* 72: 135–152.
- MC NEILL J ET AL. 2007. International Code of Botanical Nomenclature (Vienna Code) adopted by the Seventeenth International Botanical Congress Vienna, Austria, July 2005. In: GANTNER VERLAG AND RUGGELL LICHTENSTEIN (Eds), *Regnum Vegetabile*, 146, XVIII, 568 p.
- PARKER D. 1943. Comparison of aquatic and terrestrial plants of *Isoetes engelmanni* in the Mountain Laguna area. *Am Midl Nat* 30: 452–455.
- PFEIFFER NA. 1922. Monograph of the Isoetaceae. *Bot Gard* 9: 79–232.
- PONCE MM. 1996. Pteridophyta. In: ZULOAGA ET AL. MORRONE O (Eds), Catálogo de las Plantas Vasculares de la República Argentina I: Pteridophyta, Gymnospermae y Angiospermae (Monocotyledoneae). *Monographs in Systematic Botany from the Missouri Botanical Garden* 60: 1–79.
- PRADA C. 1979. Estudio de la anatomía foliar de las Isoetes españolas del género *Isoetes* L. *Lagascalia* 9: 107–114.
- PRADA C AND ROLLER C. 2003. Caracteres diagnósticos de las Isoetes ibéricas en taxones ibéricos de *Isoetes* L. (Isoetaceae, Pteridophyta). *Anales Jard Bot Madrid* 60: 371–386.
- TAKAMIYA M, WATANABE M AND ONO K. 1997. Systematic studies on the genus *Isoetes* (Isoetaceae) in Japan. IV. *Acta Phytotaxon et Geobotánica* 48(2): 89–124.
- TRYON AF AND LUGARDON B. 1991. Spores of the Isoetes (Lycophyta). Surface, wall structure and diversity by electron microscope studies. Springer, New York.