



Anais da Academia Brasileira de Ciências

ISSN: 0001-3765

[aabc@abc.org.br](mailto:aabc@abc.org.br)

Academia Brasileira de Ciências

Brasil

Martins, Maurício L.; Azevedo, Tatiana M.P.; Ghiraldelli, Luciana; Bernardi, Neuza  
Can the parasitic fauna on Nile tilapias be affected by different production systems?  
Anais da Academia Brasileira de Ciências, vol. 82, núm. 2, junio, 2010, pp. 493-500  
Academia Brasileira de Ciências  
Rio de Janeiro, Brasil

Available in: <http://www.redalyc.org/articulo.oa?id=32713482024>

- How to cite
- Complete issue
- More information about this article
- Journal's homepage in [redalyc.org](http://redalyc.org)

[redalyc.org](http://redalyc.org)

Scientific Information System  
Network of Scientific Journals from Latin America, the Caribbean, Spain and Portugal  
Non-profit academic project, developed under the open access initiative



Anais da Academia Brasileira de Ciências (2010) 82(2): 493-500  
(Annals of the Brazilian Academy of Sciences)  
ISSN 0001-3765  
www.scielo.br/aabc

## Can the parasitic fauna on Nile tilapias be affected by different production systems?

MAURÍCIO L. MARTINS<sup>1</sup>, TATIANA M.P. AZEVEDO<sup>1</sup>, LUCIANA GHIRALDELLI<sup>1</sup> and NEUZA BERNARDI<sup>2</sup>

<sup>1</sup>Laboratório AQUOS, Sanidade de Organismos Aquáticos, Departamento de Aquicultura, CCA  
Universidade Federal de Santa Catarina (UFSC), Rod. Admar Gonzaga, 1346, 88040-900 Florianópolis, SC, Brasil

<sup>2</sup>Departamento de Engenharia Química, UFSC, Rod. SC 404, Trindade, 88040-900 Florianópolis, SC, Brasil

Manuscript received on July 2, 2008; accepted for publication on June 26, 2009

### ABSTRACT

This study compared the parasitic fauna on Nile tilapias kept with swine dejects and tilapia from fee fishing with commercial ration. A total of 360 fish were analyzed from August 2003 to July 2004 in a facility situated in Niterói, Trento, Santa Catarina, Brazil. No significant difference was observed in both systems. The parasite fauna in both systems were slightly similar, with the presence of the following parasites: *Trichodina magna* and *T. compacta* (Ciliophora); *Cichlidogyrus sclerosus* and *Cichlidogyrus* sp. (Monogenoidea); and *Lamproglana* sp. (Lernaeidae). Parasitological analyzes did not differ among fishes kept with swine dejects and in fee fishing with commercial diet ( $P > 0.05$ ). The total prevalence of trichodinids in the gills of fish kept with swine dejects was 1.7% as compared to the one kept in fee fishing with commercial diet (0.6%). The Monogenoidea prevalence and mean intensity in the gills and body of fish from fee fishing was 16.5% and 2.6, and compared to other system was 13.2% and 0.8, respectively. This study showed that low stocking density and low water temperature in that region were responsible for the maintenance of good health and lower parasitism rate.

**Key words:** *Oreochromis niloticus*, fee fishing, swine dejects, Monogenoidea, *Trichodina*, prevalence.

### INTRODUCTION

In 1990, an economy based on the integrated production of fish and swine culture has been developed, in which today, the most raised fish species are tilapia (*Oreochromis niloticus*) and carp (*Cyprinus carpio*) (Souza et al. 2003).

During the last years, there has been a significant increase in the facilities that practice the fee fishing system (sportive fish culture). The intensive system of fish maintenance is characterized by an increasing in stocking density, which leads to problems related to inappropriate handling, feeding, infectious and parasitic diseases (Martins et al. 2002, 2008). These problems are

industrial and sportive fish culture (Tavares-Dias 2001a).

In Brazil, Békési (1992) pointed out the first report of parasitism in raised fishes from the Northeast (Souza et al. 2001) and Varella et al. (2003) from the Southeast; Martins et al. (2002) from the Southeast; and Fortes et al. (1998), and Ghiralde (2006a) from the South Brazil.

According to Martins et al. (2002), the most common parasites recorded in the Southeast Brazil are: *Pillulina pillulare* (Dinoflagellida) (Schäperclaus 1981); trichodinids, *Trichodina* (Ciliophora) Fouquet, 1876



poda) Linnaeus, 1758; and, more recently, *Lamprogle-na* von Nordmann, 1832 as the second important copepod parasite in freshwater fish (Piasecki et al. 2004). In the South Brazil, Ghiraldelli et al. (2006a) and Martins and Ghiraldelli (2008) identified trichodinids and Monogenoidea as the most dominant fish-cultured parasites.

Nile tilapias are widely reared in the tropics, representing the second species reared all over the world (Cavichiole et al. 2002). In the Southeast region, the tilapia showed an incidence of 40% incidence of diseases (Martins et al. 2002) that, in several times, influenced its health and haematological parameters (Ghiraldelli et al. 2006b).

There are no epidemiological data concerning ectoparasites from reared tilapia in the South Brazil. To date, comparative field studies among parasitic fauna on tilapia kept in an environment that receives swine dejects and commercial ration are not registered around the world. Thus, this study was carried out for 12 months to compare ectoparasites from tilapia kept with swine dejects and in fee fishing fed commercial diet in the South Brazil.

#### MATERIALS AND METHODS

From August 2003 to July 2004, 360 Nile tilapias were caught by net in Nova Trento town (27°17'09"S, 48°55'47"W), S.C., Brazil. The animals kept with swine dejects (n=180) presented  $26.8 \pm 1.1$  cm of total length and  $385.0 \pm 37.5$  g of mean weight, at a stocking density of 3 fish/m<sup>2</sup>. The others, from fee fishing (n=180), measured  $28.5 \pm 2.0$  cm and weighted  $456 \pm 86.8$  g, at a stocking density of 4 fish/m<sup>2</sup>. During the sampling days, water temperature, dissolved oxygen, pH, alkalinity and transparency were all obtained. A 250 mL water sample from the ponds was frozen for ammonia analyses according to Koroleff (1976).

After sacrifice and biometrical evaluation being sacrificed and evaluated biometrically for parasitological examination (Ethical Committee no. 23080.017229/2004-78/UFSC), according to Ghiraldelli et al. (2006a)


a 0.05 significance rate, and the ANOVA F statistic for a variance analysis, whereas the average comparison of parasite number, in each month and between the both systems, was performed by Tukey test (Mendes 1999).

#### RESULTS

The water quality measured during the whole period did not show a significant difference ( $P > 0.05$ ) when both the pond that received swine dejects and fee fishing were compared (Tables I and II). The mean values of water temperature were the lowest in August 2003 and in May-July 2004; the highest ones were observed in November-December 2003 and in January-February 2004. Although the dissolved oxygen concentration in fee fishing was elevated from August to October 2003, no difference was observed in relation to the swine dejects system. As shown in Table I, pH and alkalinity did not differ in both systems. Although the fish were kept with swine dejects, the ammonia concentration was kept in tolerable levels. Interestingly, no difference was observed between the systems. Swine dejects supplementation in the water was responsible for the decrease in water transparency, as observed in the majority of the analyzed months. No significant difference among parasitological data in both systems was observed.

From 360 examined animals, 66 (18.3%) were parasitized during the whole period, 20 (5.5%) with *Trichodina magna* Van As and Basson, 1989 and *T. compacta* (Protozoa: Ciliophora) Van As and Basson, 1989; 64 (17.8%) with *Cichlydogyrus* Paperna, 1960 and *Cichlydogyrus sclerosus* (Monogenoidea: Dactylogyridae) Paperna and Thurston, 1969 on body and gills; and 4 (1.1%) with *Lamproglana* (Crustacea: Lernaecidae) von Nordmann, 1832 in the gill arches.

The total prevalence rate of trichodinids was higher in fish's gills from the pond with swine dejects than those of fee fishing (Tables I and II). In this case, this protozoan was present only in August and September 2003. When analyzing the averages of the whole period, Monogenoidea prevalence in the gills was higher (16.5%) in fish from fee fishing than those kept with swine dejects (13.2%). Although without significant dif-



PARASITIC FAUNA ON NILE TILAPIAS

TABLE I

Water quality and parasitic characteristics of Nile tilapias (n=180) kept in fee fishing system from Valley of Tijucas River, Santa Catarina, Brazil. T: temperature; O<sub>2</sub>: dissolved oxygen; NH<sub>3</sub>: ammonia; Alkal: alkalinity; Tran: transparency; P: prevalence; MI: mean intensity; G: gill; B: body surface.

Months	Water quality						Monogenoidea				Trichodinid		<i>Lamproglanis</i>	
	T °C	O <sub>2</sub> mg/L	pH	NH <sub>3</sub> mg/L	Alkal mg/L	Tran cm	P (%)		MI		P (%)		P (%)	MI
							G	B	G	B	G	B	G	B
Aug/03	19.3	9.3	7.1	0.500	20	25	60	13	14.6	0	7	13	20	1.4
Sep/03	19.0	8.3	7.0	0	25	22	46	0	6.3	0	0	13	0	0
Oct/03	22.5	8.5	6.5	0.108	30	14	0	26	0	3.2	0	0	0	0
Nov/03	24.4	4.0	6.5	0.103	30	18	26	0	1.5	0	0	0	0	0
Dec/03	28.1	5.8	6.5	0	30	17	6	0	1.0	0	0	0	0	0
Jan/04	26.3	6.4	5.1	0.032	30	15	20	0	2.0	0	0	0	0	0
Feb/04	25.4	7.8	7.4	0.135	30	10	0	0	0	0	0	0	0	0
Mar/04	32.0	6.5	5.6	0.136	20	22	0	0	0	0	0	0	0	0
Apr/04	24.0	3.8	6.0	0.313	50	20	20	0	2.0	0	0	0	0	0
May/04	19.2	3.5	5.1	0.043	40	20	26	0	4.0	0	0	0	0	0
Jun/04	17.3	3.8	6.2	0.160	30	10	0	0	0	0	0	0	0	0
Jul/04	18.3	4.3	6.5	0.131	30	10	0	0	0	0	0	0	0	0
Mean	23.0	6.0	6.3	0.116	30	17	16.5	3	2.6	0.3	0.6	2.2	1.7	0.1

TABLE II

Water quality and parasitic characteristics of Nile tilapias (n=180) kept with swine dejects from Valley of Tijucas River, Santa Catarina, Brazil. T: temperature; O<sub>2</sub>: dissolved oxygen; NH<sub>3</sub>: ammonia; Alkal: alkalinity; Tran: transparency; P: prevalence; MI: mean intensity; G: gills; B: body surface.

Months	Water quality						Monogenoidea				Trichodinid		<i>Lamproglanis</i>	
	T °C	O <sub>2</sub> mg/L	pH	NH <sub>3</sub> mg/L	Alkal mg/L	Tran cm	P (%)		MI		P (%)		P (%)	MI
							G	B	G	B	G	B	G	B
Aug/03	18.2	5.7	7.0	0.500	40	4	66	0	3.3	0	13	0	0	0
Sep/03	20.0	6.8	6.0	0	20	7	80	0	3.3	0	7	7	6.6	2
Oct/03	22.7	5.8	6.5	0.098	20	16	0	0	0	0	0	0	0	0
Nov/03	25.3	3.8	6.0	0.107	30	12	6	0	1.0	0	0	0	0	0
Dec/03	28.8	3.8	6.5	0.072	30	22	0	6	1.0	0	0	0	0	0
Jan/04	26.9	4.1	4.8	0.048	30	18	13	0	1.5	0	0	0	0	0
Feb/04	25.1	5.9	7.8	0.194	30	18	0	0	0	0	0	0	0	0
Mar/04	27.6	5.5	5.4	0.151	25	9	0	0	0	0	0	0	0	0
Apr/04	24.5	3.1	5.5	0.073	80	12	0	0	0	0	0	0	0	0
May/04	19.0	3.1	5.1	0.199	30	10	0	0	0	0	0	0	0	0
Jun/04	18.0	3.2	6.3	0.348	30	10	0	0	0	0	0	0	0	0



Monogenoidea prevalence and mean intensity were higher in August and September 2003 when compared to the other months, not only in fishes from fee fishing, but also in those from the pond with swine dejects. It is not possible to confirm whether there was a relationship between parasite number and prevalence with temperature. But, it was observed that the copepod *Lamproglena* sp. was present in fishes from fee fishing, in August 2003 (prevalence 20% and mean intensity 1.4) and September 2003 (prevalence 6.6% and mean intensity 2).

#### DISCUSSION

The rapid development of aquaculture in Brazil has caused low water quality, handling stress, and inadequate source of feeding, which has led to an increased number of parasites. Other economic losses have occurred as a result of imbalance in the host/parasite/environment system (Ghiraldelli et al. 2006b). The Nile tilapia is one of the most raised fish in the South Brazil. Only in the last decade parasitological studies from raised fish have been reported (Békési 1992, Tavares-Dias et al. 2001a, b, Ghiraldelli et al. 2006a). The presence of trichodinids associated to low water quality or fish mortality was reported in São Paulo (Tavares-Dias et al. 2001a, Martins et al. 2002), Paraná (Vargas et al. 2000) and Ceará States and in the Amazon region (Békési 1992, Andrade et al. 2001).

Trichodinids are considered as commensals, which are found on the skin surface and gills, feeding on mucus, cells, and bacteria. They include an important group of ectoparasites in freshwater fish since their presence is directly related to water quality. Experimental infections with *Trichodina truttae* Müller, 1937 revealed fish mortality as related by Urawa (1992), and with *Trichodina jadranica* Raabe, 1958 caused disease in eel from recirculating aquaculture (Møllergaard and Dalsgaard 1987). Mixed infections, which were observed in this study, corroborated the findings of Barker et al. (2002) who reported *Trichodina murmanica* Polyanski, 1955 and *Gyrodactylus pleuronecti* Cone, 1981

et al. (2000). Their prevalence in August and September 2003 corroborated the observations of Tavares-Dias et al. (2001a) and Martins et al. (2002) in tilapias. Although seasonality in the trichodinid's prevalence was not observed, the highest values were recorded in August and September 2003 for both systems. It can also be mentioned the lack of trichodinids' pathogenicity in this study, as confirmed by Madsen et al. (2000) observations. However, we concluded that the pathogenicity was directly related to the number of parasites on fish (category 1 of Madsen et al. 2000) and is associated to water quality as reported by Afif et al. (2000).

Several species of the genus *Oreochromis* have been introduced in Africa and America with Monogenoidea (Kritsky and Thatcher 1974). In this work, Monogenoidea total prevalence was lower than the one observed by Flores-Crespo et al. (1992). In this work, despite the highest mean intensity in August and September 2003, symptoms of disease were not observed, which was different of what was found by Kaneko et al. (1988). Buchmann et al. (1995) emphasized the importance of Monogenoidea in Danish rainbow trout culture, since some of them occur in a specific season and the others present high values during the whole year. In the present study, tilapias showed low infestation by Monogenoidea and trichodinids, corroborating the results obtained by Tavares-Dias et al. (2001b). The total prevalence rate of Monogenoidea in fishes from fee fishing and pond that received swine dejects was similar to that found by Tavares-Dias et al. (2001b), which was also in fee fishing situated in the Southeast Brazil. But, it must be emphasized that the prevalence was higher (46-80%) in August and September 2003 than that one mentioned on the same fish from the Southeast Brazil. Monogenoidea's mean intensity showed the highest values in the fall and spring, while Tavares-Dias et al. (2001b) reported them in the spring and summer in *P. mesopotamicus* and *L. macrocephalus*.

The genus *Lamproglena* is the most primitive group of Lernaeidae (Thomas and Hameed 1984) and the second major gill freshwater parasite cyclopoid (Piasecki et al. 2004). It occurs in Africa (Marx and



#### PARASITIC FAUNA ON NILE TILAPIAS

ality of *Lamproglena minuta* Capart 1943 in *Puntius binotatus* Valenciennes 1842 and its relation with the fish size observed by Leong (1986), in this study, there was no relation between the parasitism and the sampling months.

According to Duëllou and Erlwanger (1994), *Tilapia rendalli* Boulenger, 1897 is one of the most preferred hosts for *Lamproglena monodi* Capart, 1944, with high values of prevalence and parasite number when compared to the findings of the present study.

Similar prevalence and mean intensity of *Lamproglena* were reported in *Chondrostoma regium* Heckel, 1843 parasitized by *Lamproglena pulchella* von Nordmann 1832 (Saglam 1998). *Lamproglena clariae* (Fryer 1956) is endemic in Africa, and Marx and Avenant-Oldewage (1996) assigned the infestation to unpolluted sites. It can also be added that in the majority of the cases, *Lamproglena* occurs in intensities from 1 to 5 parasites. Moreover, *L. monodi* has been found in *O. niloticus*, reared in the Philippines, at the mean intensity of 1 to 3.5 parasites per fish (Yambot and Lopez 1996), which was similar to this trial. In Brazil, the lack of data in raised fish during the last 15 years does not allow any conclusion on its introduction. The low intensity of parasitism may be responsible for unknown effects on survival and growth of the hosts. Six percent prevalence of *Lamproglena* in tilapia reared in ponds and cages in São Paulo, Brazil (Martins et al. 2002) were not directly associated to mortality.

Except for the lowest dissolved oxygen and the highest ammonia concentration in the pond that received swine dejects, the analyses of water quality was similar in both systems. It must be noted that the presence of swine dejects was probably responsible for the worst parameters. Although the free fishing system had presented a higher fish density than pond with swine dejects, oxygen and ammonia did not increase. This is especially true when analyzing the results obtained by Dhawan and Kaur (2002) that concluded any effects of swine dejects in the water quality. In this study, a great amount of algae was observed in the surface of the pond that received swine dejects. Low temperature was prob-

In this context, Skinner (1982) observed a high infection rate by Monogenoidea in fishes from environments with high amounts of ammonia. This trial did not prove this fact probably because ammonia concentration was not that high. On the contrary, high prevalence rates of trichodinids were associated with temperature in fishes from India (Singhal et al. 1992) and from Finland lakes (Halmetoja et al. 1992). This study was not able to establish a relationship between water quality and parasites. Considering the free system, high stocking density caused a slight increase in the prevalence and number of Monogenoidea. This result is in accordance to Koskivaara et al. (1991) found in roaches from eutrophic and polluted lakes in Finland. Moreover, the mean intensity of this work was significantly higher than that related to roaches. The effects of water quality on the parasite community were also studied in polluted and unpolluted Lake in Central Italy (Bagge and Valtonen 1996). Although direct relationship with our results was not possible, it can be inferred that water quality may affect the parasite community. In fact, Galli et al. (2001) have observed the negative effects of the pollution on the infection with *L. pulchella* in fishes from lakes in Italy. In intensive fish farming, this condition must be considered by the fact that high water flow and high organic matter contents could be better substrates for parasite reproduction.

Parasitologists have been closely observing the mechanisms involved in host response, which depend on interactions among host/parasite/environment. Several studies have been conducted on specific substances (Buchman and Lindstrom 2002). Some of the factors that are responsible for the decrease of fish resistance are: stocking temperature, water quality, stress level, quantity of feeding, age, and natural immunity depending on fish species. Finally, the severity of disease depends on the parasite capacity of infection and environmental quality in which fishes are kept. Data collected here indicate the difference in the parasite numbers observed by tilapias reared in the Brazil (Tavares et al. 2001a, b). In the Southeast Brazil, 20-30°C water temperature might improve parasite reproduction.



#### ACKNOWLEDGMENTS

The authors thank Dr. Walter A. Boeger and Rogério T. Viana (Departamento de Zoologia, Universidade Federal do Paraná, PR, Brasil) who kindly identified Monogenoidea and “Pesca e Lazer Tridapalli” facility for fish donation. This study was financially supported by “Conselho Nacional de Desenvolvimento Científico e Tecnológico/CNPq” – 300281/04-8, 470100/04-4).

#### RESUMO

Este estudo comparou a fauna parasitária de tilápia do Nilo mantida em consorciação com suínos e alimentada com ração comercial mantida em pesque-pague. Um total de 360 peixes foi analisado, entre agosto de 2003 e julho de 2004, em uma propriedade de Nova Trento, Santa Catarina, Brasil. Não houve diferença significativa entre os dois sistemas de cultivo. A fauna parasitária em ambos foi ligeiramente semelhante com a presença de *Trichodina magna* e *T. compacta* (Ciliophora); *Cichlidogyrus sclerosus* e *Cichlidogyrus* sp. (Monogenoidea) e *Lamproglana* sp. (Lernaeidae). A análise parasitológica não mostrou diferença entre os peixes dos dois sistemas. A prevalência total de tricodinídeos nas brânquias dos peixes mantidos com dejetos de suínos foi de 1,7% quando comparada com os do pesque-pague de 0,6%. A taxa de prevalência e a intensidade média de Monogenoidea nas brânquias dos peixes do pesque-pague foi de 16,5% e 2,6, respectivamente comparada com os mantidos com suínos de 13,2% e 0,8, respectivamente. Este estudo demonstrou que a baixa densidade de estocagem de peixes nesta propriedade e a baixa temperatura na região colaboraram para que a saúde dos animais fosse mantida e o nível de parasitismo baixo.

**Palavras-chave:** *Oreochromis niloticus*, pesque-pague, dejetos de suínos, Monogenoidea, *Trichodina*, prevalência.

#### REFERENCES

AFIF SH, THOBIATI AL AND HAZAA MS. 2000. Parasitic gill lesions in Nile tilapia *Oreochromis niloticus* from fish farms in Saudi Arabia. *Assiut Vet Med J* 42(84): 183–194.

ANDRADE SMS, MALTA JCO AND FERRAZ E. 2001. Fauna parasitológica de alevinos de matrinxã *Brycon cephalus* (Günther, 1869) coletados nos rios Negro e Solimões, na

parasite communities of roach (*Rutilus rutilus*). *Parasitology* 112: 499–508.

BARKER DE, CONE DK AND BURT MDB. 2002. *Trichodina murmanica* (Ciliophora) and *Gyrodactylus pleuronecti* (Monogenea) parasitizing hatchery-reared winter flounder, *Pleuronectes americanus* (Walbaum): effects on host growth and assessment of parasite interaction. *J Fish Dis* 25: 81–89.

BÉKÉSI L. 1992. Evaluation of data on ichthyopathological analyses in the Brazilian Northeast. *Ci e Cult* 44: 400–403.

BUCHMANN K AND LINDESTROM T. 2002. Interactions between monogenean parasites and their fish hosts. *Int J Parasitol* 32: 309–319.

BUCHMANN K, ULDAL A AND LYHOLT HCK. 1995. Parasite infections in Danish trout farms. *Acta Vet Scandinavica* 36: 283–298.

BUSH AO, LAFFERTY KD, LOTZ JM AND SHOSTAK W. 1997. Parasitology meets ecology on its own terms: Margolis et al. revisited. *J Parasitol* 83: 575–583.

CAVICHIOLO F, VARGAS L, RIBEIRO RP, MOREIRA HLM AND LEONARDO JM. 2002. Níveis de suplementação de vitamina C na ração sobre a ocorrência de ectoparasitas, sobrevivência e biomassa em alevinos de tilápia do Nilo (*Oreochromis niloticus* L.). *Acta Scientiarum* 24: 957–964.


DHAWAN A AND KAUR S. 2002. Effect of swine dejects on water quality and polyculture of carp species during winter and summer. *Aquac Int* 10: 297–307.

DOUËLLOU L AND ERLWANGER KH. 1994. Crustacean parasites of fishes in Lake Kariba, Zimbabwe, preliminary results. *Hydrobiologia* 287: 233–242.

FLORES-CRESPO J, VELARDE FI, FLORES-CRESPO R AND PELAEZ CGV. 1992. Variación estacional de *Dactylogyrus* sp. en dos unidades productoras de tilapia del estado de Morelos. *Tecnica Pecuaria en Mexico* 30: 109–118.

FORTES E, HOFFMANN RP AND SCARIOT J. 1998. *Lernaea cyprinacea* Linnaeus, 1758 (Crustacea, Copepoda) parasitando peixes de água doce da grande Porto Alegre, RS, Brasil. *Rev Bras Med Vet* 20: 64–65.

GALLI P, CROSA G, BERTOGLIO S, MARINIELLO L, ORTIS M AND D'AMELIO S. 2001. Populations of *Lamproglana pulchella* von Nordmann 1832 (Copepoda: Eudactylinidae) in cyprinid fish in rivers with different pollution



#### PARASITIC FAUNA ON NILE TILAPIAS

- sites communities from *Oreochromis niloticus* raised in the State of Santa Catarina, Brazil. J Fish Aquat Sci 1: 181–190.
- GHIRALDELLI L, MARTINS ML, YAMASHITA MM AND JERÔNIMO GT. 2006b. Ectoparasites influence on the haematological parameters of Nile tilapia and carp cultured in the State of Santa Catarina, South Brazil. J Fish Aquat Sci 1: 270–276.
- HALMETOJA A, VALTONEN ET AND TASKINEN J. 1992. Trichodinids (Protozoa) on fish from central finnish lakes of differing water quality. Aqua Fenica 22: 59–70.
- KANEKO JJ, YAMADA R, BROCK JA AND NAKAMURA RM. 1988. Infection of tilapia, *Oreochromis mossambicus* (Trewavas), by a marine monogenean, *Neobenedenia melleni* (MacCallum, 1927) Yamaguti, 1963 in Kaneohe Bay, Hawaii, USA, and its treatment. J Fish Dis 11: 295–300.
- KOROLEFF F. 1976. Determination of nutrients. In: GRASSHOF K (Ed), Methods of Seawater Analysis. Verlag Chemie Weinheim, Germany, p. 117–181.
- KOSKIVAARA M, VALTONEN ET AND PROST M. 1991. Seasonal occurrence of gyrodactylid monogeneans on the roach (*Rutilus rutilus*) and variation between four lakes of differing water quality in Finland. Aqua Fenica 21: 47–55.
- KRITSKY DC AND THATCHER VE. 1974. Monogenetic trematodes (Monopisthocotylea: Dactylogyridae) from freshwater fishes of Colombia, South America. J Helminthol 48: 59–66.
- KUMARI P, KHERA S AND GUPTA NK. 1989. On six new species of the genus *Lamproglena* Nordmann (Copepoda: Eudactylinidae) ectoparasitic on freshwater fishes of India. Res Bull Panjab Univ Sci 40: 9–23.
- LEONG TS. 1986. Seasonal occurrence of metazoan parasites of *Puntius binotatus* in na irrigation canal. Pulau Pinang, Malaysia. J Fish Biol 28: 9–16.
- MADSEN HCK, BUCHMANN K AND MELLERGAARD S. 2000. Association between trichodiniasis in eel (*Anguilla anguilla*) and water quality in recirculating systems. Aquaculture 187: 275–281.
- MALTA JCO, GOMES ALS, ANDRADE SMS AND VARELLA AMB. 2001. Infestações maciças por acantocéfalos *Neoechinorhynchus buttnerae* Golvan, 1956, (Eoacanthocephala: Neoechinorhynchidae) em tambaquis jovens.
- MARTINS ML AND GHIRALDELLI L. 2008. *Trichodinids* magna Van As and Basson, 1989 (Ciliophora: Peritrichia) from cultured Nile tilapia in the State of Santa Catarina, Brazil. Braz J Biol 68: 169–172.
- MARTINS ML, ONAKA EM, MORAES FR, BOZELLI R, PAIVA AMFC AND GONÇALVES A. 2002. Recirculating systems on parasitic infections of freshwater raised fish in the State of São Paulo, Brazil. Acta Scientiarum 24: 9–12.
- MARTINS ML, MOURIÑO JLP, AMARAL GV, VIEIRA L, DOTTI G, BEZERRA AJM, PEDROTTI FS, JERÔNIMO GT, BUGLIONE NETO CC AND PEREIRA JR C. 2008. Haematological changes in Nile tilapia experimentally infected with *Enterococcus* sp. Braz J Biol 68: 631–634.
- MARX HM AND AVENANT-OLDEWAGE A. 1996. Description of *Lamproglena clariae* Fryer, 1956 (Copepoda: Lernaecidae) with notes on its occurrence and distribution. Crustaceana 69: 509–523.
- MELLERGAARD S AND DALSGAARD I. 1987. Diseases and problems in Danish eel farms. Aquaculture 67: 139–144.
- MENDES PP. 1999. Estatística Aplicada a Aquicultura. Editora Bagaço, 265 p.
- PIASECKI W, GOODWIN AE, EIRAS JC AND NOWAKI M. 2004. Importance of copepoda in freshwater fish diseases. Biological Studies 43: 193–205.
- SAGLAM N. 1998. Investigation of *Lamproglena pinnata* (Nordmann, 1832) on *Capoeta trutta* and *Chondrostoma toxostoma* caught in Keban Dam Lake (Elazig, Turkey). Appl Ichthyol 14: 101–103.
- SINGHAL RN, JEET S AND DAVIES RW. 1986. The relationships between changes in selected physico-chemical properties of water and the occurrence of fish parasites in Haryana, India. Trop Ecol 27: 1–9.
- SKINNER RH. 1982. The interrelation of water quality, parasites and gill pathology of some fishes from Biscayne Bay, Florida. Fish Bull 80: 269–280.
- SOUZA FJ, SCHAPPO CL, TAMASSIA STJ AND BOZELLI R. 2003. Estudo de competitividade da piscicultura em Vale do Itajaí. Instituto Cepa/SC/Epagri/Acaq, 76 p.
- TAVARES-DIAS M, MARTINS ML AND MORAES FR. 2008. Fauna parasitária de peixes oriundos de “pesque-paga” município de Franca, São Paulo, Brasil. I. Protozoários. Rev Bras Zool 18: 67–79.
- TAVARES-DIAS M, MORAES FR, MARTINS ML AND MORAES FR. 2008b. Fauna parasitária de peixes oriundos de “pesque-paga” município de Franca, São Paulo, Brasil. II. Nematódeos. Rev Bras Zool 18: 81–91.





- THOMAS S AND HAMEED S. 1984. Description of new species of *Lamproglena* (Copepoda: Lernaeidae) from Kerala. Indian J Fish 31: 223–227.
- URAWA S. 1992. *Trichodina truttae* Mueller, 1937 (Ciliophora: Peritrichida) on juvenile Chum Salmon (*Oncorhynchus keta*): pathogenicity and host-parasite interactions. Gyobyo Kenkyu 27: 29–37.
- VARELLA AMB, PEIRO SN AND MALTA JCO. 2003. Monitoramento da parasitofauna de *Colossoma macropomum* (Cuvier, 1818) (Osteichthyes: Characidae) cultivado em tanque-rede em um lago de várzea na Amazônia Central. XII Simp Bras Aqüic, Goiânia 2: 95–106.
- VARGAS L, POVH JA, RIBEIRO RP AND MOREIRA HLM. 2000. Prevalência de ectoparasitos en tilapia del Nilo (*Oreochromis niloticus*) de origen tailandesa de Maringá, Paraná. Arq Ciên Vet Zool Unipar 3: 32–37.
- YAMBOT AV AND LOPEZ EA. 1996. Gill parasite, *Lamproglena monodi* Capart, infecting the Nile tilapia, *Oreochromis niloticus* L., cultured in the Phillippines. Proceedings Third Symp Dis Asian Aquac, 26 Jan to 2 Feb, Bangkok, Thailand, p. 175–177.