



Anais da Academia Brasileira de Ciências

ISSN: 0001-3765

aabc@abc.org.br

Academia Brasileira de Ciências

Brasil

Cardoso, Fernanda C.G.; Salvalaggio, Ana P.B.; Marques, Márcia C.M.

Population structure of *Rudgea parquoides* (Rubiaceae), a shade-tolerant shrub species, in Southern
Brazil

Anais da Academia Brasileira de Ciências, vol. 82, núm. 3, septiembre, 2010, pp. 637-642

Academia Brasileira de Ciências

Rio de Janeiro, Brasil

Available in: <http://www.redalyc.org/articulo.oa?id=32717619011>

- How to cite
- Complete issue
- More information about this article
- Journal's homepage in redalyc.org

redalyc.org

Scientific Information System

Network of Scientific Journals from Latin America, the Caribbean, Spain and Portugal

Non-profit academic project, developed under the open access initiative



Anais da Academia Brasileira de Ciências (2010) 82(3): 637-642
(Annals of the Brazilian Academy of Sciences)
ISSN 0001-3765
www.scielo.br/aabc

Population structure of *Rudgea parquoides* (Rubiaceae), a shade-tolerant shrub species, in Southern Brazil

FERNANDA C.G. CARDOSO, ANA P.B. SALVALAGGIO and MÁRCIA C.M. MARQUES

Laboratório de Ecologia Vegetal, Departamento de Botânica, Setor de Ciências Biológicas
Universidade Federal do Paraná, Caixa Postal 19031, 81531-980 Curitiba, PR, Brasil

Manuscript received on April 28, 2009; accepted for publication on April 12, 2010

ABSTRACT

The expectations that shade-tolerant forest species show 1) a population structure composed by a high amount of small individuals, and 2) biomass allocation for diameter higher than for height growth, were tested for *Rudgea parquoides*, a typical shrub in Southern Brazil. We described the size structure (height and stem diameter) and allometrical relationship of a *R. parquoides* population by counting and measuring all the individuals in a 725m² area in the municipal forest of Curitiba (25°25'S; 49°19'W). A total of 916 individuals (12,634 ind.ha⁻¹) were recorded in the area. The first expectation was supported, since distribution by height and diameter classes showed a predominance of small individuals (skewness coefficients > 1). On the other hand, the regression between height and stem base diameter showed slope $\beta < 1$, which indicates that growth in height is higher than in diameter, not supporting the second expectation. The results show that life strategies in shade-tolerant species may imply in more trade-off combinations than previously described.

Key words: allometry, *Araucaria* forest, size distribution, subtropical forest, understorey.

INTRODUCTION

Plant populations can be described in terms of age, size and shape of its individuals and genetic structure (Harper and White 1974). Differences in size can be caused by factors such as age, genetic variability, resources heterogeneity, competition, effect of herbivore, parasites or pathogens and disturbance (Weiner and Solbrig 1984, Coomes et al. 2003, Muller-Landau et al. 2006). If size structure is related to the age of individuals, it gives information on the current regenerating capacity, past disturbances (Knowles and Grant 1983, Agren and Zackrisson 1990) and future expected occurrences (Odum 2001).

Light quantity and quality are strong factors determining plant growth and population structure in plant

Poorter et al. 2006). Irradiance can fluctuate from 1% of light at the forest floor to 100% above the canopy (Poorter et al. 2006). To take most advantage of the available light, species partition access to light through trade-offs between growth-rate, stem-persistence and shade-tolerance (Wright 2002, Falster and Westoby 2003).

For tropical plant communities, it is suggested that gap-dependent tree species are usually characterized by size distributions with many large individuals and a long tail of relatively rare, small individuals with large diameter, seed and seedling mortalities (Wright et al. 2000). On the other hand, size distributions with many small individuals and a long tail of relatively rare, large individuals characterized shade-tolerant species with opposite traits (Poorter et al. 1996, Kohira and Niimura 2002, Wright et al. 2003). The



The dimensional relationships (allometry) in plants have been used as an indicator of growth strategy of trees (Niklas 1994, Bohlman and O'Brien 2006). Tree architecture is a strong determinant of mechanical safety (van Gelder et al. 2006). As shade-tolerant species, understorey trees have morphologies that increase light interception and persistence by allocating more biomass for diameter growth than height. In contrast, saplings from canopy trees allocate their biomass for efficient height growth and pass through the understorey to reach the high-light environment of the canopy (King 1990, van Gelder et al. 2006). Even though this relationship is well known for trees (King 1990), no information is available for understorey shrubs. It is supposed that, as a shade-tolerant species, shrubs increase light interception and persistence by allocating more biomass for stem (or stems) radial growth.

Rudgea parquoides (Cham.) Muell. Arg. (Rubiaceae) is an endemic shrub of Araucaria forest, occurring in South and Southeast Brazil, as well as in portions of Paraguay and Argentina (Smith and Downs 1956). Its habitat comprises the understorey of the densest regions of *Araucaria* forest and, in Southern Brazil, is frequently one of the most important species in terms of abundance (Kozera et al. 2006). Here, we studied the population of *R. parquoides*, a shade-tolerant species, in Southern Brazil. Specifically, we addressed the following questions: 1) if this population shows a juvenile-dominated size structure; 2) if the relationship between diameter and height indicates higher biomass allocation for diameter.

MATERIALS AND METHODS

This study took place in a 50-ha *Araucaria* forest remnant in Curitiba, Eastern Paraná State, Brazil (25°25'S, 49°19'W, 900 m a.s.l.). The *Araucaria* forest is distributed in Southern Brazil, Eastern Paraguay and Northern Argentina. This rain forest is dominated by the endemic species *Araucaria angustifolia* (Bertol.) Kuntze, and lies in a transition zone between tropical and temperate climates, with rainfall throughout the whole year and moderate temperatures (Hueck 1953). Since monthly

only a small fraction of their former range and are rapidly disappearing, with some contiguous remaining forests currently found in the state of Paraná, Southern Brazil (e.g. 1.3% in Paraná, Brites and Castella 2004). These forests show particular features by virtue of its ecotonal location (Marques et al. 2004), but little is known about its local species dynamics (Oliveira and Pillar 2004).

The climate in the region is Cfb in Köppen's classification system, i.e. humid subtropical and mesothermic with mild winters and occasional hard frosts and no pronounced dry season (Iapar 1978). The average monthly temperature ranges from 13°C (in June to July) to 21°C (in February). Annual precipitation during the 1970-95 period averaged 1389 mm, with greatest rainfall in January and February and least in July and August (Marques et al. 2004).

The studied fragment is characterized by having three distinct strata, with Lauraceae species in the superior layer and many Rubiaceae and Myrtaceae species in the intermediate and inferior layers. The only emergent species occurring is *Araucaria angustifolia* which, with its unique crown architecture (candelabrum), gives a peculiar physiognomic aspect to this kind of forest (Dittrich et al. 1999, Kozera et al. 2006).

The study was conducted in a 725 m² plot, subdivided in 29 (5×5 m) non-contiguous sub-plots (two blocks apart 200 m from each other). All individuals of *Rudgea parquoides* were marked and had their height and stem base diameter measured. As all juveniles and adults (height >32.2 cm) were branched at a minimum of five centimeters above soil level, only one measure of stem diameter was necessary for each individual.

The data were compiled and analyses were performed to assess the pattern of size structure and growth of the individuals in the population. To evaluate the frequency distribution for height and diameter, we first tested for normality and log-normality of data using Shapiro-Wilk and Kolmogorov-Smirnov-Lilliefors (KSL) tests, respectively (Zar 1999). After that, we chose among coefficient of variation, Gini coefficient or skewness coefficient for verifying the size distribution



POPULATION STRUCTURE OF A SHADE-TOLERANT SHRUB

the size distribution has long tails to the left ($g_1 < 0$), or to the right ($g_1 > 0$), or whether the distribution is symmetrically bell-shaped ($g_1 = 0$). We presented size distribution in size classes (following Vandermeer 1978) because they are more comparable to the other population studies.

To examine the growth pattern, we verified the relationship between height and diameter by performing a standardized major axis (SMA). This analysis is indicated for allometrical relationships in plants and transform (log) the equation $y = \gamma \cdot x^\beta$ to a linear regression (Warton et al. 2006). The 1:1 slope ($\beta = 1$) in the height (x) by diameter (y) regression indicates the symmetrical growth in height and diameter; $\beta > 1$ indicates a higher growth in diameter than in height, and $\beta < 1$ shows a higher growth in height than in diameter. We also performed SMA for juveniles and adults separately to assess possible differences in growth pattern among the stages.

RESULTS

A total of 916 *Rudgea parquoides* individuals was recorded, representing an estimated density of 12,634 ind.ha⁻¹. The median height of the population was 44.5 cm (maximum = 370 cm; minimal = 1.42 cm), and the median diameter was 0.54 cm (maximum = 4.56 cm; minimal = 0.06 cm).

Height and diameter were not normally distributed ($W = 0.84$ and 0.85 , respectively, $P < 0.001$) but approximately log-normally distributed ($D = 0.09$, $P < 0.01$). The skewness coefficient (g_1) for height and diameter were, respectively, $g_1 = 1.12$ and $g_1 = 1.37$, confirming the distribution with a long tail to the right (all $g_1 > 0$). The distribution by height and diameter classes (Figs. 1a and 1b) showed a predominance of individuals in the first classes (1.42–32.2 cm in height and 0.06–0.38 cm in diameter).

Plant height and stem base diameter were highly related ($r^2 = 0.76$; $P < 0.0001$; Fig. 2), which is expected for all allometrical relationships. Nevertheless, the slope (\pm CI) was lower than 1 (0.74 ± 0.02), which

this tendency was confirmed (juveniles: $r^2 = 0.76$; $P < 0.0001$; slope = 0.92 ± 0.08 ; adults: $r^2 = 0.76$; $P < 0.0001$; slope = 0.97 ± 0.05).

DISCUSSION

The expectations that shade-tolerant species show a juvenile-dominated size structure and allocate more mass for diameter than for height were partially supported for the studied population of *Rudgea parquoides*. These results show that the growth strategies among shade-tolerant shrubs and trees, as pointed out for several tropical and sub-tropical species (Went 1949, al. 2003, Souza 2007).

The *R. parquoides* population showed a positive skewness coefficient for height and diameter, which represents a long tail of individuals in the right side of the curve. Since larger individuals are distributed in the long tail, the population also fitted a “J-reverse” distribution in size classes (Leak 1965). Both situations indicate a strong hierarchy among individuals (Went and Solbrig 1984), where a small number of large individuals impose a stronger competitive pressure on small individuals. This behavior has been associated with shade-tolerant tree species, which have size distributions with many small individuals and a long tail of rare, large individuals, the larger ones contributing in a more significant way to the population total biomass (Poorter et al. 1996, Kohira and Ninomiya 2003, et al. 2003). The juvenile dominated size structure of *parquoides* indicates that understory shrubs also grow slowly, which results in the seedling permanence on the forest floor for long periods.

The high density of seedlings and juveniles in the population reflects the occurrence of constant and frequent reproductive events (Ramirez and Arroyo 1990) and a seedling bank (Harper and White 1974). In fact, the population produced fruits during the two preceding years of the demographic study (Marques et al. 2004). They are animal-dispersed between December and March, a period with the best temperature and humidity conditions for germination (Marques et al. 2004). Factors




Fig. 1 – Number of individuals in height (a) and diameter (b) classes of a *Rudgea parquoides* (Rubiaceae) population in an *Araucaria* forest fragment in Curitiba, Southern Brazil.

community, could cause mortality of small individuals, reducing the density of larger individuals in the studied population.

R. parquoides plants showed an allometric growth, demonstrated by the strong relationship between height and diameter. Tropical species often show significant regressions between height and diameter (Alvarez-Buylla and Martinez-Ramos 1992, Santos 2000). This trend seems to be genetically determined (Santos 2000) and related to an equal resource distribution for growth in width and height (Niklas 1994). It was expected that, as

light interception and persistence (King 1990). Nevertheless, the studied population (all stages) tended to allocate higher biomass to height than to diameter (slope < 1), although juveniles and adults showed a marginally 1:1 allometric relationship. The higher diameter allocation is believed to improve stem persistence (Wright et al. 2003, Falster and Westoby 2005). This result suggests that, despite shade-tolerant tree species allocate more biomass to stem growth to sustain the weight of the branches and leaves (Poorter et al. 2006), in shade-tolerant shrub species, the stem biomass can be inferior



POPULATION STRUCTURE OF A SHADE-TOLERANT SHRUB




Fig. 2 – Relationship between stem diameter and height of the *Rudgea parquoides* (Rubiaceae) population in an Araucaria forest fragment in Curitiba, Southern Brazil.

Whereas light is a limiting resource for plants, the growth and survival in the intensely shaded understorey demand a complex strategy to optimize light uptake and, thus, allow persistence and reproduction (Wright et al. 2003). Even if persistence (measured by higher diameter growth) is unimportant, the population structure dominated by small individuals suggests a considerable auto-regenerating potential for *R. parquoides*. Then, regarding growth and size structures, life strategies for shade-tolerant species may imply in more trade-off combinations than previously described.

ACKNOWLEDGMENTS

Thanks to the Prefecture of the Municipality of Curitiba for permission to carry out this research in Barigüi Park. Thanks to F.H. Mello Leitão for helping in the data organization. A.P.B. Salvallaggio received support from the program Iniciação Científica at Universidade Federal do Paraná.

RESUMO

Foram testadas duas hipóteses relativas a espécies florestais tolerantes à sombra em uma população de *Rudgea parquoides*, um arbusto característico do sul do Brasil: 1) espécies tole-


que para o crescimento em altura. Foram descritas estruturas de tamanho (altura e diâmetro do tronco) e alométricas através da contagem e medição de todos os indivíduos de *R. parquoides* em uma área de 725m² no município de Curitiba (25°25'S; 49°19'W). Um total de 916 indivíduos (12.634 ind.ha⁻¹) foi medido nesta área. A primeira hipótese foi aceita, uma vez que a distribuição em classes de diâmetro mostrou um predomínio de indivíduos pequenos (coeficiente de assimetria >1). Por outro lado, a relação entre altura e diâmetro na base do caule mostrou inclinação $\beta < 1$, o que indica que o crescimento em altura é mais rápido do que o crescimento em diâmetro, não suportando a segunda hipótese. Esses resultados mostram que as estratégias de crescimento em altura e diâmetro em espécies tolerantes à sombra podem implicar em outras relações alométricas além das já descritas.

Palavras-chave: alometria, floresta com Araucária, distribuição de tamanho, floresta subtropical, sub-bosque.

REFERENCES

- AGREN J AND ZACKRISSON O. 1990. Age and size structure of *Pinus sylvestris* populations on mines in central and northern Sweden. *J Ecol* 78: 1049–1062.
- ALVAREZ-BUYLLA ER AND MARTINEZ-RAMOS M. 2000. Demography and allometry of *Cecropia obtusifolia*, a tropical pioneer tree – an evaluation of the climax-succession paradigm for tropical forests. *J Ecol* 80: 275–290.
- AUGSPURGER CK. 1983. Seed dispersal of tropical trees: *Platypodium elegans*, and the escape of its seedlings from fungal pathogens. *J Ecol* 71: 759–771.
- BENDEL RB, HIGGINS SS, TEBERG JE AND PYLON M. 1989. Comparison of skewness coefficient, coefficient of variation, and Gini coefficient as inequality measures within populations. *Oecologia* 78: 394–400.
- BOHLMAN SA AND OBRIEN ST. 2006. Allometry, growth, stature and regeneration requirements of 65 tree species on Barro Colorado Island, Panama. *J Trop Ecol* 22: 131–136.
- BRITEZ RM AND CASTELLA PR. 2004. A Floresta de Araucária no Paraná: conservação e diagnóstico de remanescentes florestais. Brasília: Ministério do Meio Ambiente, 233 p.

CLARK DB AND CLARK DA. 1987. Population dynamics and microhabitat distribution of *Dypterix panamensis* in a tropical rain forest. *Ecology* 68: 1000–1010.



- butions in natural forests from following scaling relationships. *Ecol Lett* 6: 980–989.
- DITTRICH VAO, KOZERA C AND SILVA SM. 1999. Composição florística de epífitos vasculares em área de Floresta Ombrófila Mista (Parque Barigüi), Curitiba, PR. *Iheringia Ser Bot* 52: 11–22.
- FALSTER DS AND WESTOBY M. 2005. Alternative height strategies among 45 dicot rainforest species from tropical Queensland, Australia. *J Ecol* 93: 521–535.
- FRANKLIN J. 2003. Regeneration and growth of pioneer and shade-tolerant rain forest trees in Tonga. *New Zeal J Bot* 41: 669–684.
- HARPER JL AND WHITE J. 1974. The demography of plants. *Annu Rev Ecol Sys* 5: 419–463.
- HUECK K. 1953. Distribuição e habitat natural do Pinheiro-do-Paraná (*Araucaria angustifolia*). *Bol Fac Fil Ci Let USP- Botânica* 10: 1–24.
- IAPAR. 1978. Cartas climáticas básicas do estado do Paraná. Fundação Instituto Agrônômico do Paraná, Londrina.
- JANZEN DH. 1970. Herbivores and the number of tree species in tropical forest. *Am Nat* 940: 501–528.
- KING DA. 1990. Allometry of saplings and understorey trees of a Panamanian Forest. *Funct Ecol* 4: 27–32.
- KNOWLES P AND GRANT M. 1983. Age and size structure analyses of engelmann spruce, ponderosa pine, lodgepole pine, and limber pine in Colorado. *Ecology* 64: 1–9.
- KOHIRA M AND NINOMIYA I. 2003. Detecting tree populations at risk for forest conservation management: using single-year vs. long-term inventory data. *Forest Ecol Manag* 174: 423–435.
- KOZERA C, DITTRICH VAO AND SILVA SM. 2006. Fitossociologia do componente arbóreo de um fragmento de Floresta Ombrófila Mista Montana, Curitiba, PR, Brasil. *Rev Floresta* 36: 225–237.
- LEAK WB. 1965. The J-shaped probability distribution. *Forest Sci* 11: 405–19.
- MARQUES MCM, ROPER JJ AND SALVALAGGIO APB. 2004. Phenological patterns among plant life forms in a Subtropical Forest in Southern Brazil. *Plant Ecol* 173: 191–201.
- MULLER-LANDAU HC ET AL. 2006. Comparing tropical forest tree size distributions with the predictions of metabolic ecology and equilibrium models. *Ecol Lett* 9: 589–602.
- ODUM E. 2001. Fundamentos de Ecologia. Lisboa: Fundação Calouste Gulbenkian, 434 p.
- OLIVEIRA JM AND PILLAR VD. 2004. Vegetation dynamics on mosaics of Campos and Araucaria forest between 1974 and 1999 in Southern Brazil. *Community Ecol* 5: 197–202.
- POORTER L, BONGERS F, VAN ROMPAEY RSAR AND KLERK M. 1996. Regeneration of canopy tree species at five sites in West African moist forest. *Forest Ecol Manag* 84: 61–69.
- POORTER L, BONGERS L AND BONGERS F. 2006. Architecture of 54 moist-forest tree species: traits, trade-offs and functional groups. *Ecology* 87: 1289–1301.
- RAMIREZ N AND ARROYO MK. 1990. Estructura poblacional de *Copaifera pubilifera* Benth. (Leguminosae; Caesalpinioideae) en los altos llanos centrales de Venezuela. *Biotropica* 22: 124–132.
- SANTOS FAM. 2000. Growth and leaf demography of two Cecropia species. *Rev Bras Bot* 23: 133–141.
- SMITH LB AND DOWNS RJ. 1956. Resumo preliminar das Rubiaceas de Santa Catarina. *Sellowia* 7: 13–86.
- SOUZA AF. 2007. Ecological interpretation of multiple population size structures in trees: The case of *Araucaria angustifolia* in South America. *Austral Ecol* 32: 524–533.
- VAN GELDER HA, POORTER L AND STERCK FJ. 2006. Wood mechanics, allometry, and life-history variation in a tropical rain forest tree community. *New Phytologist* 171: 367–378.
- VANDERMEER J. 1978. Choosing categories size in a stage projection matrix. *Oecologia* 32: 79–84.
- WARTON DI, WRIGHT IJ, FALSTER DS AND WESTOBY M. 2006. Bivariate line-fitting methods for allometry. *Biol Rev* 81: 259–291.
- WEINER J AND SOLBRIG OT. 1984. The meaning and measurement of size hierarchies in plant populations. *Oecologia* 61: 334–336.
- WRIGHT SJ. 2002. Plant diversity in tropical forests: a review of mechanisms of species coexistence. *Oecologia* 130: 1–14.
- WRIGHT SJ AND VAN SCHAIK CP. 1994. Light and the phenology of tropical trees. *Am Nat* 143: 193–199.
- WRIGHT SJ, MULLER-LANDAU H, CONDIT R AND HUBBELL SP. 2003. Gap-dependent recruitment, realized vital rates, and size distributions of tropical trees. *Ecology*