



Anais da Academia Brasileira de Ciências

ISSN: 0001-3765

aabc@abc.org.br

Academia Brasileira de Ciências

Brasil

KELLNER, ALEXANDER W.A.; AZEVEDO, SERGIO A.K.; MACHADO, ELAINE B.; DE CARVALHO, LUCIANA B.; HENRIQUES, DEISE D.R.

A new dinosaur (Theropoda, Spinosauridae) from the Cretaceous (Cenomanian) Alcântara Formation, Cajual Island, Brazil

Anais da Academia Brasileira de Ciências, vol. 83, núm. 1, março, 2011, pp. 99-108

Academia Brasileira de Ciências

Rio de Janeiro, Brasil

Available in: <http://www.redalyc.org/articulo.oa?id=32717681005>


- How to cite
- Complete issue
- More information about this article
- Journal's homepage in redalyc.org

redalyc.org

Scientific Information System

Network of Scientific Journals from Latin America, the Caribbean, Spain and Portugal

Non-profit academic project, developed under the open access initiative



A new dinosaur (Theropoda, Spinosauridae) from the Cretaceous (Cenomanian) Alcântara Formation, Cajual Island, Brazil

ALEXANDER W.A. KELLNER, SERGIO A.K. AZEVEDO, ELAINE B. MACHADO,
LUCIANA B. DE CARVALHO and DEISE D.R. HENRIQUES

Departamento de Paleontologia e Geologia, Museu Nacional/UFRJ, Quinta da Boa Vista s/n
São Cristóvão, 20940-040 Rio de Janeiro, Brasil

Manuscript received on November 25, 2010; accepted for publication on December 22, 2010

ABSTRACT

A new spinosaurid taxon, *Oxalaia quilombensis* gen. et sp. nov., is described based on the anterior part of a snout and a fragment of a maxilla. These specimens were collected at the Laje do Coringa site, Late Cretaceous (Cenomanian) of the São Luis Basin. Unlike *Cristatusaurus* and *Suchomimus*, *Oxalaia quilombensis* lacks serrations on the teeth. The new species differs from *Angaturama limai* by having the anterior part of the premaxillae more expanded and lacking a sagittal premaxillary crest. It further differs from *Spinosaurus* cf. *S. aegyptiacus* and the Algerian spinosaurid by the rounder shape of the terminal expansion. Furthermore, *Oxalaia quilombensis* has one functional tooth followed by two replacement teeth, a feature not previously observed in theropods. *Oxalaia quilombensis* appears to be more closely related to the spinosaurids found in North Africa than to the Brazilian members of this clade and thus further increases the diversity of these enigmatic predatory dinosaurs in this country.

Key words: Dinosauria, Spinosauridae, *Oxalaia quilombensis*, Cenomanian, Brazil.

INTRODUCTION

Overall the dinosaur record from Brazil is extremely meager and consists mostly of isolated and incomplete remains (see Kellner and Campos 2000 for a review). In terms of non-avian theropods, other than Cretaceous footprints (e.g., Leonardi 1994) and fossilized feathers (e.g., Kellner et al. 1994), only seven species have been described so far: two from the Triassic deposits of Rio Grande do Sul whose phylogenetic position has been questioned (e.g., Galton 2000, Langer 2004, Langer et al. 2009, Bittencourt and Kellner 2009), one abelisaurid from the Late Cretaceous deposits of Mato Grosso (Gibson et al. 1997, Kellner and Campos 2002, Candeiro et al. 2006, Weska 2006) and four from the Aptian-Albian

Romualdo Formation of the Araripe Basin (Martins et al. 1996, Kellner and Campos 1996, 1999, Kellner and Naish et al. 2004).

Here we report on a new theropod, *Oxalaia quilombensis* gen. et sp. nov., the first formally described dinosaur species from Brazilian Cenomanian (early Cretaceous) deposits. The remains of this taxon were collected from the locality known as Laje do Coringa at the Cajual Island, Maranhão State (Corrêa et al. 1997). This site is one of the few bone-beds from the Cenomanian and has yielded hundreds of isolated elements (Corrêa and Schultz 2002, Van Tomme et al. 2008), which are probably reworked from previous deposits (Kellner et al. 2009). The holotype of *Oxalaia quilombensis* sp. nov. consisting of premaxillae (MN 6117-V)



strong tides that affect this deposit. An isolated portion of a left maxilla showing the same general features of spinosaurids is also referred to this species.

Despite being incomplete, this material shows the main characters of the theropod clade Spinosauridae and is a new addition to this bizarre and poorly known group of theropod dinosaurs.

ABBREVIATIONS

MNSM – Museo di Storia Naturale di Milano, Italy.

MN – Museu Nacional/Universidade Federal do Rio de Janeiro, Brazil.

MNHN SAM – Muséum National d’Histoire Naturelle, Paris, France.

SYSTEMATIC PALEONTOLOGY

THEROPODA Marsh 1881

SPINOSAUROIDEA Stromer 1915

sensu Sereno et al. 1998

SPINOSAURIDAE Stromer 1915

sensu Sereno et al. 1998

Oxalaia gen. nov.

Etymology: The generic name comes from Oxalá, the most respected masculine deity in the African pantheon, introduced in Brazil during slavery.

Type species: *Oxalaia quilombensis* sp. nov., type by monotypy.

Diagnosis: as for the type and only species.

Oxalaia quilombensis sp. nov.

Etymology: The generic name is derived from the Portuguese expression *quilombo*, the place where the *quilombola* (the descendants of former Brazilian slaves) live. The Cajual Island, where the specimens of this new taxon were collected, is one of these places.

Holotype: Fused premaxillae of a very large individual (ca. 12-14m, 5-7 tons), housed at the Museu Nacional/UFRJ under the collection number MN 6117-V (Figs. 1-3).

Referred material: Isolated and incomplete left maxilla


Island, in Maranhão State, Northeast Brazil. The rocks of this site belong to the Alcântara Formation, Itapecuru Group of the São Luís Basin whose age is regarded as Cenomanian (Late Cretaceous) (Góes and Rossetti 2001).

Diagnosis: Spinosaurid with the following combination of characters that distinguish it from other members of this clade (autapomorphies are marked with an asterisk): teeth unserrated (differing from *Cristatusaurus* and *Suchomimus*); maximum expansion of the distal end of the premaxillae between the 3rd and 4th alveoli*; anterior projection of the maxillae between the premaxillae in the palatal region very thin*; presence of two replacement teeth associated with the 3rd functional tooth*; diastema between the 5th and 6th premaxillary teeth present but shorter than in *Spinosaurus*; ventral portion of the premaxillae very sculptured*.

DESCRIPTION AND COMPARISONS

The preserved segment of the premaxilla (MN 6117-V) of *Oxalaia quilombensis* is about 201 mm long, 115 mm wide (maximum estimated width: 126 mm) and 103 mm high (Figs. 1-3). Based on comparisons with other spinosaurids (e.g., Dal Sasso et al. 2005), the total length of the skull is estimated around 1350 mm. This bone is broken on the region of the 7th alveolus and, despite the fact that a second portion of an upper jaw is here tentatively referred to this taxon (MN 6119-V), the information of the posterior extension of the upper jaw in this new species is limited. The left side was partially embedded in the matrix and is the better-preserved portion (Fig. 1b). Some of the bone surface was abraded and the ventral edges that form the lateral surface of the alveoli of the right side are broken in some parts. The anterior margin of the rostrum shows some large and deep pits interpreted as neurovascular foramina. A row of nutrient foramina just above the bases of the teeth on the left side is observed.

The premaxilla has seven alveoli, the last one incompletely preserved on both sides (Fig. 2). This is the same number as found in *Angaturama*, *Suchomimus*, the



Oxalaia quilombensis sp. nov.




Fig. 1 – *Oxalaia quilombensis* gen. et sp. nov. (MN 6117-V). **a** – right lateral view; **b** – left lateral view. Abbreviations: **alv** – alveoli; **nfo** – nutritive foramen; **pm** – premaxilla; **te** – tooth. Scale bar: 50 mm.

attributed to *Spinosaurus* cf. *S. aegyptiacus* by Dal Sasso et al. (2005), in which only six premaxillary teeth were found. There is a marked variation in the teeth size, with the first one (incompletely preserved) rather small, and the 2nd and 3rd alveoli being the largest. One marked diastema separating the 3rd from the 4th tooth is observed, which is present in all other spinosaurids, being smaller in *Suchomimus*. A second diastema between the 5th and the 6th alveoli is also observed in *Oxalaia*, similar to the Algerian spinosaurid (MNHN SAM 124, Taquet and Russell 1998). This diastema is absent in *Suchomimus* and *Cristatusaurus* and is much longer in *Spinosaurus* cf. *Spinosaurus aegyptiacus* described by

tween the 6th and 7th alveoli. This terminal expansion of *Oxalaia* is similar to that observed in *Suchomimus* and *Cristatusaurus* by having the lateral margins gradually thinner posteriorly, lacking the small constriction present between the 3rd and 4th alveoli in the Algerian spinosaurid (MNHN SAM 124) and *Spinosaurus* cf. *S. aegyptiacus* that in those taxa give the anterior portion of the rosetta a rather triangular shape (Dal Sasso et al. 2005: Fig. 1b, Taquet and Russell 1998: Fig. 1).

Ventrally, the maxillae extend anteriorly, forming two elongated processes that are encased laterally by the premaxillae. These rostral extensions of the maxillae, which border a triangular opening, are rather





Fig. 2 – *Oxalaia quilombensis* gen. et sp. nov. (MN 6117-V) in ventral view. **a** – holotype; **b** – drawing of the holotype. Abbreviations: **idp** – interdental plate; **m** – maxilla; **pm** – premaxilla; 1 to 7 – numbers of alveoli. Scale bar: 50 mm.

illae is well ornamented, differing from the smoother condition observed in other spinosaurids.

Except for the replacement teeth, no tooth shows the apical portion of the crown. Teeth fragments are present in the 3rd and 5th alveoli of the right side and in the 2nd and 3rd alveoli of the left side. All preserved portions lack serrations. Where observable, the teeth are not lateromedially compressed as in other theropods (e.g., Currie et al. 1990), but show an oval transverse section, a typical feature of spinosaurid teeth (e.g., Stromer 1915, Kellner and Mader 1997). In the 3rd alveolus of the right side, a tooth fragment is visible, showing a

alveoli of the left side and on the 2nd and 3rd of the right side. The more apical section of the crown of the replacement teeth is straight, slightly flattened labiolingually. The interdental plates are prominent especially between the 2nd and 3rd; and 3rd and 4th alveoli.

The second specimen referred to *Oxalaia quilombensis* gen. et sp. nov. is the fragment of a left maxilla (MN 6119-V), containing two complete alveoli and part of a 3rd one (Fig. 4). The specimen was found as surface float, a common condition of the fossils in this deposit (e.g., Kellner et al. 2009), and shows extensive abrasion



Oxalaia quilombensis sp. nov.




Fig. 3 – *Oxalaia quilombensis* gen. et sp. nov. (MN 6117-V). **a** – ventral view, showing the replacement teeth in the 3rd alveolus; **b** – dorsal view. Arrows indicate the larger functional and the smaller replacement teeth. Scale bar: 50 mm.

suggests that this specimen was likely forming part on the posterior end of the external naris. A row of nutritive foramina above the base of the teeth is also preserved on the lateral surface of this bone.

DISCUSSION

Overall spinosaurids are one of the less well-known groups of theropod dinosaurs. Introduced in the literature by Stromer (1915), all specimens of *Spinosaurus aegyptiacus* collected in the early Cenomanian Bahariya Formation of Egypt were destroyed during the Second World War (Taquet 1984). No upper jaw was present in the material originally described (Stromer 1915, Kellner and Campos 1996), and therefore no di-

(2005) have referred a partial skull found in the Cenomanian strata of Morocco to *Spinosaurus aegyptiacus*. Providing this assignment is correct, the Moroccan species has only 6 premaxillary teeth, whereas 7 are present in all other spinosaurids (including *Oxalaia quilombensis*) which might be a diagnostic feature for this taxon. Another plausible explanation for the reduced number of premaxillary teeth in *Spinosaurus aegyptiacus* (MSNM V4047) is ontogeny, based on the fact that some reptiles tend to lose one or more teeth as they grow older (e.g., recent crocodylomorphs, *Tyrannosaurus rex*). However, due to the lack of a considerable sample size which is paramount in determining individual and ontogenetic variations (e.g., 2010), the reduction of premaxillary teeth as a





Fig. 4 – *Oxalaia quilombensis* gen. et sp. nov. (MN 6119-V), fragment of the left maxilla. **a** – in lateral view; **b** – in medial view; **c** – in ventral view. Abbreviations: **dep** – depression; **nfo** – nutritive foramen; **rid** – ridge; **te** – tooth. Arrows indicate the channel that ends inside the medioposterior part of the alveoli. Scale bar: 50 mm.

the spacing of the alveoli, much larger in the latter, and the small constriction between the 3rd and 4th alveoli of the Moroccan taxon, giving the anterior portion of the rosetta a rather triangular shape in dorsal view (see description).

A second taxon, *Spinosaurus maroccanus* from supposed Albian deposits of Morocco, was erected by Russel (1996) based on one cervical vertebra whose centrum is quite elongated. This species was considered by Sereno et al. (1998) and Dal Sasso et al. (2005) as *nomen dubium*. Taquet and Russell (1998) referred an incomplete rostrum collected in Albian deposits of Algeria to *Spinosaurus maroccanus*, but since there is no material in common that would allow a detailed comparison, this allocation is questionable and we refer to MNHN SAM 124 as the Algerian spinosaurid. In any case, *Oxalaia* differs from the latter by the thinner an-

tion of the rosetta in the Algerian spinosaurid has a triangular shape in dorsal view.

Differences between *Oxalaia quilombensis* and the Baryonychinae (*sensu* Sereno et al. 1998) – a clade that includes *Baryonyx walkeri*, *Cristatusaurus lapparenti* and *Suchomimus tenerensis* – are the teeth with serrated carinae and the absence of a marked diastema between the 5th and 6th alveoli (Charig and Milner 1986, 1997, Taquet and Russell 1998, Sereno et al. 1998). Besides these, *Oxalaia quilombensis* bears two replacement teeth in the 3rd alveoli, an unusual feature that, to our knowledge, is here reported for the first time in theropods.

Two additional spinosaurid dinosaurs were described from Brazil, both from the Albian Romualdo Formation of the Santana Group: *Irritator challengeri* and *Angaturama limai*. Based on different portions of



Oxalaia quilombensis sp. nov.

comparisons of the new species with *Irritator* that lacks the anterior end of the rostrum (Sues et al. 2002) cannot be made. Regarding *Angaturama limai*, *Oxalaia quilombensis* differs from this taxon by the lack of a sagittal crest and by having a wider rostrum, among others. Other spinosaurid material from the Romualdo Formation, consisting of vertebrae (Bittencourt and Kellner 2004) cannot be compared with the new species described here.

Based on the morphology of the rostrum and the dentition, two main groups of Spinosauridae can be recognized (e.g., Kellner and Campos 1996, Sereno et al. 1998). The Baryonychinae share teeth with finely serrated carinae, posterior end of the terminal rosetta less constricted than in spinosaurins, and lack a marked diastema between the 5th and 6th alveoli. The monospecific *Cristatusaurus* and *Suchomimus* are very similar and came from the same deposit, raising the possibility that they are congeneric or even conspecific, in which case *Cristatusaurus lapparenti* has priority over *Suchomimus tenerensis*.

The second group, the Spinosaurinae, is united by the unserrated carinae, a very unusual feature within theropods. They also show a comparatively smaller first premaxillary pair of teeth and a larger spacing in the upper jaw (and, where observable, also in the lower jaw, Sereno et al. 1998). Within spinosaurins, *Spinosaurus* cf. *Spinosaurus aegyptiacus* and the Algerian spinosaurid (found in deposits of different age – Cenomanian and Albian, respectively) apparently are closely related in respect to *Oxalaia quilombensis*, which is based on the particular triangular shape of the rostral end. Also part of the spinosaurin clade is *Angaturama limai* that differs from the new species described here by having a sagittal crest and a comparatively more compressed rostrum (features also possibly present in *Irritator chalengeri*, see Machado and Kellner 2005). Morphologically, *Angaturama* (and likely *Irritator*) is very distinct from other spinosaurins and perhaps represents a different group of spinosaurid theropods.

From the same deposits of *Oxalaia quilombensis*, Medeiros (2006) reported the presence of hundreds of

verse section. Morphotype 2 also has the general shape of spinosaurin teeth (e.g., unserrated carinae, sublabiolingually compressed) but presents the tooth surface smooth. Although fragmentary, the dentition observed in *Oxalaia quilombensis*, compares well with that of morphotype 1 of Medeiros. The specimens of morphotype 1 apparently show a broad range of variations, particularly regarding compression, and represent either worn teeth of morphotype 1 or do indeed indicate a yet undescribed spinosaurin theropod in those deposits, as Medeiros (2006) supposed.

Several typical spinosaurid teeth have been reported in other stratigraphic units, some mistaken as belonging to crocodylomorphs (see Buffetaut 2010 for a recent review). In Brazil, Candeiro et al. (2004) reported some isolated teeth from the Bauru Basin to this group, and latter Candeiro et al. (2006: fig.12c) dismissed their occurrence but still labeled one tooth as Spinosaurid. Compared to published spinosaurid teeth (e.g., Sereno and Mader 1997, Sereno et al. 1998, Medeiros 2006) this material most likely belongs to crocodylomorphs. It has been pointed out before (Machado et al. 2008, Medeiros et al. 2009) reported one tooth from the Lisandro Formation (lower Turonian, Early Cretaceous) as possible referable to the Spinosauridae. However, despite the fluting and the oval transverse section, this is rather unusual for spinosaurids in being small and having coarsely serrated carinae, contrary to what is reported in the Baryonychinae (e.g., Sereno et al. 1998). Although any assessment is difficult without cranial material, the Lisandro specimen either belongs to a distinctive spinosaurid or to a bizarre crocodylomorph, the latter being favored here.

Comparisons between the premaxillae of *Oxalaia quilombensis* and *Spinosaurus* cf. *S. aegyptiacus* (MSNM V4047) suggest that the new Brazilian spinosaurid had a total length between 12 and 14 meters and a mass of 5 to 7 tons, making it the largest theropod recovered from the country so far.

CONCLUSIONS

Although mainly known from the premaxillae



its incompleteness, the material further shows a distinct morphology from other spinosaurids and comes from a distinct deposit, which is located in a different geographical area from other spinosaurid occurrences, criteria regarded as determinant for the recognition of a new species in the fossil record (see discussion in Kellner 2010).

Regarding the phylogenetic position, *Oxalaia quilombensis* is part of the Spinosaurinae, closely related to *Spinosaurus* cf. *Spinosaurus aegyptiacus* and the Algerian spinosaurid (Dal Sasso et al. 2005, Taquet and Russell 1998). It is further the largest theropod recovered from Brazil so far and demonstrates that this clade of bizarre theropods was present on the coastal area of the country from the Albian (e.g., *Angaturama*) to the early Cenomanian.

ACKNOWLEDGMENTS

Gareth Dyke (University College Dublin, Ireland) and Jonathas de Souza Bittencourt (USP, Ribeirão Preto) are thanked for comments on earlier drafts of the ms. We thank the Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq, grant number 307276/2009-0 to AWAK) and Fundação Carlos Chagas Filho de Amparo à Pesquisa do Estado do Rio de Janeiro (FAPERJ, grant number E-26/102.779/2008 to AWAK) that partially funded this project. The drawing of the type-specimen was done by André Pinheiro (Instituto de Geociências, UFRJ). We would also like to acknowledge João Nogueira and Leonardo Eddie (Rio de Janeiro) for support during the development of this project.

RESUMO

Um novo espinossaurídeo, *Oxalaia quilombensis* gen. et sp. nov., é descrito com base na parte anterior de um rostró e de um fragmento da maxila. Os espécimes são procedentes do depósito Laje do Coringa, formado durante o Cretáceo Superior (Cenomaniano) da Bacia de São Luís. Ao contrário de *Cristatusaurus* e *Suchomimus*, *Oxalaia quilombensis* não possui dentes serrilhados. A nova espécie difere de *Angaturama limai* pela ausência de crista sagital e por possuir a região do rostró mais expandida. Também difere de *Spinosaurus*

maxilares, uma feição até então desconhecida em terópodes. *Oxalaia quilombensis* está aparentemente mais proximamente relacionado aos espinossaurídeos da região norte da África do que às formas brasileiras e aumenta a diversidade deste grupo enigmático de dinossauros predadores no país.

Palavras-chave: Dinosauria, Spinosauridae, *Oxalaia quilombensis*, Cenomaniano, Brasil.

REFERENCES

- BITTENCOURT JS AND KELLNER AWA. 2004. On a sequence of sacrocaudal theropod dinosaur vertebrae from the Lower Cretaceous Santana Formation, northeastern Brazil. *Arq Mus Nac* 62: 309–320.
- BITTENCOURT JS AND KELLNER AWA. 2009. The anatomy and phylogenetic position of the Triassic dinosaur *Staurikosaurus pricei* Colbert, 1970. *Zootaxa* 2079: 1–54.
- BUFFETAUT E. 2010. Spinosaurids before Stromer: early finds of spinosaurid dinosaurs and their interpretation. In: MOODY R, BUFFETAUT E, NAISH D AND MARTILL D (Eds), *Dinosaurs and Other Extinct Saurians: A Historical Perspective*. Geol Soc, London, Spec Publ 343: 175–188.
- CANDEIRO CRA, ABRANCHES CT, ABRANTES EA, AVILLA LS, MARTINS VC, MOREIRA AL, TORRES SR AND BERGQVIST LP. 2004. Dinosaurs remains from western São Paulo state, Brazil (Bauru Basin, Adamantina Formation, Upper Cretaceous). *J South Am Earth Sci* 18: 1–10.
- CANDEIRO CRA, MARTINELLI AG, AVILLA LS AND RICH TH. 2006. Tetrapods from the Upper Cretaceous (Turonian-Maastrichtian) Bauru Group of Brazil: a reappraisal. *Cretaceous Res* 27: 923–946.
- CHARIG AJ AND MILNER AC. 1986. *Baryonyx*, a remarkable new theropod dinosaur. *Nature* 324: 359–361.
- CHARIG AJ AND MILNER AC. 1997. *Baryonyx walkeri*, a fish-eating dinosaur from the Wealden of Surrey, *Bull Nat Hist Mus, Geology Series* 53(1): 11–70.
- CORRÊA-MARTINS FJ. 1997. A Laje do Coringa: um boneco da Formação Itapecuru (Cretáceo Superior da Bacia de São Luís – MA). *An Acad Bras Cienc* 63(3): 436–437.
- CURRIE PJ, RIGBY JK JR AND SLOAN RE. 1990. Theropod teeth from the Judith River Formation of southern Alberta, Canada, p. 107–125. In: CARPENTER K AND CURRIE PJ (Eds), *Dinosaur Systematics: Perspectives and Approaches*. Cambridge University Press, Cambridge.



Oxalaia quilombensis sp. nov.

- GALTON PM. 2000. Are *Spondylosoma* and *Staurikosaurus* (Santa Maria Formation, Middle-Upper Triassic, Brazil) the oldest saurischian dinosaurs? *Paläont Zeit* 74: 393–423.
- GIBSON SA, THOMPSON RN, WESKA RK, DICKIN AP AND LEONARDOS OH. 1997. Late Cretaceous rift-related upwelling and melting of the Trindade starting mantle plume head beneath western Brazil. *Contr Mineral Petrol* 126: 303–314.
- GÓES AM AND ROSSETTI DF. 2001. Gênese da Bacia de São Luís-Grajaú, meio norte do Brasil. In: ROSSETTI DF, GÓES AM AND TRUCKENBRODT W (Eds), *O Cretáceo na Bacia de São Luís-Grajaú*. Belém, Museu Goeldi, p. 15–30.
- KELLNER AWA. 1999. Short note on a new dinosaur (Theropoda, Coelurosauria) from the Santana Formation (Romualdo Member, Albion), Northeastern Brazil. *Bol Mus Nacl, Geol* 49: 1–8.
- KELLNER AWA. 2010. Comments on Pteranodontidae with the description of two new species. *An Acad Bras Cienc* 82: 1063–1084.
- KELLNER AWA AND CAMPOS DA. 1996. First Early Cretaceous theropod dinosaur from Brazil with comments on Spinosauridae. *N Jahr Geol Palaeont Abh* 199(2): 151–166.
- KELLNER AWA AND CAMPOS DA. 1999. Vertebrate paleontology in Brazil – a review. *Episodes* 22: 238–251.
- KELLNER AWA AND CAMPOS DA. 2000. Brief review of dinosaur studies in Brazil. *An Acad Brasil Cienc* 72: 509–538.
- KELLNER AWA AND CAMPOS DA. 2002. On a theropod dinosaur (Abelisauria) from the continental Cretaceous of Brazil. *Arqui Mus Nac* 60: 163–170.
- KELLNER AWA AND MADER BJ. 1997. Archosaur teeth from the Cretaceous of Morocco. *J Paleont* 71: 525–527.
- KELLNER AWA, MAISEY JG AND CAMPOS DA. 1994. Fossil down feather from the Lower Cretaceous of Brazil. *Paleontology* 37: 489–492.
- KELLNER AWA, PINHEIRO AEP, AZEVEDO SAK, HENRIQUES DDR, CARVALHO LB AND OLIVEIRA G. 2009. A new crocodyliform from the Alcântara Formation (Cenomanian), Cajual Island, Brazil. *Zootaxa* 2030: 49–58.
- LANGER MC. 2004. Basal Saurischians. In: WEISHAMPEL DB, DODD JR R, OSMUNDSON H (Eds), *The Dinosauria*. Berkeley, University of California Press, 1–10.
- LANGER MC, EZCURRA MD, BITTENCOURT J AND NOVAS FE. 2009. The origin and early evolution of dinosaurs. *Biol Rev* 84: 1–56.
- LEONARDI G. 1994. Annotated atlas of South American pod footprints (Devonian to Holocene). *Compact Pesquisa e Recursos Minerais – CPRM, Brasília*, 1–100.
- MACHADO EB, CAMPOS DA AND KELLNER AWA. 2005. On a theropod scapula (Upper Cretaceous) from the rília Formation, Bauru Group, Brazil. *Paläont* 78: 308–313.
- MACHADO EB AND KELLNER AWA. 2005. Nota sobre a Spinosauridae (Theropoda, Dinosauria). *An In UFRJ* 28: 158–173.
- MARTILL DM, CRUICKSHANK ARI, FREY E, SMITH AND CLARKE M. 1996. A new crested maniraptor dinosaur from the Santana Formation (Lower Cretaceous of Brazil). *J Geol Soc London* 153: 5–8.
- MEDEIROS MA. 2006. Large theropod teeth from the cenomanian of northeastern Brazil and the occurrence of Spinosauridae. *Rev Brasil Paleont* 9: 333–338.
- MEDEIROS MA AND SCHULTZ CL. 2002. A fauna sauriana da Laje do Coringa, Cretáceo médio do Nordeste do Brasil. *Arq Mus Nac* 60: 155–162.
- NAISH D, MARTILL DM AND FREY E. 2004. Ecology, Systematics and Biogeographical Relationships of Dinosauria Including a New Theropod, from the Santana Formation (?Albian, Early Cretaceous) of Brazil. *Historical Biology* 16: 57–70.
- RUSSEL DA. 1996. Isolated dinosaur bones from the Cretaceous of the Tafilalet, Morocco. *Bull Musé National d’Histoire Naturelle* 4: 349–402.
- SALGADO L, CANUDO JI, GARRIDO AC, RUIZ-OMATE JI, GARCÍA RA, FUENTE MS, BARCO JL AND LATI R. 2009. Upper Cretaceous vertebrates from the fit teatro area, Río Negro, Patagonia, Argentina. *Cret Res* 30: 767–784.
- SERENO PC ET AL. 1998. A long snouted predatory dinosaur from Africa and the evolution of Spinosaurids. *Nature* 391: 1298–1302.
- STROMER E. 1915. Ergebnisse der Forschungsreisen von E. Stromer in den Wüsten Ägyptens. II. Wirbeltiere der Baharije-Stufe (unterstes Cenoman). 3. Das Skelett des Theropoden *Spinosaurus aegyptiacus* nov. gen. *Abhandlungen der Königlich Preussischen Akademie der Wissenschaften* 1: 1–100.



- SUES HD, FREY E, MARTILL DM AND SCOTT DM. 2002. *Irritator challengeri*, a spinosaurid (Dinosauria: Theropoda) from the Lower Cretaceous of Brazil. *J Vert Paleont* 22: 535–547.
- TAQUET P. 1984. Une curieuse spécialisation du crâne de certains Dinosaures carnivores du Crétacé: le museau long et étroit des Spinosauridés. *CR Acad Sci (Paris)* 299: 217–222.
- TAQUET P AND RUSSELL DA. 1998. New data on spinosaurid dinosaurs from the Early Cretaceous of the Sahara. *CR Acad Sci (Paris)* 327: 347–353.
- VAN TOMME MPA, MEDEIROS MA AND MACHADO EB. 2008. On an enigmatic theropod tibia from the Alcântara Formation (Late Cretaceous, Early Cenomanian) of Maranhão, NE Brasil, p. 199–200. In: *BOLETIM DE RESUMOS DO VI SIMPÓSIO BRASILEIRO DE PALEONTOLOGIA DE VERTEBRADOS. Paleontologia em Destaque, boletim especial*.
- WESKA RJ. 2006. Uma síntese do Cretáceo Superior mato-grossense. *Geociências* 25: 71–81.