



Anais da Academia Brasileira de Ciências

ISSN: 0001-3765

aabc@abc.org.br

Academia Brasileira de Ciências

Brasil

MATEUS, OCTÁVIO; JACOBS, LOUIS L.; SCHULP, ANNE S.; POLCYN, MICHAEL J.; TAVARES, TATIANA S.; BUTA NETO, ANDRÉ; MORAIS, MARIA LUÍSA; ANTUNES, MIGUEL T.

Angolatitan adamastor, a new sauropod dinosaur and the first record from Angola
Anais da Academia Brasileira de Ciências, vol. 83, núm. 1, marzo, 2011, pp. 221-233

Academia Brasileira de Ciências
Rio de Janeiro, Brasil

Available in: <http://www.redalyc.org/articulo.oa?id=32717681011>


- How to cite
- Complete issue
- More information about this article
- Journal's homepage in redalyc.org

redalyc.org

Scientific Information System

Network of Scientific Journals from Latin America, the Caribbean, Spain and Portugal

Non-profit academic project, developed under the open access initiative



Anais da Academia Brasileira de Ciências (2011) 83(1): 221-233
(Annals of the Brazilian Academy of Sciences)
Printed version ISSN 0001-3765 / Online version ISSN 1678-2690
www.scielo.br/aabc

***Angolatitan adamastor*, a new sauropod dinosaur and the first record from Angola**

OCTÁVIO MATEUS^{1,2}, LOUIS L. JACOBS³, ANNE S. SCHULP⁴, MICHAEL J. POLCYN³,
TATIANA S. TAVARES⁵, ANDRÉ BUTA NETO⁵, MARIA LUÍSA MORAIS⁵ and MIGUEL T. ANTUNES⁶

¹CICEGe, Faculdade de Ciências e Tecnologia, FCT, Universidade Nova de Lisboa, 2829-516 Caparica, Portugal

²Museu da Lourinhã, Rua João Luis de Moura, 2530-157 Lourinhã, Portugal

³Huffington Department of Earth Sciences, Southern Methodist University, Dallas, TX, 75275, USA

⁴Natuurhistorisch Museum Maastricht, de Bosquetplein 6-7, NL6211 KJ Maastricht, The Netherlands

⁵Geology Department, Universidade Agostinho Neto, Av. 4 Fevereiro, Luanda, Angola

⁶Academia de Ciências de Lisboa, Rua da Academia das Ciências, 19, 1242-122 Lisboa, Portugal

Manuscript received on October 19, 2009; accepted for publication on June 21, 2010

ABSTRACT

A forelimb of a new sauropod dinosaur (*Angolatitan adamastor* n. gen. et sp.) from the Late Turonian of Iembe (Bengo Province) represents the first dinosaur discovery in Angola, and is one of the few occurrences of sauropod dinosaurs in sub-Saharan Africa collected with good chronological controls. The marginal marine sediments yielding the specimen are reported to be late Turonian in age and, thus it represents a non-titanosaurian sauropod in sub-Saharan Africa at a time taken to be dominated by titanosaurian forms. Moreover, *Angolatitan adamastor* is the only basalmost Somphospondyli known in the Late Cretaceous which implies the existence of relict forms in Africa.


Key words: Angola, Cretaceous, Turonian, Dinosaurs, Sauropoda, *Angolatitan*.

INTRODUCTION

Although very rich in Cretaceous vertebrates, Angola has remained unexplored in recent decades. Beginning in 2005, the field work conducted by the PaleoAngola Project has significantly enhanced the palaeontological record of the Cretaceous of Angola (Jacobs et al. 2006, 2009a, b, Polcyn et al. 2009, Mateus et al. 2008, Schulp et al. 2008). The PaleoAngola Project mounted the first systematic paleontological expeditions that have been conducted since the early 1960s, when Miguel Telles Antunes performed an extensive and systematic survey on the Angolan fossil vertebrate fauna the most relevant work published in 1964 (Antunes 1964). In May 2005 we (O.M. and L.J.) prospected the coastal area near Iembe which resulted in the discovery, in May 25th (by

OM), of the new taxon reported here. Follow-up excavations were performed in May and August 2005. To our knowledge, it was the only systematic prospecting for fossil vertebrates in this area since 1960 by Miguel Antunes (in September) and M. Mascarenhas Neto.

While the lush vegetation in most inland areas precludes fossil prospecting, the exposures along the Atlantic coastline in Ambriz municipality allow systematic prospecting. The material collected by the PaleoAngola Project is largely housed at Museu Geológico da Universidade Agostinho Neto (MGU) in Luanda, while some of the collected mosasaur, dinosaur, pterosaur, and turtle specimens are temporarily housed at Museu da Lourinhã (ML), Portugal. The Southern Methodist University (SMU) in Dallas



GEOLOGY AND ASSOCIATED FAUNA

The term ‘Tadi beds’ was first coined in the 1960 unpublished study of G. Brognon, G. Verrier, and R Thiers “Géologie du Bassin du Cuanza et du Bas-Congo” from the “Missão de Pesquisas de Petróleo Petrofina”. It was accepted and used by Antunes (1964) and Antunes and Cappetta (2002). The original study considered the ‘Tadi Beds’ to represent the base of the ‘Senonian’ (= Coniacian), but the fish fauna led Antunes (1961, 1964: 56) to consider the Tadi Beds to be of late Turonian age. In a revision of the Cretaceous selachians of Angola, Antunes and Cappetta (2002: 95) reconfirm the late Turonian age of these beds, and place them in the ‘Itombe Formation’ (see also Jacobs et al. 2006). The Tadi Beds are about 50 m thick (Antunes 1964: 56). G. Brognon et al. (cited in Antunes 1964: 56) provide a list of the invertebrate fauna, including the ammonite genera *Hemitissotia*, *Romaniceras* and *Pachydiscus*, and echinoderms. The fish and shark fauna (Antunes and Cappetta 2002: 95) includes *Hexanchus* cf. *microdon*, *Ptychodus mortoni*, *P.* cf. *paucisulcatus*, *P. whipplei*, *Scapanorhynchus raphiodon*, *S.* cf. *lewesii*, *Cretodus semiplicatus*, *Cretodus* sp., *Cretoxyrhina mantelli*, *Paranomotodon angustidens*, *Squalicorax falcatus*, *Squalicorax* sp., *Onchosaurus pharaoh*, and *Enchodus* cf. *elegans*.

The tetrapod fauna from the Iembe locality (Antunes 1964, Jacobs et al. 2006; and new unpublished data) includes an eucryptodiran turtle *Angolachelys mbaxi* (Mateus et al. 2009), the mosasaurs *Angolasaurus bocagei* Antunes 1964 (see also Polcyn et al. 2009) and *Tylosaurus iembeensis* (Antunes 1964, Lingham-Soliar 1992), Plesiosauria indet., and the sauropod material described here. The discovery here reported is the first dinosaur and the first Cretaceous terrestrial animal from Angola.

SYSTEMATICS PALEONTOLOGY

Dinosauria Owen, 1842

Sauropoda Marsh, 1878

Neosauropoda Bonaparte, 1986




Fig. 1 – Map of Angola, showing the Tadi site (star; 08°S/13°E) and the distribution of Cretaceous strata (grey) in Angola.

ETYMOLOGY


Angolatitan means Angolan giant. *Adamastor* was a mythological sea giant from the South Atlantic feared by the Portuguese sailors.

GENERIC AND SPECIFIC DIAGNOSIS

Titanosauriform sauropod with: posteroventral eminence in scapula in the proximal one-quarter, anterior to the acromial ridge; acute medioproximal margin of humerus; and rectangular lateral corner of proximal humerus; proximal outline of the ulna with anteromedial ridge (process); posterior facet of the distal epiphysis of metacarpal I with two small splint-like projections (Figs. 3 and 5).

TYPE LOCALITY AND HORIZON

Tadi, near Iembe, Tábi community, municipality of Ambriz, Bengo Province, Angola. The precise coordinates of collecting sites are on file at the participating institu-



NEW SAUROPOD DINOSAUR FROM ANGOLA

		Age	M.a.	Unit	Lithology	Paleo-environment	Max. thickness	Formation
CRETACEOUS	LATE	MAAS	70		Shale with interbedded calcite-cemented sandstone and siltstone	MARINE	900	RIO DANDE
		CMP	80		Limestone, calcareous sandstone interbedded with silty shale			TEBA
		SAN CON			Shale		100	N'GOLOME
		TUR	90		Limestone, shale, sandstone, siltstone, silty shale		600	ITOMBE
		CEN			Silty shale Banded limestone		400	CABO LEDO
	EARLY	ALBIAN	-100		Limestones		250	QUISSONDE
					Calcareous limestone		100	CATUMBELA
					Dolomites		1000	TUENZA
		APTIAN	-110		Dolomites and salt			
					Dolomite, limestones		150	BINGA
					Calcareous limestone, dolomite, and anhydrite		1400	QUANANGA
		BRM	-120		Salt			MASSIVE SALT
					Dolomite and argillite			UPPER (Grey) CUVO
		NEOCOMIAN	-130		Red argillaceous sandstone		2000	LOWER (Red) CUVO
					Sandstone and shale		2000	MACULUNGO
			-140					

Compiled by Octávio Mateus based on:
Coward et al. 1999, <http://books.google.pt/books?id=YVooEivHwIMC&printsec=frontcover>
Brownfield and Charpentier 2006, <http://www.usgs.gov/bul/2207/8/>

Fig. 2 – Summary of the geology of the Cretaceous of Kwanza Basin (Angola) based on Brownfield and Charpentier (2006) and Coward et al. (1999).

MATERIAL

The material includes a partial right forelimb including the scapula, humerus, ulna, radius and metacarpals I, III, and IV (Table I). All material described here is curated at Museu de Geologia da Universidade Agostinho Neto – PaleoAngola Project, Luanda, Angola (MGUAN-PA), and temporarily registered under field number MGUAN-PA-003 (replicas will be housed at Museu da Lourinhã, Portugal, and at Southern Methodist University, USA).

DESCRIPTION

The **scapula** is missing its posterior end and part of

However, this could be a case of taphonomical distortion. The proximal end is incomplete, but does not appear very expanded. The scapular blade is 16.5 cm, with a very expanded ventral supraglenoid expansion just above it, and on the maximum dorsal expansion. The ventral expansion of the glenoid expansion is 65 cm in its maximum anterior width, therefore representing nearly four times the minimum width of the blade. For the description, the orientation of the scapula adopted here is with the scapular blade in a vertical dorsal position as in Brownfield (1956).

The blade is rectangular in cross section, with a slightly expanded anterior margin. The anterior margin (acromion)





Fig. 3 – *Angolatitan adamastor* n. gen. et sp., A) right scapula in medial view; B) right humerus in proximal(a), anterior (b), distal (c), medial (d), and posterior (e) views. Autapomorphic features: 1, posteroventral eminence in scapula in the proximal one-quarter, anterior to the acromial ridge; 2, acute medioproximal margin of humerus; and 3, rectangular lateral corner of proximal humerus.


anteriorly, forming a gentle curve in its anterodorsal corner due to the absence of the dorsal hook.

The medial surface is gently concave in all its extension, except in a small (ca. 6 cm) shallow bump placed at the anterior part at the base of the scapular blade, and in the bump in the posteromedial margin of the scapula about 20 cm above the glenoid.

The contact surface between the coracoid and glenoid is perpendicular with the blade in a posterior or anterior perspective, but the contact with the coracoid runs slightly dorsally, forming a 20° angle with the horizontal (if the scapula is arranged upwards). The

ally glenoid is apomorphic of *Somphospondyli* according to Wilson 2002 (character 153).

The ventroposterior margin of the scapula, adjacent to the glenoid, projects posteriorly and has a small (ca. 9 cm) vertical ridge that borders a groove somewhat similar to *Cetiosaurus oxoniensis*. A posteroventral bump in the scapula in the proximal one-quarter is shared/convergent with *Dystrophaeus* and *Supersaurus*, *Ultrasaurus mcintoshi*, *Lourinhasaurus alenquerensis*, and *Camarasaurus supremus*, but, unlike other sauropods, it is anterior to the acromial ridge. The glenoid foramen is not visible.



NEW SAUROPOD DINOSAUR FROM ANGOLA




Fig. 4 – *Angolatitan adamastor* n. gen. et sp., A) right ulna in medial (a), anterior (b), lateral (c), posterior (d), proximal (e) and distal (f) views and (B) right radius in proximal (a), distal (b), anterior (c), and posterior (d) views.

gracile with a straight shaft. The proximal end is L-shaped due to the well-defined deltopectoral crest. The humeral head is well developed and more projected anteriorly than the lateral corner, which gives an asymmetrical outline in anterior view. The lateral corner forms a rectangular corner (rather than a round corner more commonly seen in sauropods) and a straight line to the most proximal part of the humerus. The medioproximal corner of the humerus is very pointy due to a 4 cm

margin. The tip of the deltopectoral crest projects anteromedially, invading the anterior side of the humerus. A medially projected deltopectoral crest is present in Titanosauriformes more derived than *Brachiosaurus*. The bump placed lateromedially to the tip of the deltopectoral crest gently projects laterally, making a wide margin for muscle attachment. Except for this bump, the lateral margin is totally straight. A hemispherical humeral head projects well posteriorly. The diaphyseal cross-





Fig. 5 – *Angolatitan adamastor* n. gen. et sp., right metacarpals I (A), III (B) and IV (C), in proximal (a), anterior (b), medial (c), palmar (d), lateral (e), and distal (f) views. 4, posterior facet of the distal epiphysis of metacarpal I has two small splint-like projections (autapomorphy).

ysis, being well separated by posterior and anterior intercondylar grooves. The distal articular facet is slightly convex, nearly planar, without condylar expression. At the anterior face, the distal condyles are directed antero-medially without significant dorsal projection. There is a vertical groove between the lateral condyle and the lateral margin, that creates a notch aspect in distal view. In the posterior face, the lateral condyle is round, while the medial condyle forms an acute vertical edge. The posterior intercondylar groove is wide and deep.

The right **ulna** is complete, except for a small part of the lateral face of the diaphysis. The general

width at its distal expansion. In proximal view, the ulna bears a tri-radiate outline, with prominent anterolateral and anteromedial processes, and a less-developed posterior process. The proximal outline shows an expanded anteromedial ridge (process) constricted in the base. The anteromedial ridge is nearly twice the lateral one. The olecranon process is poorly developed and does not projects significantly more proximally than the anteromedial one. The shaft is moderately straight and less robust than in most titanosaurs. The distal epiphysis is sub-round to sub-triangular, with a slight transverse expansion.



NEW SAUROPOD DINOSAUR FROM ANGOLA

TABLE I
Measurements of *Angolatitan adamastor* (in cm).

	Length	Minimal diaphyseal circumference	Maximal proximal expansion	Maximal distal expansion
Scapula	95	–	61	–
Humerus	110	43.2	36	29
Ulna	69	27*	23	14.5
Radius	63	22.5	16.5	13
Metacarpal I	30.0	–	10.5	8.5*
Metacarpal III	30.9	13*	9	8.5*
Metacarpal IV	28.5	12	9*	–

*Estimated.


epiphyseal expansions. The maximum proximal width equals about 25% of the full length of the radius. The proximal end is sub-round, while the distal end is flattened and elongated with an anteroposterior expansion. The medial edge of the distal epiphysis is acute, but the lateral is round. The posterior face of the distal epiphysis bears a groove at the ulnar articulation procuding two shallow posterior condyles. The medial part projects more distally forming a bevelled angle with the main shaft.

Three **metacarpals** (I, III and IV) were present near the distal end of the ulna. With the exception of the bone surfaces exposed to weathering prior to discovery, the bones do not show signs of crushing or deformation. The metacarpal position was identified as I, III, and IV using Janensch (1961), Apesteguía (2005) and Bonnan and Wedel (2004). The metacarpals are elongated, much more than in titanosaurs. All metacarpals are nearly of the same size, but III is the largest and IV is the shortest preserved element. The distal end bears subtle but separated distal condyles which is a plesiomorphic condition because the presence of divided condyles is normally associated with the presence of phalanges, lost in titanosaurs (Wilson 2002, Upchurch et al. 2004). The metacarpals bound together by the extensive diaphyseal articulation to form a tubular arrangement, U-shaped in proximal view. The length ratio between metacarpal III and the radius is 0.48. A ratio higher than 0.45 is used to be considered a character

medial corner more rounded than the lateral. The carpal presents a crest along the whole lateral face with a sinusoidal area at midshaft. The distal end is transversely broad, with two well-defined condyles, less visible than in non-macronarian sauropods) (Fig. 1) by distal and posterior grooves, the lateral condyle is smaller than the medial. The posterior facet of the epiphysis has two small splint-like projections: one above the intercondylar groove and the other near the lateral condyle, which is autapomorphic.

The right **metacarpal III** is complete, with the exception of a portion of the lateral distal condyle. It is subtriangular in proximal view. Ligament attachments on the palmar surface occupy almost the entire proximal half of the metacarpal – which also represents the area of contact with metacarpal IV. The medial face bears an acute crest along the diaphysis that is divided into a proximal third, and rims a large facet facing anteroposteriorly. The distal end shows defined condyles, being the lateral one the largest.

The **metacarpal IV** is robust. In the proximal view it is triangular, with the anterior margin being the widest. The proximal surface is flat, but the distal corner which is projected medially deflects distally forming a 45° angle with the proximal surface. The distal end is transversely broad and presents two condyles divided by a medial vertical intercondylar groove. There is no distal intercondylar groove. This bone resembles the metacarpal IV of cf. *Laplatasaurus*



There are no signs of other carpals, metacarpals or phalanges, but the lateral flattening on metacarpal IV suggests the possible presence of metacarpal V, and the distal condyles suggest the presence of phalanges.

COMPARISONS AND PHYLOGENY

The olecranon process of the ulna is more proximally projected than in Jurassic sauropods *Lourinhasaurus*, *Lusotitan*, *Apatosaurus*, *Camarasaurus*, and *Diplodocus*, but less than in most Titanosauria, when compared with *Saltasaurus loricatus* “*Titanosaurus*” *colberti*, *Tornieria robusta* (Janensch 1961), *Epachthosaurus sciuttoi* (Martínez et al. 2004), and *Malawisaurus dixeyi* (Gomani 2005), and somehow with the same extension as in *Rapetosaurus krausei* (Curry Rogers 2009). Ulna robustness (proximal width/length) is 0.33 in *Angolatitan adamastor*, which is lower than 0.35 in *Lusotitan atalaiensis*, 0.35 in aff. *Turiasaurus* ML368, 0.36 in *Malawisaurus dixeyi* (Gomani 2005: Table 7), 0.34 in *Rapetosaurus krausei* (Curry-Rogers 2009: Table 3), and strikingly lower than 0.41–0.43 in *Epachthosaurus sciuttoi* (Martínez et al. 2004: Table 2). The data for establishing the robustness is unreported or unknown in most titanosaurs (e.g. *Limaysaurus tessonei*, *Rinconosaurus cadamirus*, *Muyelensaurus pecheni*, and *Saltasaurus loricatus*), for which the ulnae is clearly more robust. Wilhite (2003) provides measures for many Late Jurassic ulnae, which, in average, gives the following means for robustness: 0.40 for *Apatosaurus*, 0.23 for *Barosaurus*, 0.39 for *Brachiosaurus*, 0.38 for *Camarasaurus*, and 0.33 for *Diplodocus*.

In *Angolatitan adamastor*, the metacarpals do not show three titanosaurian characters considered by Apesteguía (2005): bowed metacarpal I, the flat distal metacarpal trochlea (although this is difficult to quantify), and reduced intermetacarpal contact that is 1/6 of the metacarpal length.


The data matrix was based in Wilson (2002), but the integration of characters in Upchurch et al. (2004) for the forelimb (*Angolatitan* coding using Wilson 2002, characters 150 to 200: 1 0000????10 0101110010 1???-

using WinClada ver 1.0 (www.cladistics.com) using a heuristic search with 100 replicates. *Angolatitan adamastor* n.gen. et sp. was placed as a basal titanosauriform Somphospondyli more derived than *Brachiosaurus*, but less derived than *Euhelopus* (see Fig. 6). The metacarpals are slender (close to *Camarasaurus* proportions and alike titanosaurs as *Opisthocoeleicaudia*, when compared in the plot provided by Bonnan and Wedel 2004). The robustness (ratio minimum diaphyseal perimeter/length) is 0.42, which is lower than the range of 0.5–0.6 in most titanosaurs (Apesteguía 2005: Table 15.2). The size of metacarpals is equivalent (being of variable length in Titanosauridae, see Apesteguía 2005), and the distal end is round, rather than flat as in Titanosauridae.

The *Angolatitan adamastor* is supported as Somphospondyli by the following unambiguous synapomorphies: humeral square proximolateral corner (character 159 in Wilson 2002), deltopectoral crest medially expanded across the cranial face of the humerus (ch. 220 in Upchurch et al. 2004), and distalmost part of the caudal surface of the humerus deeply concave between the lateral and medial prominent vertical ridges (ch. 221 in Upchurch et al. 2004). However, the scapular glenoid surface faces cranioventrally (ch. 203 in Upchurch et al. 2004) unlike *Euhelopus* and other derived Somphospondyli. The Angolan sauropod does not bear titanosaurian synapomorphies such as the ulna with prominent olecranon process (Wilson 2002: character 167) or extremely robust ulna and radius (Upchurch et al. 2004: 309).

GEOGRAPHY AND CHRONOLOGY

Although sauropod dinosaurs have been reported from elsewhere in Africa, (e.g. Algeria, Cameroon, Egypt, Kenya, Malawi, Mali, Morocco, Niger, South Africa, Sudan, Tunisia, Zambia, Zimbabwe, and Madagascar), no others are known from the Late Turonian. The only described Late Cretaceous sauropods from Africa are the lithostrotian titanosaurs *Aegyptosaurus baharijensis* from Egypt and Niger (Stromer 1932: early Cenoman-



NEW SAUROPOD DINOSAUR FROM ANGOLA




Fig. 6 – Phylogenetic relationships of *Angolatitan adamastor* and other sauropods. See text for explanation.

rus krausei and “*Titanosaurus*” *madagascariensis* from Madagascar (Campanian; Curry Rogers and Forster 2001, Weishampel et al. 2004). Globally, the circa-Turonian sauropods were found only in Argentina, Brazil, and China (Table II), and comprise mostly titanosaurs or diplodocoids. Thus, *A. adamastor* fills a 15My gap in the African sauropod record, and an even wider gap in the global record of non-titanosaur sauropods.

Non-titanosaur sauropods, which were dominant during the Jurassic, were being replaced by titanosaurs and derived diplodocoids, becoming rarer in the Late Cretaceous when most identifiable African sauropods were titanosaurs. The reasons for the late Cretaceous dominance of titanosaurs over other sauropods are not fully understood, but it seems to be a general pattern in South America, in Madagascar, and probably in Africa

chronology and even taxonomy of these sauropods are not well understood for the Late Cretaceous of Africa. Diplodocoids were the most successful non-titanosaur sauropods in the Cretaceous, but the presence of possible brachiosaurids (Rauhut and Werner 1997) raises questions regarding the paleogeography of sauropods.

The age of the sediments yielding the specimen is Upper Turonian (~90Ma) based on the fish and plant teeth (Antunes and Cappetta 2002) and, thus, is the best-dated Late Cretaceous sauropod specimen from sub-Saharan Africa. Further work in this area, based on ammonites collected with the specimen, will almost certainly provide refined chronology and paleogeography of the Late Cretaceous dinosaur fauna of Africa.

During the Late Turonian, the coast of Iembe (located at 8°S latitude) was at about 24°S latitude due to





TABLE II
Circa-Turonian sauropod species.

Taxon	Country	Early_stage	Late_stage	Reference	Systematics	
<i>Andesaurus delgadoi</i>	Argentina	Albian	Turonian	Calvo and Bonaparte 1991, Calvo 1999	Titanosauria	Humerus
<i>Argentinosaurus huinculensis</i>	Argentina	Albian	Turonian	Bonaparte and Coria 1993	Titanosauria	Unknown
<i>Argyrosaurus superbus</i>	Argentina	Cenomanian	Turonian	Huene 1929	Titanosauria	Complete forelimb
<i>Borealosaurus wimani</i>	China	Cenomanian	Turonian	You et al. 2004	?Opisthocoelicaudiinae	Humerus
<i>Cathartesaura anaerobica</i>	Argentina	Cenomanian	Coniacian	Gallina and Apesteguía 2005	Rebbachisauridae	Scapula and femur
<i>Dongyangosaurus sinensis</i>	China	Cenomanian	Santonian	Lü et al. 2008	Titanosauria	Unknown
<i>Epachthosaurus sciutoi</i>	Argentina	Late Cenomanian	Early Turonian	Martínez et al. 2004	Titanosauria	Complete forelimb
<i>Futalognkosaurus dukei</i>	Argentina	Late Turonian	Early Coniacian	Calvo et al. 2007a	Titanosauria	Unknown
<i>Gondwanatitan faustoi</i>	Brazil	Turonian	Santonian	Kellner and Azevedo 1999	Lithostrotia	Humerus
<i>Huabeisaurus allocotus</i>	China	Cenomanian	Maastrichtian	Pang and Cheng 2000	Titanosauria	Scapula, coracoid, humerus, radius
<i>Huanghetitan ruyangensis</i>	China	Cenomanian	Santonian	Lü et al. 2007	Titanosauriformes	Unknown
<i>Limaysaurus tessonei</i>	Argentina	Albian	Turonian	Calvo 1999, Salgado et al. 2004	Rebbachisauridae	Forelimb
<i>Muyelensaurus pecheni</i>	Argentina	Late Turonian	Coniacian	Calvo et al. 2007b	Titanosauria	Forelimb
<i>Qingxiusaurus youjiangensis</i>	China	Cenomanian	Maastrichtian	Mo et al. 2008	Titanosauria	Sternal plates, humerus
<i>Rinconsaurus caudamirus</i>	Argentina	Late Turonian	Coniacian	Calvo and González Riga 2003	Titanosauria	Forelimb
<i>Sonidosaurus saihangaobiensis</i>	China	Turonian	Maastrichtian	Xu et al. 2006	Titanosauria	Unknown
<i>Sonidosaurus saihangaobiensis</i>	China	Turonian	Maastrichtian	Xu et al. 2006	Titanosauria	Unknown

15° and 30° mainly because of the atmospheric air convection Hadley Cells, which has remained stable since the Cretaceous (Jacobs et al. 2009a). Therefore, Iembe was an arid zone during the Turonian, which is further supported by the sediments at Iembe that show no signs of terrestrial productivity. *A. adamastor* was probably a sauropod adapted to arid/desert-like environment in a situation similar to the desert elephants today. Rubilar-Rogers (2006) reported a Cretaceous titanosaur sauro-

ACKNOWLEDGMENTS

Many thanks to Eduardo Morais and Agostinho Vaz (Universidade Agostinho Neto), Manuel João Fonseca and Margarida Ventura (Universidade Privada de Angola), Nancy Stevens, Luís Rocha, Rosário Sarzedas, and Pedro Serrano for their support in the field. The field expeditions were funded by the Petroleum Research Institute. Institute for the Study of Earth and



NEW SAUROPOD DINOSAUR FROM ANGOLA

and Phil Mannion for the comments in the phylogeny and anatomy. Preparation was conducted by O.M., A.S., Ricardo Araújo and C. Tomás. To João Marinheiro who edited some pictures.

RESUMO


Um membro anterior de um novo dinossauro saurópode (*Angolatitan adamastor* n. gen. et sp.) do Turoniano Superior de Iembe (Bengo) representa a descoberta do primeiro dinossauro em Angola e é uma das poucas ocorrências de dinossauros saurópodes na África Subsaariana recolhidos com bons controlos cronológicos. Os sedimentos marinhos marginais de onde provém o espécime estão datados do Turoniano superior e, portanto, trata-se de um saurópode não-titanossauro na África subsaariana onde predominavam titanossauros. Além disso, *Angolatitan adamastor* é o único Somphospondyli basal conhecido no Cretáceo Superior, o que implica a sobrevivência de formas-reliquia na África.

Palavras-chave: Angola, Cretáceo, Turoniano, dinossauros, sauropoda, *Angolatitan*.


REFERENCES

- ANTUNES MT. 1961. Sur la Faune de vertébrés du Cretacé de Iembe (Angola). CR Acad Sciences Paris 253: 513–514.
- ANTUNES MT. 1964. O Neocretácico e o Cenozóico do litoral de Angola. Junta de Investigações do Ultramar, 255 p.
- ANTUNES MT AND CAPPETTA H. 2002. Sélaciens du Crétacé (Albien-Maastrichtien) d’Angola. Palaeontographica 12: 85–146.
- APESTEGUÍA S. 2005. Evolution of the titanosaur metacarpus. *Thunder-Lizards, The Sauropodomorph Dinosaur*. Indiana University Press, p. 321–345.
- BONAPARTE JF AND CORIA RA. 1993. Un nuevo y gigantesco sauropodo titanosaurio de la Formacion Rio Limay (Albiano-Cenomaniano) de la Provincia de Neuquén, Argentina. Ameghiniana 30(3): 271–282.
- BONNAN MF AND WEDEL MJ. 2004. First occurrence of *Brachiosaurus* (Dinosauria: Sauropoda) from the Upper Jurassic Morrison Formation of Oklahoma. *PaleoBios* 24(2): 13–21.

- CALVO JO. 1999. Dinosaurs and other vertebrates from the Lake Ezequiel Ramos Mexía area, Neuquén – Patagonia, Argentina in Proceedings of the Second Gondwanan Dinosaur Symposium, National Science Museum, Tokyo, Graphs #15, In: TOMIDA Y, RICH TH AND VIGNATI P, RICH P (Eds), Tokyo: x + 296 p. P. 13–46.
- CALVO JO AND BONAPARTE JF. 1991. *Andesaurus* gen. et sp. nov. (Saurischia-Sauropoda), Dinosauria, Titanosauridae de la Formacion Rio Limay (Albiano-Cenomaniano), Neuquén, Argentina. Ameghiniana 28(4): 303–310.
- CALVO JO AND GONZÁLEZ RIGA BJ. 2003. *Rincosaurus caudamirus* gen. et sp. nov., a new titanosaurid dinosaur (Saurischia, Sauropoda) from the Late Cretaceous of Neuquén, Argentina. Rev Geol Chile 30(2): 3–23. doi:10.1007/s00716-02082003000200011. http://www.scielo.org/scielo.php?script=sci_arttext&pid=S0716-02082003000200011&lng=en&nrm=iso&tlang=en
- CALVO JO, GONZÁLEZ-RIGA BJ AND PORFIRI JD. 2004. A new titanosaur sauropod from the Late Cretaceous of Neuquén, Patagonia, Argentina. Arq Mus Nac 5(4): 504.
- CALVO JO, PORFIRI JD, GONZÁLEZ-RIGA BJ AND MANNION PHILIP A. 2007a. A new Cretaceous terrestrial sauropod from Gondwana with the description of a new titanosaurid dinosaur. An Acad Bras Cienc 79: 529–541.
- COWARD MP, PURDY EG, RIES AC AND SMITH D. 2007. The distribution of petroleum reserves in basins along the South Atlantic margins. Geol Soc, London, Sp 153(1): 101–131. (Available at <http://www.google.com/books?hl=pt-PT&lr=&id=YVooEivHwIMC>).
- CURRY ROGERS K. 2009. The postcranial osteology of *Rapetosaurus krausei* (Sauropoda: Titanosauria) from the Late Cretaceous of Madagascar. J Vert Paleontol 29: 1046–1086.
- CURRY ROGERS K AND FORSTER CA. 2001. The titanosaur titans: a new sauropod from Madagascar. Nature 412: 530–534.
- GALLINA PA AND APESTEGUÍA S. 2005. *Cathartes anaerobica* gen. et sp. nov., a new rebbachisaurid dinosaur (Saurischia, Sauropoda) from the Huincul Formation (Late Cretaceous), Rio Negro, Argentina. Rev Mus Arq Cienc Nat 7: 153–166.
- GOMANI EM. 2005. Sauropod dinosaurs from the Late Cretaceous of Malawi. Paleontologica Electronica 2(1): 1–10.



- JACOBS LL, MATEUS O, POLCYN MJ, SCHULP AS, ANTUNES MT, MORAIS ML AND TAVARES T. 2006. The occurrence and geological setting of Cretaceous dinosaurs, mosasaurs, plesiosaurs, and turtles from Angola. *J Paleo Soc Korea* 22: 91–110.
- JACOBS LL, MATEUS O, POLCYN MJ, SCHULP AS, SCOTSE CR, GOSWAMI A, FERGUSON KM, ROBINS JA, VINEYARD DP AND NETO AB. 2009a. Cretaceous paleogeography, paleoclimatology, and amniote biogeography of the low and mid-latitude South Atlantic Ocean. *B Soc Geol Fr* 180(4): 333.
- JACOBS LL, POLCYN MJ, MATEUS O, SCHULP AS AND NETO AB. 2009b. The Cretaceous Skeleton Coast of Angola. *J Vert Paleont* 29(Suppl 3): 121A.
- JANENSCH W. 1961. Die gliedmaszen und gliedmaszengürtel der Sauropoden der Tendaguru-schichten. *Palaeontographica* 3(Suppl 7): 177–235.
- KELLNER AWA AND AZEVEDO SAK. 1999. A new sauropod dinosaur (Titanosauria) from the Late Cretaceous of Brazil. In: TOMIDA Y, RICH TH AND RICH PV, *Proceedings of the Second Gondwanan Dinosaur Symposium*. *Natl Sci Mus Monographs* 15: 111–142.
- LINGHAM-SOLIAR T. 1992. The tylosaurine mosasaurs (Reptilia, Mosasauridae) from the upper Cretaceous of Europe and Africa. *Bull Inst R Sci Nat Belg Sci terre* 62: 171–194.
- LÜ J, AZUMA Y, CHEN R, ZHENG W AND JIN X. 2008. A new titanosauriform sauropod from the Early Late Cretaceous of Dongyang, Zhenjiang Province: *Acta Geol Sin* 82(2): 225–235.
- LÜ J, XU L, ZHANG X, HU W, WU Y, JIA S AND JI K. 2007. A new gigantic sauropod dinosaur with the deepest known body cavity from the Cretaceous of Asia. *Acta Geol Sin* 81(2): 167–176.
- MATEUS O, JACOBS LJ, POLCYN M, SCHULP AS, NETO AB AND ANTUNES MT. 2008. Dinosaur and turtles from the Turonian of Iembe, Angola In: *LIVRO DE RESUMOS DE “TERCER CONGRESO LATINOAMERICANO DE PALEONTOLOGÍA DE VERTEBRADOS”* 156 Neuquén, Argentina.
- MATEUS O, JACOBS LJ, POLCYN M, SCHULP AS, VINEYARD D, NETO AB AND ANTUNES MT. 2009. The oldest African eucryptodiran turtle: *Angolachelys mbaxi*, from the Cretaceous of Angola. *Acta Paleo Polonica* 54(4): 581–588.
- thosaurus sciuttoii* from the early Late Cretaceous Bajo Barreal Formation of Chubut Province, Argentina. *J Vert Paleont* 24(1): 107–120.
- MO JIN-YOU JIN-YOU, MO XIN, HUANG CHUO-LIN, ZHAO ZHONG-RU, WANG WEI AND XU. 2008. A new titanosaur (Dinosauria: Sauropoda) from the Late Cretaceous of Guangxi, China. *Vertebrat Palasiatic* 46(2): 147–156.
- PANG Q AND CHENG Z. 2000. A new family of sauropod dinosaur from the Upper Cretaceous of Tianzhen, Shanxi Province, China. *Acta Geol Sin* 74(2): 117–125.
- POLCYN M, JACOBS LJ, MATEUS O AND SCHULP AS. 2009. New specimens of *Angolasaurus bocagei* and comments on the early radiations of plioplatecarpine mosasaurs. *J Vert Paleont* 29(3): 165A.
- RAUHUT OWM AND WERNER C. 1997. First record of Maastrichtian sauropod dinosaur from Egypt. *Palaeo Africana* 34: 63–67.
- ROMER AS. 1956. *Osteology of the reptiles*. Chicago, University of Chicago, 772 p.
- RUBILAR-ROGERS D. 2006. Titanosaurs from the Atacama Desert. *J Vert Paleont* 26(3): 117A.
- SALGADO L, GARRIDO A, COCCA SE AND COCCA JR. 2004. Lower Cretaceous rebbachisaurid sauropods from Cerro Aguada Del León, Neuquén Province, northwestern Patagonia, Argentina. *J Vert Paleont* 24(4): 903–912.
- SCHULP AS, POLCYN MJ, MATEUS O, JACOBS LJ AND MORAIS ML. 2008. A new species of *Prognathodon* (Squamata, Mosasauridae) from the Maastrichtian of Angola, and the affinities of the mosasaur genus *Liodon*. *Proceedings of the Second Mosasaur Meeting, Fort Hays Studies Special Issue* 3, Fort Hays State University, Hays, Kansas, p. 1–12.
- SMITH JB, LAMANNA MC, LACOVARA KJ, DODSON P, SMITH JR, POOLE JC, GIEGENGACK R AND ATTIA Y. 2001. A giant sauropod from an Upper Cretaceous Mangrove Deposit in Egypt. *Science* 292: 1704–1706.
- STROMER E. 1932. *Ergebnisse der Forschungsreisen Prof. E. Stromer in den Wüsten Ägyptens*. II. Wirbeltier-Reste der Baharijestufe (unterstes Cenoman). #11. Sauropoda. *Abh. Bayer. Akad. Wissensch. Math.-naturwiss. Abteilung* 10: 1–21.
- UPCHURCH P, BARRETT PM AND DODSON P. 2004. Sauropoda. p. 259–322 In: WEISHAMPEL DB, DODSON P AND OSMÓLSKA H (Eds). *The Dinosauria*. 2nd ed.



NEW SAUROPOD DINOSAUR FROM ANGOLA

- In: WEISHAMPEL DB, DODSON P AND OSMOLSKA H (Eds), *The Dinosauria*, 2nd ed., California University Press, p. 517–606.
- WILHITE R. 2003. Biomechanical reconstruction of the appendicular skeleton in three North American Jurassic sauropods. Ph.D. dissertation. Louisiana State University, Baton Rouge, USA.
- WILSON J. 2002. Sauropod dinosaur phylogeny: critique and cladistic analysis. *Zool J [London]* 136(2): 215–275.
- XU X, ZHANG X, TAN Q, ZHAO X AND TAN L. 2011. A new titanosaurian sauropod from Late Cretaceous of Inner Mongolia, China. *Acta Geol Sin* 80: 20–26.
- YOU H, JI Q, LAMANNA MC, LI J AND LI Y. 2011. A new titanosaurian sauropod dinosaur with opisthocoelous vertebral centra from the Early Late Cretaceous of Inner Mongolia, China: *Acta Geol Sin* 78(4): 907–911.