



Anais da Academia Brasileira de Ciências

ISSN: 0001-3765

aabc@abc.org.br

Academia Brasileira de Ciências

Brasil

Cunha, Fabiane; Gómez, Daniel R. Sosa; Silva, Jose J. da; Alexandre, Talita M.; Moscardi, Flávio
Genetic diversity of the sunflower caterpillar (*Chlosyne lacinia saundersii* Doubleday and Hewitson)
(Lepidoptera: Nymphalidae) populations determined by molecular RAPD markers
Anais da Academia Brasileira de Ciências, vol. 82, núm. 4, 2010, pp. 1127-1136
Academia Brasileira de Ciências
Rio de Janeiro, Brasil

Available in: <http://www.redalyc.org/articulo.oa?id=327117686028>


- How to cite
- Complete issue
- More information about this article
- Journal's homepage in redalyc.org

redalyc.org

Scientific Information System

Network of Scientific Journals from Latin America, the Caribbean, Spain and Portugal

Non-profit academic project, developed under the open access initiative



Anais da Academia Brasileira de Ciências (2010) 82(4): 1127-1136
(Annals of the Brazilian Academy of Sciences)
ISSN 0001-3765
www.scielo.br/aabc

**Genetic diversity of the sunflower caterpillar
(*Chlosyne lacinia saundersii* Doubleday and Hewitson)
(Lepidoptera: Nymphalidae) populations determined
by molecular RAPD markers**

FABIANE CUNHA¹, DANIEL R. SOSA GÓMEZ², JOSE J. DA SILVA²,
TALITA M. ALEXANDRE¹ and FLÁVIO MOSCARDI³

¹Pós-Graduação em Entomologia, UFPR, Embrapa Soja, Laboratório de Patologia de Insetos
Caixa Postal 231, 86001-970 Londrina, PR, Brasil

²Embrapa Soja, Laboratório de Patologia de Insetos, Caixa Postal 231, 86001-970 Londrina, PR, Brasil

³Universidade Estadual de Londrina – Departamento de Agronomia Laboratório de Controle Microbiano de Insetos
Caixa Postal 6001, 86051-990 Londrina, PR, Brasil

Manuscript received on September 17, 2009; accepted for publication on September 17, 2010

ABSTRACT


Chlosyne lacinia saundersii is one of the most important pests of sunflower and it is the main target of insecticide applications. Larvae were collected in Londrina (PR), Santa Maria (RS), Dourados (MS), Ribeirão Preto (SP), Brasília (DF), Barreiras (BA), Uberaba (MG) and Vilhena (RO). Genomic DNA was extracted and amplified with ten-multiplex primers, which produced 101 loci. The size of the RAPD amplicons ranged from 180 to 2564 bp. Polymorphism among populations ranged from 31% to 67%, with the highest polymorphisms of 57% and 67% being detected in Uberaba and Vilhena populations, respectively. Populations with the highest similarity determined with Dice coefficient were from Ribeirão Preto and Barreiras, while insects from Londrina showed the highest similarity among them. Gene flow of *C. lacinia saundersii* 1.1 was lower than those previously observed for the noctuid *Anticarsia gemmatalis* Hübn. suggesting that *C. lacinia saundersii* populations are more isolated than the ones of this noctuid. Through the Analysis of Molecular Variance (AMOVA), RAPD variance was 33.64% among geographical populations and 66.36% within populations. These results suggest that populations of *C. lacinia saundersii* are genetically structured.

Key words: amova, gene flow, molecular marker, sunflower pest.

INTRODUCTION

The sunflower caterpillar *Chlosyne lacinia saundersii* Doubleday and Hewitson (1849) (Lepidoptera: Nymphalidae), a Neotropical species, is the main sunflower defoliator in Brazil (Ungaro 1981, Nakano et al. 1981, Gallo et al. 2002). It was first cited in Brazil by Maranhão (1945), in Piracicaba, SP, being further found in other states, as Paraná (Silva et al. 1968, Moscardi 1983, G. L. Villas Boas et al. unpublished data) Rio de Janeiro

1984) and São Paulo (Silva et al. 1968, Boiça Jr. 1984). Depending on the infestation level, larvae can cause up to 100% defoliation (Nakano et al. 1981). Literature on *C. lacinia saundersii* is scanty and most of the papers is related to controlling this insect with chemical insecticides, with very few studies on its biology, ecology and none on the genetic structure of populations of this insect. Papers related to genetic aspects are restricted to the genetics of color variation in larvae



Rica reported that the larva of *C. lacinia saundersii* is polymorphic, with coloration ranging from black to orange. In Brazil, Gallo et al. (2002) described the larva of *C. lacinia saundersii* as of black coloration, while Zucchi et al. (1993) described the larva as orange in color and covered with black scoli and setae. As this insect is an agricultural pest and considering the sunflower expansion (from 77.000 ha to 94.000 ha. i.e., 22.1%) (Lazzaroto et al. 2005) in the last six years in Brazil, it has a favorable habitat in terms of food resources, which can make possible the increase in the genetic flow thus, reducing the genetic differentiation. Therefore, studies related to the population genetics would be important for the development of management strategies mainly in species that faced changes in their habitats (Lougheed et al. 2000).

In the present study we have chosen, among the available molecular techniques, to use the RAPD (Random Amplified Polymorphic DNA), as it identifies the DNA polymorphism amplified at random through the use of a sole initiator of arbitrary sequence without previous knowledge of the species genome moreover this technique is of reduced cost (Williams et al. 1990). Although this technique results in amplification of random sequences, several authors have demonstrated its efficiency to distinguish genotypes in insect populations (eg. Carvalho and Vieira 2001, Oliveira et al. 2002, Lopes-da-Silva et al. 2004). Considering that *C. lacinia saundersii* is an insect of distinct geographical and climatic conditions, this study aimed to evaluate the intra and interspecific genetic variation in this insect, which can contribute to the development of proper strategies for the integrated management of this pest.

MATERIALS AND METHODS

INSECT COLLECTION


Second to fourth-instar *C. lacinia saundersii* larvae were collected in eight Brazilian states, (Table I). Collections were made from December 2007 to June 2008 in sunflowers or in other host plants and sent to the Laboratory of Entomology of Embrapa Soja, Londrina, PR.

ited to record for the occurrence of parasitoids and entomopathogens. Last instar larvae were placed in 15 mL sealed plastic vials containing dehydrated silica gel, which were maintained at -15°C for further DNA extraction. Thirty larvae were used for each population, being 10 of the nigra, 10 of the bicolor, and 10 of the rufa coloration.

DNA EXTRACTION

The DNA was extracted according to the protocol of Rogers and Bendich (1988), with few modifications. The extraction was made from the larval head to avoid a possible contamination with parasitoids in the larval hemocele (D.R. Sosa Gómez, unpublished data). Each sample was macerated in $480\mu\text{l}$ of extraction buffer at a final concentration of 200mM Tris-HCl (pH 8.0), 70mM EDTA, 2M NaCl, and 1% β -Mercaptoethanol. After addition of $120\mu\text{l}$ of cetyltrimethyl ammonium bromide 10%, the samples were kept at 65°C for 5 min. Then, $6\mu\text{l}$ of Proteinase K (10 mg/ml) was added and the samples were incubated at 65°C for 60 min. After that the samples were kept at environment temperature and centrifuged at 16.000 rpm for 15 min. The supernatant ($500\mu\text{l}$) was transferred to another microcentrifuge tube and added to it the same volume of chloroform/isoamyl alcohol (24:1).

After homogenization, each sample was centrifuged at 16.000 rpm for 15 min and, then, the aqueous phase ($400\mu\text{l}$) was transferred to another tube where the nucleic acids were precipitated with the same volume of cold isopropanol plus 45% of the volume of ammonium acetate (10M). DNA samples were mixed by gently shaking the tube and kept at -20°C for 2 h at -4°C overnight. The samples were, then, centrifuged at 14.000 rpm for 15 min, discarding the supernatant and washing the pellets with $300\mu\text{L}$ of cold ethanol 70%. The pellets were dry, with the tubes inverted, in environment temperature for 10 min, being, then, re-suspended with $100.8\mu\text{l}$ of RNase solution ($40\mu\text{g}/\text{ml}$) plus TE. After that, the tubes were placed in an environmental chambers from 30 min to 2 h for RNA degradation. Aliquots of DNA were kept at -15°C and defrost when



GENETIC DIVERSITY OF THE SUNFLOWER CATERPILLAR

TABLE I
Data on *Chlosyne lacinia saundersii* collection sites.

Site	Host Plant	Date	Coordinates
Barreiras – BA	<i>Helianthus annuus</i>	11/06/2008	12°07' 29 79"S 44°59' 14 59"O
Brasília – DF	<i>Tithonia diversifolia</i>	10/03/2008	15°43' 43 76"S 47°53' 58 61"O
Dourados – MS	<i>Parthenium hysterophorus</i>	18/01/2008	22°16' 59 06"S 54°49' 18 06"O
Londrina – PR	<i>Helianthus annuus</i>	13/12/2007	23°11' 29 74"S 51°10' 59 93"O
Ribeirão Preto – SP	<i>Tithonia diversifolia</i>	30/04/2008	21°10' 12 18"S 47°50' 51 55"O
Santa Maria – RS	<i>Helianthus annuus</i>	31/01/2008	29°43' 41 34"S 53°44' 40 89"O
Uberaba – MG	<i>Helianthus annuus</i>	10/03/2008	19°29' 54 64"S 48°02' 19 10"O
Vilhena – RO	<i>Oryza sativa</i>	05/05/2008	12°07' 24 85"S 60°17' 24 71"O

USA), and its integrity was evaluated through electrophoresis in agarose 0.8% gel and colored with 10 mg/ml of ethidium bromide. The electrophoresis was conducted with buffer of 10mM of NaOH and pH adjusted to 8.5 with boric acid (Brody and Kern 2004). The DNA was visualized and the image was digitalized using a transilluminator with a system of image capture L-PIX – ST and L-PIX IMAGE 7.1M Pixel. Images were captured with the software L-PIX IMAGE 1.0.1 (Loccus Biotecnologia, São Paulo – SP).

RAPD REACTIONS

The amplification reactions were conducted in a volume of 25µl containing about 9ng of DNA, Milli Q sterile water, buffer 10×, 0.4µM of primer, 2.4 mM of MgCl₂, 0.1mM of dNTP and Taq polymerase enzyme (Gibco BRE) (1U). The independent control of the reactions was conducted without the template DNA, when the amplification reaction with each primer was done. The primers that did not produce clear amplification were not considered in the analysis. The PCR was done using a thermocycler PTC 200 DNA Engine™ (MJ Research, Scientific Support, Hayward, CA) 3.8 version, with the following program: 45 cycles at 94°C for 15 sec, 39°C for 30 sec, and 72°C for 1 min, and a final extension of 72°C for 7 min. With a volume of 25µl, the product of RAPD was submitted to electrophoresis in agarose 1.3% and buffer SB 1× at 120 volts. The Lambda DNA, digested with the endonucleases *EcoRI*, *HindIII* and *BamHI*, was used as a marker

DATA ANALYSIS

Digital photos from gels were visually analyzed indicating the presence (1) and absence (0) of amplified bands with the different primers, assuming the occurrence of two alleles per locus. Only conspicuous and consistent bands, as well as their absence, were considered to generate the binary matrix. The similarity among samples was calculated using the Dice Coefficient by the formula $2h/(a+b)$, being “h” the number of observed bands and “a+b” the total number considering presence and absence of bands (Dice 1945). This matrix was submitted to cluster analysis using the unweighted pair group method with arithmetic mean (UPGMA). The procedures were conducted through the program NTSYSpc (Numerical Taxonomy (Rohlf 2000)). To verify consistency between the similarity matrix and the obtained dendrogram, the coefficient of cophenetic (r) was calculated according to Sokal and Rohlf (1962). The local stability of the groupings was estimated by the bootstrap analysis, with 1.000 replications, using the computer program Applied Maths Bionumerics (1999).

The structure of the populations was determined by the analysis of variance for molecular data (AMOVA) by using the program Arlequin (Schneider et al.

RESULTS

The ten used primers generated 101 loci, characterizing a high level of polymorphism in the *C. lacinia saundersii* populations. The number of obtained bands ranged from





Fig. 1 – Intra specific variation OPA-05 obtained by genomic DNA amplification of *Chlosyne lacinia saundersii* from Santa Maria, RS and Uberaba, MG. As molecular marker the fago Lambda DNA, digested with the *EcoRI*, *HindIII*, and *BamHI* endonucleases, was used.

from 180 to 2.564 pb. No monomorphic loci were observed among the studied populations despite being observed within each population (Table II).

TABLE II
Number of RAPD loci for *Chlosyne lacinia saundersii*.

Primer	Sequence	Number of bands
OPA – 01	5'-CAGGCCCTTC-3'	8
OPA – 05	5'-AGGGGTCTTG-3'	19
OPA – 09	5'-GGGTAACGCC-3'	8
OPA – 17	5'-GACCGCTTGT-3'	10
OPB – 10	5'-GTGAGGCGTC-3'	11
OPC – 02	5'-GTGAGGCGTC-3'	5
OPC – 06	5'-GAACGGACTC-3'	12
OPC – 10	5'-TGTCTGGGTG-3'	8
OPF – 03	5'-CCTGATCACC-3'	11
OPM – 02	5'-ACAACGCCTC-3'	9

The polymorphism among the populations ranged from 31% to 67%, with the ones with the greater number of polymorphic loci (57% e 67%) being from Uberaba and Vilhena, respectively. However the smaller polymorphism was detected for the population from Brasília (31%), followed by the one from Ribeirão Preto (35%) (Table III).

The largest proportion of individuals from Barreiras, Brasília, Ribeirão Preto, Dourados, Londrina, and Santa Maria formed six main groups according to their geographical location (Fig. 2). However, those from Uberaba and Vilhena presented similarities with indi-


the highest similarity were collected in the region of Londrina (Fig. 2).

The populations with a close similarity among them were those from Londrina, Dourados and Santa Maria, which presented Dice coefficients ranging from 0.765 to 0.775 (Fig. 2). The most divergent population was the one from Vilhena, RO, with a Dice coefficient of 0.695. This population presented the highest value of polymorphic loci differently from the Brasília and Ribeirão Preto populations, which were the least polymorphic ones (Table III).

TABLE III
Number of polymorphic bands produced by the primers used for the characterization of *Chlosyne lacinia saundersii* populations.


Origin of the population	Number of polymorphic loci (%)
Barreiras	42 (41)
Brasília	31 (30)
Dourados	48 (47)
Londrina	40 (39)
Ribeirão Preto	35 (34)
Santa Maria	46 (45)
Uberaba	58 (57)
Vilhena	68 (67)

Larval color polymorphism was presented by orange larvae (rufa), those the ones with black as general color with orange dorsal spots (bicolor), and the ones



GENETIC DIVERSITY OF THE SUNFLOWER CATERPILLAR





The bootstrap analysis was consistent with the Dice UPGMA dendrogram for the clusters (Fig. 2). The AMOVA results presented a fixation index Φ_{st} of 0.34 indicating that 33.64% of the genetic variation occurred among populations, and 66.36% occurred within them ($P < 0.001$) (Table IV).

DISCUSSION

Populations from Londrina, Dourados and Santa Maria were genetically close (Dice coef. = 0.76) compared to populations from Vilhena, Uberaba, Barreiras and Ribeirão Preto. The highest genetic dissimilarity (Dice coef. = 0.71–0.74) was observed between populations from Vilhena and Uberaba. The most heterogeneous population was from Vilhena, as samples from this population are found interspersed among populations from other localities.

The gene flow indexes (N_m) ranged from 0.9 between the populations of Londrina and Ribeirão Preto, up to 6.7 between the populations of Vilhena and Uberaba (Table V). The gene flow index considering all the *C. lacinia saundersii* studied populations was 1.1, which is relatively low when compared, for instance, to populations of *Anticarsia gemmatilis* (Lepidoptera: Noctuidae), with a gene flow of 3.05 (Sosa-Gómez 2004). According to this author, the great flying capacity of this species would help to explain the high similarity among populations, since the geographical distances among them were large. Conversely, the similarity among populations of *C. lacinia saundersii* is lower and, consequently, there are lower interchange of genetic characteristics among them probably because the sunflower cultivated area is still small and patchy, which hampers the gene flow among populations.


The area cultivated with sunflower in Brazil is currently of 94.000 hectares, while soybean is grown over 21 million hectares (CONAB 2008). The continuity of soybean areas could be favoring a greater migration and, thus, a larger gene flow index of the velvetbean caterpillar (*A. gemmatilis*) in relation to the sunflower caterpillar due to the relative discontinuity of sunflower cultivated areas in the country. On the other hand,

there have been no studies related to the association of genotypes of *C. lacinia saundersii* with these natural plant hosts. On the other hand, there are known studies with the fall armyworm, *Spodoptera frugiperda* (Lepidoptera: Noctuidae), which confirmed a greater genetic similarity among populations obtained from a same plant host (McMichael and Pashley 1999, Busato et al. 2004). It suggests that the host plant represents an important factor in the natural selection process. Therefore, more detailed studies on geographical barriers, distribution, and abundance of these hosts can help to clarify the impact of host plants on *C. lacinia saundersii* genetic flow.

Busato et al. (2004) evaluated the genetic diversity of *S. frugiperda* using AFLP markers and observed that studied populations were not different as there were overlapping of individuals from distinct populations in rice and/or corn. Almost nothing is known about the migration behavior of the sunflower caterpillar. Therefore, the origin and frequency of possible migrations are not known for this insect when occurring in sunflower and wild host plants in Brazil. *C. lacinia saundersii* occurs in diverse regions where sunflower is present in this country. Thus, it is possible that this species is well adapted to the different conditions of climate and topography.

Through the AMOVA, 34.64% of the total genetic variability could be attributed to differences among populations of *C. lacinia saundersii*, and 66.3% to differences within populations, which is an indicative that the studied populations were structured. Similar values were obtained with *Sternechus subsignatus* (Coleoptera: Curculionidae) that presented difference, among populations of 29% (Sosa-Gómez et al. 2008). For *A. gemmatilis* and *S. frugiperda* the variation among populations were of 14% and 18%, respectively (Sosa-Gómez 2004, Martinelli et al. 2006). These lower values may be related to the high gene flow in populations of these lepidopterous species.

The RAPD markers used in the present study were efficient for the genetic differentiation of *C. lacinia saundersii* populations. This type of work is the only one done for this species up to the moment. However, addi-



GENETIC DIVERSITY OF THE SUNFLOWER CATERPILLAR



Fig. 3 – Map with the geographical distances among the populations of *Chlosyne lacinia saundersii*. Was based on the coordinates collection site and expressed in kilometers through the software GPS TrackMaker[®] 13.4 (<http://www.gpstm.com/>).

TABLE IV
Analysis of molecular variance (AMOVA) for eight geographical populations
of *Chlosyne lacinia saundersii*.

Source of variation	g.l.	Sum of squares	Components of variance	Percent of variation	Probability	Fixation index Φ_{st}
Among populations	7	739.452	3.66 (V_a)*	33.64	$P < 0.001$	0.34
Within populations	207	1 495 888	7.23 (V_b)*	66.36	$P < 0.001$	–




TABLE V
Paired comparisons among populations of *Chlosyne lacinia saundersii*
through the Nei coefficient for the gene flow (*Nm*) estimate.

Values of Nm	Vilhena	Uberaba	Brasília	Ribeirão Preto	Santa Maria	Londrina	Dourados	Barreiras
Vilhena	xxxxxx	6.7	2.3	2.1	2.7	2.1	3.2	2.0
Uberaba		xxxxxx	2.5	2.2	2.9	2.0	2.8	2.1
Brasília			xxxxxx	1.0	1.7	1.2	1.7	1.5
Ribeirão Preto				xxxxxx	1.2	0.9	1.3	1.6
Santa Maria					xxxxxx	1.8	3.5	1.6
Londrina						xxxxxxx	1.9	1.1
Dourados							xxxxxx	1.8
Barreiras								xxxxxx

ACKNOWLEDGMENTS

The authors thank Dr. Vanoli Fronza for revising the manuscript, Dr. Eliseu Binneck for the statistical support, Silvana R. Rockenbach for the technical assistance; for insect collections, the following researchers: Débora Pires de Paula (Embrapa Recursos Genéticos e Biotecnologia), Mônica C. Pires and Pedro V. Lima (Fundação Bahia), Marino Alves (UNESP de Ribeirão Preto), Marley Utumi (Embrapa Rondonia), Talita M. Alexandre and Jovenil J. da Silva (Embrapa Soja), Rejane C.R.K. Roggia (UFMS), and Neylson Eustáquio Arantes (Embrapa Soja/Epamig – Uberaba); Adair V. Carneiro and Danilo Estevão for the support with the pictures and figures. This work was supported by Embrapa Soja and Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (Capes) and was approved for publication by the Publications Committee of Embrapa Soja with number 01/2009.

RESUMO

Chlosyne lacinia saundersii é uma das mais importantes pragas da cultura do girassol e o principal alvo das aplicações de inseticidas. As larvas foram coletadas em Londrina (PR), Santa Maria (RS), Dourados (MS), Ribeirão Preto (SP), Brasília (DF), Barreiras (BA), Uberaba (MG) e Vilhena (RO). O DNA genômico foi extraído e amplificado com dez primers, que produziram 101 locos. O tamanho das amplificações de

respectivamente. As populações com maior similaridade determinada com o coeficiente de Dice foram de Ribeirão Preto e Barreiras, enquanto os insetos coletados em Londrina apresentaram maior similaridade entre eles. O fluxo gênico de *C. lacinia saundersii* de 1,1 foi menor que o observado para *Anticarsia gemmatalis* Hübner Noctuidae, sugerindo que as populações de *C. lacinia saundersii* estão mais isoladas do que estes noctuídeos. Através da análise de variância molecular (AMOVA), RAPD a variação foi de 33,64% entre as populações geográficas e 66,36% dentro das populações. Estes resultados sugerem que as populações de *C. lacinia saundersii* são geneticamente estruturadas.

Palavras-chave: amova, fluxo gênico, marcador molecular, praga do girassol.


REFERENCES

APPLIED MATHS BIONUMERICS. 1998. The integral study of biological relationships. Manual, version 1.01, Kortrijk, Belgium, 154 p.

BOIÇA JR AL, BOLONHEZI AC AND PACCINI NETO J. 1984. Levantamento de insetos pragas e seus inimigos naturais em girassol (*Helianthus annuus* L.), cultivada em primeira e segunda época, no município de Seviria-MS. An Soc Entomol Bras 13: 189–196.


BRODY JR AND KERN SE. 2004. Sodium boric acid: a Tris-free, cooler conductive medium for DNA eletrophoresis. BioTechniques 36: 214–216.

DUNN, G.B., GRANTHAM, J.E., AD. OLIVER, A.C., MORGAN, J.



GENETIC DIVERSITY OF THE SUNFLOWER CATERPILLAR

- (J.E. Smith, 1797) (Lepidoptera: Noctuidae) associadas ao milho e arroz no Rio Grande do Sul. *Neotrop Entomol* 33: 709–716.
- CAMPOS-FARINHA AEC, PINTO NPO AND GOVONE JS. 1997. Estudo do comportamento e desenvolvimento de lagartas de *Chlosyne lacinia saundersii*, Doubleday and Hewitson (1849) (Lepidoptera: Nymphalidae), no ataque a uma planta de girassol (*Helianthus annuus* L.). *Arq Inst Biol* 64: 143–147.
- CARVALHO AOR AND VIEIRA LGE. 2001. Determinação das condições ótimas para análises de PCR-RAPD em *Atta sexdens rubropilosa* Forel (Hymenoptera: Formicidae). *Neotrop Entomol* 30: 593–600.
- COMPANHIA NACIONAL DE ABASTECIMENTO, CONAB. 2008. Acompanhamento da safra de girassol 2007/2008. Disponível em <<http://www.conab.gov.br/conabweb/download/moc/titulos/T51s2007-2008.pdf>>.
- DE VRIES PJ. 1987. The butterflies of Costa Rica and their natural history Papilionidae, Pieridae, Nymphalidae. Princeton University Press, 327 p.
- DICE LR. 1945. Measures of the amount of ecologic association between species. *Ecology* 26: 297–302.
- EDWARDS WH. 1893. Notes on a polymorphic butterfly, *Synchlora lacinia*, Geyer (in Hub. Zutr.), with description of its preparatory stages. *Can Ent* 25: 286–291.
- GALLO D ET AL. 2002. Entomologia Agrícola, Piracicaba, FEALQ, 920 p.
- GORODENSKI SA. 1969. The genetics of three polymorphic larval color forms of *Chlosyne lacinia* (Lepidoptera, Nymphalidae). *Genet Res* 14: 332–336.
- LAZZAROTO JJ, ROESSING AR AND MELLO HC. 2005. O agronegócio do girassol no mundo e no Brasil. In: GIRASSOL NO BRASIL (Ed), Londrina: Embrapa Soja, p. 15–42.
- LOPES-DA-SILVA M, TONET GEL AND VIEIRA LGE. 2004. Characterization and genetic relationships among brazilian biotypes of *Schizaphis graminum* (Rondani) (Hemiptera: Aphididae) using RAPD markers. *Neotrop Entomol* 33: 043–049.
- LOUGHEED SC, GIBBS HL, PRIOR KA AND WEATHERHEAD PJ. 2000. A comparison of RAPD versus microsatellite DNA markers in population studies of the *Matsigena rattlesnake*. *J Hered* 91: 458–463.
- MARANHÃO ZC. 1945. *Chlosyne lacinia saundersii*, praga *doptera frugiperda* (Lepidoptera: Noctuidae) Pop Associated to Maize and Cotton Crops in Brazil. *Entomol* 99: 519–526.
- MCMICHAEL M AND PASHLEY DP. 1999. Different amplified fragment-length polymorphisms in fall armyworm (Lepidoptera: Noctuidae) host strains. *Ann Entomol Soc Am* 92: 175–181.
- MOSCARDI F. 1983. Plantas Hospedeiras da lagarta do girassol, *Chlosyne lacinia saundersii*, no Estado do Paraná. EMPRESA BRASILEIRA DE PESQUISA AGROPECUÁRIA, Centro Nacional de Pesquisa de Soja, Londrina, Resultados de Pesquisa de Girassol Londrina, p. 25–30.
- MOSCARDI F, SOSA-GÓMEZ DR AND CORSO IC. 1999. Invertebrados associados ao girassol e seu manejo. GIRASSOL NO BRASIL (Ed), Londrina: Embrapa Soja, p. 471–500.
- NAKANO O, SILVEIRA NETO S AND ZUCCHI RA. 1999. Entomologia Econômica. Piracicaba, Livroceres, 300 p.
- NECK RW, BUSH L AND DRUMOND BA. 1971. Effect of associated lethals and brood effect on larval color polymorphism of the patch butterfly *Chlosyne lacinia*. *Heredity* 26: 73–84.
- OLIVEIRA CM, FUNGARO MH, CAMARGO LE AND LOPES JRS. 2002. Análise comparativa da estabilidade de *Dalbulus maidis* (Delong and Wolcott) (Hemiptera: Coreidae) sobre diferentes métodos de preservação de DNA. uso de RAPD-PCR. *Neotrop Entomol* 31: 225–230.
- ROGERS SO AND BENDICH AJ. 1988. Extraction of DNA from plant tissues. *Plant Mol Biol* 6: 1–10.
- ROHLF FJ. 2000. Ntsys-Pc Numerical taxonomy and multivariate analysis system version 2.1. Exeter Software, Setauket, NY.
- SCHNEIDER S, ROESSLI D AND EXCOFFIER L. 2000. Arlequin, version 2.0: A software for population genetic analysis. Genetics and Biomtry Laboratory, University of Geneva, Geneva, Switzerland.
- SILVA AGA, GONÇALVES CR, GALVÃO DM, GONÇALVES AJL, GOMES J, SILVA MN AND SIMONI L. 1968. Catálogo dos insetos que vivem nas plantas do Brasil e seus parasitos e predadores. Rio de Janeiro, Ministério da Agricultura/laboratório central de patologia vegetal, 622 p.
- SOKAL RR AND ROHLF FJ. 1962. The comparison of



- gemmatalis* Hübner, 1818 (Insecta: Lepidoptera: Noctuidae). Genet Mol Biol 27: 378–384.
- SOSA-GÓMEZ DR, CORONEL N, BINNECK E, ZUCCHI MI AND ROSADO-NETO G. 2008. RAPD and mitochondrial DNA analysis of the soybean stalk weevil, *Sternechus subsignatus* (Coleoptera: Curculionidae). Bull Entomol Res 98: 475–481.
- UNGARO MRG. 1981. Recomendações técnicas para o cultivo do girassol. Correio Agrícola Bayer 2: 314–319.
- WILLIAMS JG, KUBELIK AR, LIVAK KJ, RAFALSKI LA AND TINGEY SV. 1990. DNA polymorphism amplified by arbitrary primers are useful as genetic markers. Nucleic Acids Res 18: 6531–6535.
- ZUCCHI RA, SILVEIRA NETO S AND NAKANO O. 1993. Guia de identificação de pragas agrícolas, Piracicaba, FEALQ, 139 p.