



Anais da Academia Brasileira de Ciências

ISSN: 0001-3765

aabc@abc.org.br

Academia Brasileira de Ciências

Brasil

Silva, Cesar A.; Madureira, Luiz A.S.

Source correlation of biomarkers in a mangrove ecosystem on Santa Catarina Island in southern Brazil

Anais da Academia Brasileira de Ciências, vol. 84, núm. 3, septiembre, 2012, pp. 589-604

Academia Brasileira de Ciências

Rio de Janeiro, Brasil

Available in: <http://www.redalyc.org/articulo.oa?id=32723112018>

- How to cite
- Complete issue
- More information about this article
- Journal's homepage in redalyc.org

redalyc.org

Scientific Information System

Network of Scientific Journals from Latin America, the Caribbean, Spain and Portugal

Non-profit academic project, developed under the open access initiative



Source correlation of biomarkers in a mangrove ecosystem on Santa Catarina Island in southern Brazil

CESAR A. SILVA and LUIZ A.S. MADUREIRA

Laboratório de Química Ambiental e Geoquímica Orgânica, Departamento de Química,
Universidade Federal de Santa Catarina (UFSC), 88040-900 Florianópolis, SC, Brasil

Manuscript received on July 28, 2010; accepted for publication on September 28, 2011

ABSTRACT

The relative distribution of several compounds identified in four samples of recently deposited sediments of the Itacorubi Mangrove located on the Santa Catarina Island, southern Brazil, was compared with similar data on compounds extracted from fresh leaves of three mangrove species (*Avicennia schaueriana*, the dominant species in the area, *Rhizophora mangle* and *Laguncularia racemosa*) and the Gramineae *Spartina alterniflora*. Terpenols, previously identified in mangrove species in northern Brazil, were also found. *A. schaueriana* mainly contains β -amyrin (90.6 $\mu\text{g g}^{-1}$ of extractable organic matter); low amounts of friedelin, betulin and germanicol were detected only in the leaf extract of this species. *R. mangle* also contained a significant amount of β -amyrin and it was the only species where taraxerol was detected. In contrast to the leaves, sediment extracts were dominated by germanicol, α -amyrin and campesterol. Despite its chemical lability, betulin was also detected. Two homologous series of α and ω -hydroxy fatty acids were detected in the acid-alkaline fraction. In spite of being reported in the literature as components of terrigenous plants, saturated ω -hydroxy acids were not identified. Our results indicate that although triterpenols may be used as biomarkers for mangrove-derived organic matter, their relative distribution can change according to the region.

Key words: biomarkers, leaf, mangrove, Santa Catarina, sediment.

INTRODUCTION

Mangrove ecosystems are characterized by high primary productivity, high levels of organic carbon, abundant detritus and anoxic/reduced conditions. The forests are best developed under tropical climates, where the coldest winter temperature is above 20°C and temperatures are fairly constant throughout the year. Mangroves are salt-tolerant trees and grow mainly along low-lying

depositional coasts and deltas, where the substrate is predominantly made of clay and silt rather than sand. Since nutrients are often adsorbed onto sediment particles, inputs of mangrove-derived organic matter into tidal creeks and estuaries are important for coastal zooplankton and fish populations (Lacerda 1998, Koch et al. 2003). Previous studies have emphasized the importance of leaf litter as the major component of the total litterfall of mangrove forests as a carbon source and thus in the identification of specific biomarkers (Koch et al. 2003, 2011). Hence, mangroves represent favorable environments where

Correspondence to: Luiz Augusto dos Santos Madureira
E-mails: lasmadureira@bol.com.br /
luiz.madureira@ufsc.br

an important uptake, conservation and accumulation of biomarkers such as hydrocarbons, triterpenols, sterols and fatty acids may occur (Mater et al. 2004, Koch et al. 2005, Farias et al. 2008).

The presence of biomarkers in sediments has been widely used as a proxy to evaluate possible sources of organic matter (Rieley et al. 1991). Organic compounds such as terpenols, *n*-alkanols, *n*-alkanes, fatty acids and hydroxy acids (Keinänen et al. 2003) and lignin phenols (Pempkowiak et al. 2006, Rezende et al. 2010) are particularly important to estuaries. However, it is well known that it is difficult to find correlations between organic compounds and a specific living organism, and other biogenic parameters (e.g. C, N, and P) and the isotope composition of individual compounds (Vogts et al. 2009) may be used for the assessment of specific sources (Madureira et al. 1997, Xu et al. 2001, Ruiz-Fernandéz et al. 2002, Bouillon et al. 2003). In the particular case of mangroves, the vegetation is well characterized and compounds like taraxerol and germanicol have been used as a lipid biomarker for mangrove-derived organic matter (Koch et al. 2005, 2011). Estuarine sediments are of great complexity due to the tidal influx of allochthonous organic matter and also the input of terrigenous debris from the local vegetation (Bouillon et al. 2003). Thus, it is important to employ as many parameters as possible to evaluate the dominant indicators for terrestrial and aquatic inputs. Apart from natural contributions, mangroves are also susceptible to inputs of pollutants from diffuse sources. In most coastal areas, mangroves are important receptors of pollutants from the atmosphere, continental runoff and tidal currents.

In recent years, the pressures of increasing population, food production, and industrial and urban development have caused a significant proportion of the world's mangrove resources to be destroyed. There are five well-defined mangrove areas at Santa Catarina Island, but they are under the constant threat of losing their protection due to changes in municipal and federal laws. Several studies have already empha-

sized the great importance of the mangroves located at Santa Catarina Island as natural habitats and food sources for many crustaceans, fishes and reptiles (Dutra 1998, Soriano-Sierra and Ledo 1998). In this regard, our contribution was to analyze biogenic parameters, as well as organic compounds, extracted from the four predominant plant species and sediment samples taken from the Itacorubi Mangrove ecosystem. The aim was to obtain information on the relative contribution of the leaves of mangroves species, bacteria and microalgae to the mangrove sediments. Sterols, triterpenols, *n*-alkanes, hydroxy and *n*-alkanoic acids are the principal compound classes that will be addressed in this study. Moreover, following the procedure described by Garcette-Lepecq et al. (2004), all of the leaf and sediments samples were submitted to sequential treatment with base and acid hydrolysis after being solvent-extracted. According to the authors, the bound acid fractions reflect differences in the linkage mode of the acid moieties and/or in the protection provided by the (macro) molecular structures into which they are incorporated.

Sterols are widely distributed in living organisms. Potential sources include sewage matter, estuarine phytoplankton and zooplankton and terrestrial vascular plants. Some types of sterols may also originate from a synthesis within the sediments and from secondary transformation products arising from the bioconversion of primary sterols (Volkman 1986). C_{28} and C_{29} desmethyl sterols in sediments that are believed to originate predominantly from vascular plants include: campesterol (24-methyl-5-cholesten-3 β -ol), β -sitosterol (24 α -ethyl-5-cholesten-3 β -ol) and stigmasterol (24 α -ethyl-5, (22E)-cholestadien-3 β -ol) (Heftmann 1971). The distribution of triterpenols in the leaves of mangrove species may vary significantly. Large amounts of betulin, lupeol, β -amyrin and taraxerol have been reported in the literature (Koch et al. 2003).

The within-class distributions of long-chain ($> C_{20}$) *n*-alkanes, *n*-alkanoic acids and *n*-alkanols

are often used to infer terrestrial inputs to sediments. These compounds are abundant in plant epicuticular waxes, but long-chain *n*-alkanes and *n*-alkanoic acids can also be found in marine algae (Volkman et al. 1980, Venkatesan and Kaplan 1982). As previously reported, in *R. mangle* leaves odd *n*-alkanes (C_{25} , C_{27} and C_{29}) are predominant (Hernandez et al. 2001).

With regard to unsaturated fatty acids, these mainly occur in bacteria and microalgae. However, their structure, especially the double bond location, can help to discriminate among such sources (Garcette-Lepecq et al. 2004). For hydroxy acids several sources can be considered. They can be originated during the diagenesis through microbial oxidation of fatty acids. Medium and long-chain α -hydroxy acids, as well as medium-chain

β -hydroxy acids, are also produced by microalgae and seagrasses (de Leeuw and Largeau 1993, de Leeuw et al. 1995, Versteegh et al. 1997).

EXPERIMENTAL

STUDY AREA

The Itacorubi Mangrove ecosystem is located at the southern limit of the mangrove distribution area of the Brazilian coast (coordinates: $27^{\circ} 34' 14''$ - $27^{\circ} 35' 31''$ S; $48^{\circ} 30' 07''$ - $48^{\circ} 31' 33''$ W) and occupies a surface area of 1.42 km^2 in an estuarine basin of 22.12 km^2 (Figure 1). Two rivers, the Itacorubi and the Sertão, 5.9 and 4.0 km long respectively, as well as natural and artificial channels, cross a well-developed forest (Dutra 1998).




Figure 1 – Map showing the location of Itacorubi Mangrove at Santa Catarina Island and the sampling locations.

Three mangrove species are present in the forest: *Rhizophora mangle*, *Laguncularia racemosa* and *Avicennia schaueriana*, in dense populations – approximately 2 900 per ha. *A. schaueriana*, the dominant species (90%), has some individual trees over five meters high, with a density of 800 adult trees per ha (Soriano-Sierra and Ledo 1998). *Spartina alterniflora* is a species of grass that grows in tallest forms at the outermost edge of the marsh, creating mud banks like islands with approximately 133 m of radius.

This ecosystem has been progressively eliminated during the past 40 years, although it is mentioned in the current national legislation as an “Area of Permanent Preservation”. Environmental changes have taken place in this mangrove system due to the impact of anthropogenic processes (Lago 1997). The mangrove served as an area for the disposal of solid waste material until 1980, receiving a total volume of 75,000 tons of domestic garbage, industrial and hospital waste, all of which were deposited in layers superposed on a sanitary landfill occupying a surface area of 12 ha. Artificial channels, 6.1 km long in total, have been built to discharge rainwater and domestic effluents to the coastal zone (Soriano-Sierra and Ledo 1998). The Itacorubi Mangrove has been used for subsistence fishing by a reduced number of fishermen but, due to its location, it is becoming an object of urban management, resulting in the elimination of some sectors to obtain land for the construction of residences and roads.

SAMPLING

Fresh leaves from *R. mangle*, *L. racemosa*, *A. schaueriana* and *S. alterniflora* were collected along transects from the sea side to the land side of the mangrove. In order to avoid possible variations in the lipid composition among the trees and leaves of different developmental stages, only fully mature foliage from adult individuals was collected. Also, trees with the presence of epiphytic colonization were avoided due to the potential contamination

of leaves. All leaves were transferred to and stored in previously-combusted (450 °C) glass jars sealed with aluminum foil-lined lids and kept frozen until freeze-drying in the laboratory. Sediment samples were collected from near the surface (0-5 cm depth) with a van Veen grab sampler at the same intertidal zone from which the leaves were collected. The sediment samples named ITA1 and ITA4 were taken from subtidal areas, whereas ITA2 and ITA3 were sampled from intertidal areas.

BULK PARAMETERS AND PARTICLE SIZE DISTRIBUTION

Immediately prior to lipid extraction, the storage solvent was evaporated and the sediment freeze-dried (Edwards F105) under vacuum and homogenized by grinding. Aliquots were taken for elemental analysis. Samples were decarbonated using 0.1 mol L⁻¹ HCl solution. Total organic carbon (TOC) and nitrogen were measured using a Carlo Erba EA1110 CHNS-O analyzer after being centrifuged, washed with deionized water and gently dried at 60 °C. Samples from all stations were collected for granulometric analysis following the procedure described by Suguio (1973).

PHYSICO-CHEMICAL PARAMETERS

Both pH and redox potentials (Eh) were measured using a portable instrument Ysi F-1100. The pH electrode was calibrated using 4.02 and 7.00 standards and measured *in situ*, using an Ag/AgCl electrode, by inserting the probes directly into different sections of the sample (1, 3 and 5 cm from the surface). The mV meter electrode was checked using deionized water. The electrodes were equilibrated over several minutes before taking a reading to be thoroughly washed and subsequently rubbed with fine emery paper after each measurement in order to prevent their poisoning by sulphide ions. Results were converted into Eh values, by adding 202 mV to the original values read (AgCl reference electrode, 25 °C).

EXTRACTION AND FRACTIONATION

The processes of extraction, separation and subsequent identification of compounds were based on the procedures developed by Goossens et al. (1989a, b) as schematically illustrated in Figure 2. The freeze-dried sediment samples weighing about 5 g were extracted with a mixture of $\text{CH}_2\text{Cl}_2:\text{CH}_3\text{OH}$ (2:1, v/v, 3x) by ultra-sound for 30 min each. In order to remove elemental sulfur, activated copper was added. The leaves were submitted to the same process except for the removal of elemental sulfur. The extracts obtained were rotary evaporated and dried over a stream of nitrogen. An aliquot of this extract (E1) was derivatized with acetyl chloride in methanol to convert fatty acids into their methyl esters.

Subsequently, very polar compounds were removed through a Pasteur pipette minicolumn over silica gel with ethyl acetate as the eluent. The eluate

was evaporated with a stream of nitrogen. This fraction was dissolved with bis-(trimethylsilyl) trifluoroacetamide (BSTFA). This mixture was heated (60 °C; 60 min) to convert alcohols into their corresponding trimethylsilyl ethers. The compounds were analyzed by means of gas chromatography-flame ionization (GC-FID) and gas chromatography-mass spectrometry (GC-MS).

After extraction the dried residues (R1) and the remaining aliquot of the underivatized extract (E1) were saponified with a 1.0 mol L^{-1} solution of KOH in $\text{MeOH}:\text{H}_2\text{O}$ (96:4) for 1 h. After cooling to room temperature, the mixture was acidified with 2.0 mol L^{-1} $\text{HCl}:\text{CH}_3\text{OH}$ (1:1) to pH 3.5, centrifuged and transferred to a separation funnel. The residues were washed subsequently with $\text{CH}_3\text{OH}/\text{H}_2\text{O}$, CH_3OH and CH_2Cl_2 . At each time the supernatants were transferred to the separation funnel. Water was added for a better separation of the CH_2Cl_2 layer. The $\text{MeOH}/\text{H}_2\text{O}$ layers were washed with CH_2Cl_2 (x2).




Figure 2 – Flow chart of the analytical scheme used to obtain the lipid fractions.

The combined CH_2Cl_2 phases were concentrated using rotary evaporation, dried over anhydrous sodium sulfate and derivatized as described above, providing the extracts E2 and E3 (Figure 2).

The dried residues after saponification (R2) were also treated with a 4.0 mol L^{-1} HCl solution (6 h; 100°C). After cooling, the reaction mixture was neutralized with a 4.0 mol L^{-1} solution of KOH to pH 8.5, freeze-dried and saponified. The extracts (E4) were washed, separated and derivatized as described above.

The four fractions were isolated to obtain the lipids in the different forms in which they are found in the organic matter. The lipid fraction (E1), or total extract, includes the lipid compounds not linked to the biopolymers. The "OH-labile" fraction (E2 and E3), released from the solvent-extracted sediments by saponification (alkaline hydrolysis), supplied the acid-linked lipids in the ester form. The extract E4 was obtained after the saponification step via acid followed by alkaline hydrolysis, thus referring to the "H-labile" fraction linked to the acids through amide and/or ether groups.

The GC-MS analysis was carried out on a Shimadzu QP5050A MS interfaced to a model 17A Shimadzu network system quadrupole GC, using electron ionization mode (electron energy 70 eV , ion source temperature 290°C , scanning from 50 to $550 m/z$, 2 scan s^{-1}), a $30 \text{ m} \times 0.25 \text{ mm}$ i.d. capillary column coated with $0.25 \mu\text{m}$ film of DB-5MS, He as the carrier gas and an oven programmed from 70°C (hold for 4 min) to 130°C at $20^\circ\text{C min}^{-1}$, from 130 to 310°C at $15^\circ\text{C min}^{-1}$, with a 15 min isotherm at 310°C . Compounds were identified based on their mass spectra, GC retention times, authentic standards (cholesterol, β -sitosterol, stigmasterol, α -cholestane, β -amyirin from Sigma-Aldrich; mixes of linear *n*-alkanes and fatty acids from AccuStandards), terpenols isolated from plants (provided by other researchers) and comparison with library mass spectra. Compounds were quantified based on the total ion current peak area of the

individual compounds and the relative response to the internal standards added and converted to mass based on the dry weight of the extracts.

RESULTS AND DISCUSSION

ELEMENTAL COMPOSITION AND GEOCHEMICAL PARAMETERS

Bulk sediment parameters, molar ratios and the redox conditions for all locations are listed in Table I. The content of silt plus clay ranged between 87.3 and 98.1 %. According to the Shepard's triangular diagram (Shepard 1954), these sediments can be classified as silted-clay. TOC values were higher in the intertidal sediment samples (ITA2 and ITA3: 4.27 and 3.32%, respectively), which may be due to the direct contribution of leaves from mangrove trees. These results are in agreement with the greater extractable organic matter (EOM) concentrations obtained in the dry sediments of these stations (ITA2 and ITA3: 16.60 and 13.54 mg g^{-1} sed. dry weight, respectively). However, as will be discussed in the next paragraph, the percentage of nitrogen also increased at both stations (ITA2 and ITA3) and the organic carbon:nitrogen ratio remained very low. Despite the TOC variation, the range of values found in the sediments is consistent with those previously reported for this mangrove region (Mater et al. 2004) and for other subtropical lagoons and estuaries (Koch et al. 2003).

The C/N molar ratios calculated for the mangrove leaves varied from 22.1 to 26.4 and are characteristic of terrigenous plants (Table I). However, the C/N ratios were much lower in the sediments, ranging from 7.2 at station ITA4 to 10.4 at ITA1. These are typical values corresponding to the marine organic matter contribution and indicate that the organic carbon from the mangrove leaves has been considerably remineralized or mixed with the marine organic matter. In fact, a previous work has reported that C:N ratios in mangrove soils under *R. mangle* and *A. schaueriana* averaged

TABLE I
Bulk sediment parameters and molar ratios in the Itacorubi mangrove stations.

Stations and mangrove species	Silt + clay (%)	TOC (%)	N (%)	H (%)	C/N	C/H	EOM (mg g ⁻¹)
ITA1	96.8	2.04	0.22	1.20	9.27	0.15	3.40
ITA2	89.1	4.27	0.41	1.66	10.41	0.12	16.60
ITA3	87.3	3.32	0.40	1.59	8.30	0.10	13.54
ITA4	98.1	1.86	0.26	1.47	7.15	0.21	1.08
Avic		41.94	1.90	5.92	22.07	0.59	
Lagun		45.17	1.71	5.62	26.42	0.63	
Spar		39.69	1.52	5.51	26.11	0.62	
Rhiz		45.42	1.82	6.41	24.96	0.67	

TOC: total organic carbon.

EOM: Extractable organic matter (mg g⁻¹ of dry weight sediment)

Values shown are the average of three replicate analyses with RSD < 3.8 %.

near 17 (Lacerda et al. 1995). The C/H molar ratio also indicated this depletion of organic carbon (Table I). The calculated C/H values were 0.59-0.67 in the leaves and decreased to 0.10-0.21 in the sediments possibly due to a partial loss of carbon and hydrogenation of the organic material.

The redox conditions were measured at three depths (1, 3 and 5 cm) at the locations where the sediment samples were collected. With the exception of station ITA1 (first centimeter), all values were negative (ranging from -444 to -250 mV), representing very reducing conditions (Baltzer 1982, Lallier-Vergès et al. 1998).

BIOMARKERS IN LEAVES AND SEDIMENTS

In this section we will discuss data on the *n*-alkane, monocarboxylic and hydroxy fatty acids, sterol and triterpenol concentrations and the distribution patterns of individual compounds in the surface sediments and leaves of the mangrove species. In terms of hydrocarbon distribution, total *n*-alkanes

(C₂₂ – C₃₃) showed a predominance of compounds with odd carbon numbers, mainly in the leaf samples. For the sediments, the total concentration was highest at station ITA3 (34.6 µg g⁻¹ of extractable organic matter - EOM) and the lowest value was found at station ITA4 (27.5 µg g⁻¹ EOM), with a maximum peak for C₂₉, typically found in terrigenous plants. The highest concentrations of hydrocarbons were detected in the leaf sample of *Avicennia* (38.8 µg g⁻¹ of dry leaf – DL). In contrast, the lowest value was found in the Gramineae sample (8.8 µg g⁻¹ DL), which is nearly 4.4 times lower than that of *Avicennia*. The lower concentration of hydrocarbons in *S. alterniflora* can be explained because it is a non-vascular plant species with lower cellulose and lignin contents.

The Carbon Preference Index (CPI) was calculated in the range of C₂₃ - C₃₁ for *n*-alkanes, and the results showed CPI > 2 for the sediments, which indicates a large contribution from terrigenous particulate material (Table II). For comparison, CPI values of hydrocarbons from

the leaf samples were also calculated. The highest value was obtained for *Laguncularia* (CPI = 11.2), with *n*-C₂₉ concentration nearly three times higher than that of *n*-C₂₇.

With respect to total monocarboxylic acids, the distribution in the sediment samples ranged from C₁₀ to C₃₂ for the linear compounds. Long-chain components with an even-over-odd distribution are characteristic of contributions from higher plants. Several monounsaturated and isoprenoic acids were also detected (Table II). In the leaf samples the major compounds were a few monocarboxylic acids, with the predominance of C₁₆ and C₁₈. *A. schaueriana* (black mangrove) mainly contained C₁₆ monocarboxylic acid (87.5 µg g⁻¹ DL) with a very low amount of C₁₈ (0.7 µg g⁻¹ DL), whereas the mangrove species *L. racemosa* (white mangrove) showed the highest values for C₁₆ and C₁₈ (159.9 and 120.6 µg g⁻¹ DL, respectively) and also a very high content of the C_{18:3} fatty acid (206.5 µg g⁻¹ DL). The latter compound was not detected in any of the mangrove sediment samples, possibly due to the instability of this compound leading to its rapid oxidation.

Several monounsaturated fatty acids were identified in the sediment samples, but none of them was seen in the leaf samples. These compounds are probably produced by microorganisms, since most bacteria do not contain fatty acids with more than one C=C bond. Significant quantities of the acids C_{16:1}, C_{18:1}, C_{19:1}, and C_{22:1} appeared predominantly (Table II), with both *cis* and *trans* configurations. Both C_{16:1} and C_{18:1} are found in recent sediments and are often used to differentiate between isomers predominantly derived from bacteria (ω⁷) and algae (ω⁹) (Killops and Killops 2005). As described in Table II, other isomers were also detected and are usually associated with bacteria (ω¹⁰ isomers).

Hydroxy acids were obtained only from the labile acid fraction. Their biological formation is characterized by a high diversity of sources. For example, medium chain length (C₁₂ – C₁₈), iso- and

anteiso-branched β-hydroxy acids are produced by various bacteria, microalgae and seagrasses (Garcette-Lepecq et al. 2004 and references therein). The medium and long chain (C₁₆ – C₃₀) α-hydroxy acids can be related to microalgae and bacteria whereas ω-hydroxy acids, medium and long chain, are major components of cutins, suberins and higher plant epicuticular waxes (Garcette-Lepecq et al. 2004 and references therein).

The hydroxy acids identified in the four sediment samples are dominated by saturated compounds, with the highest percentage of ω-C₁₆ (37.0%) and ω-C₁₈ (38.9%) at stations ITA2 and ITA1, respectively (Table III). In spite of being reported in the literature as components of terrigenous plants, saturated ω-hydroxy acids were not identified in the mangrove leaf samples, with the exception of two hydroxy acids containing more than one hydroxyl group: ω-C₁₆ (9,10,16-trihydroxy acid) and ω-C₁₈ (9,10,16-trihydroxy acid); and one unsaturated ω-C_{18:1} hydroxy acid. The ω-C₁₆ 9,10,16-trihydroxy acid showed the highest percentage of all the hydroxy acids in two mangrove species (*Avicennia* and *Laguncularia*) and the Gramineae *Spartina* (see top mass chromatogram on Figure 3). The appearance of ω-hydroxy acids only in the sediments can be indicative of bacterial ω-oxidation of alkylchains, as previously reported in decomposing eelgrasses *Zostera marina* L. by de Leeuw et al. (1995). No ω-hydroxy acids were detected in the living eelgrass free and bound fractions investigated by the authors. For the α-hydroxy acid distribution no clear even-over-odd carbon number predominance was observed (see bottom mass chromatogram on Figure 3). The α-C₂₄ hydroxy acid was the major compound identified in the leaf samples, accounting for 37.7% in the leaves of the *Rhizophora mangle*. Previous studies on soil and surface sediments from Lake Ahmasjärvi in Finland have suggested that part of the α-hydroxy acids may originate from Gram-positive bacteria and other microbial groups (Keinänen et al. 2003).

TABLE II

Overview of the relative quantities of total *n*-alkanes, monocarboxylic saturated fatty acids, individual *iso* (*i*-C_x), *anteiso* (*ai*-C_x), unsaturated fatty acids and sterols in all sediment and leaf samples. CPI: Carbon Preference Index calculated in the C₂₂ – C₃₂ *n*-alkane range. Values shown are the average of duplicate analyses.

Compound	Stations (µg g ⁻¹ EOM) ^a				Mangrove Leaf (µg g ⁻¹ DL) ^b			
	ITA1	ITA2	ITA3	ITA4	Avic	Lagun	Spar	Rhiz
<i>ai</i> -C ₁₄	--	--	--	--	0.7	0.9	0.9	0.9
<i>ai</i> -C ₁₄ 5,9,13	1.1	0.6	43.8	--	--	--	--	--
<i>i</i> -C ₁₄	7.8	5.5	--	5.2	--	--	--	--
<i>ai</i> -C ₁₅	7.3	9.6	7.7	18.7	--	--	--	--
<i>i</i> -C ₁₅	3.5	2.2	6.7	2.3	--	--	--	--
<i>n</i> -C _{16:1} ω7- cis	1.7	--	--	--	--	--	--	--
<i>n</i> -C _{16:1} ω7- trans	0.8	2.3	--	3.4	--	--	--	--
<i>n</i> -C _{16:1} ω9 - cis	7.3	0.4	--	2.9	--	--	--	--
<i>ai</i> -C ₁₆	4.5	2.6	--	3.9	--	--	--	--
<i>i</i> -C ₁₆	--	--	5.1	--	--	--	--	--
<i>ai</i> -C ₁₇	5.1	1.8	--	1.7	--	--	--	--
<i>i</i> -C ₁₇	1.5	8.2	--	1.1	--	--	--	--
<i>n</i> -C _{18:3} - trans	--	--	--	--	1.7	206.5	22.0	6.3
<i>n</i> -C _{18:1} ω10	6.7	3.6	--	49.7	--	--	--	--
<i>n</i> -C _{18:1} ω15	8.0	4.6	5.9	31.1	--	--	--	--
<i>n</i> -C _{18:1} ω19-cis	1.3	0.4	--	1.3	--	--	--	--
<i>ai</i> -C ₁₈	6.7	5.7	--	4.7	--	--	--	--
<i>n</i> -C _{19:1} ω10	2.5	--	--	--	--	--	--	--
<i>n</i> -C _{19:1} ω10	2.1	0.9	--	--	--	--	--	--
<i>n</i> -C _{19:1} ω11	2.5	0.7	--	1.8	--	--	--	--
<i>n</i> -C _{22:1} ω13-cis	1.7	0.5	--	2.6	--	--	--	--
<i>n</i> -C _{22:1} ω13-trans	3.2	--	--	1.6	--	--	--	--
Σaliphatic ^d	29.5	27.9	34.6	27.5	38.8	26.9	8.8	24.6
CPI ^e	3.1	2.8	1.8	3.1	6.6	11.2	3.4	2.2
Σsat. acids ^f	87.3	107.3	40.9	315.8	126.9	330.3	524.5	394.8

^aµg g⁻¹ of extractable organic matter; ^bµg g⁻¹ of dry leaf; ^c(--) below detection limit

^daliphatic hydrocarbons: *n*-C₂₂-C₃₂+Pristane+Phytane; ^eCPI (Carbon Preference Index) = $2(nC_{23}+nC_{25}+nC_{27}+nC_{29}+nC_{31})/[(nC_{22}+nC_{24}+nC_{26}+nC_{28}+nC_{30})+(nC_{24}+nC_{26}+nC_{28}+nC_{30}+nC_{32})]$; ^fSaturated aliphatic acids: *n*-C₁₀-C₃₂.

These authors also showed that α-hydroxy acids with a chain length greater than 14 were generally more abundant in the sediment than in the soil. Two hydroxy acids, C₁₆ and C₂₄, appeared predominantly. Another study on surface sediments from the Danube delta has also shown that α-C₂₄ hydroxy acid was

the most abundant (Garcette-Lepecq et al. 2004). It is interesting to notice that these compounds are usually related to microbial sources and to biomass, but not specifically to leaves.

All sterols detected in the surface samples and leaf extracts are described below. Their

identifications were based on their mass spectra, relative retention times, comparison with authentic standards and key fragment ions of TMS derivatives: m/z 129 for C_{27} - C_{29} Δ^5 and $\Delta^{5,22}$ components; m/z 251 for C_{27} - C_{29} Δ^0 ; and m/z 257 for C_{27} - C_{29} Δ^{22} .

In the sediments, sterols are composed mainly of C_{27} and C_{29} components, which account for 33.2 % and 45.6 %, respectively. The C_{27} sterols identified (5β -cholestan- 3β -ol, cholest-5-en- 3β -ol, 5α -cholestan- 3β -ol, and one stanone: 5α -cholest-




Figure 3 – Partial mass chromatogram (TIC) of the extract (E4) obtained from the leaf extract of *Avicennia* (top). Partial mass chromatogram (TIC) and the main fragments employed to identify the α - and ω -hydroxy fatty acids as methyl esters from the extract (E4) of the ITA1 sediment (bottom).

TABLE III
Relative abundance (%) of hydroxy fatty acids obtained from the “acid fraction” via sequential treatment.

Compound	Stations				Mangrove Leaf			
	ITA1	ITA2	ITA3	ITA4	Avic	Lagun	Spar	Rhiz
α -OH C ₁₆	1.69	12.46	10.73	--*	--	--	--	--
α -OH C ₁₇	1.84	3.35	13.03	17.74	--	--	--	--
α -OH C ₁₈	2.46	3.76	0.42	4.74	--	--	--	--
α -OH C ₂₀	2.20	4.27	3.83	21.47	7.06	2.66	13.83	9.06
α -OH C ₂₂	2.25	5.69	--	8.18	0.21	--	--	--
α -OH C ₂₃	2.30	2.00	2.43	17.90	6.82	4.53	7.86	9.54
α -OH C ₂₄	8.62	5.09	7.71	7.03	12.36	8.09	17.76	37.73
α -OH C ₂₅	1.81	2.13	--	--	--	4.39	3.39	7.99
α -OH C ₂₆	0.15	0.80	--	14.60	0.64	1.45	4.90	4.22
α -OH C ₂₇	0.93	4.34	0.46	6.21	--	--	--	--
ω -OH C ₁₆	33.16	37.04	29.78	1.06	--	--	--	--
ω -OH C ₁₈	38.94	9.00	22.74	0.75	--	--	--	--
ω -OH C ₂₂	3.65	10.07	8.89	0.32	--	--	--	--
ω -OH 9,10,16 - C ₁₆	--	--	--	--	69.75	78.76	50.87	--
ω -OH 9,10,16 - C ₁₈	--	--	--	--	3.16	0.11	--	2.58
ω -OH C _{18:1}	--	--	--	--	--	--	--	28.87

* (--) Below detection limit

8-en-3-one) are usually associated with sewage, microbes or other marine sources since they are not commonly found in higher plant waxes (Grimalt et al. 1990). Compounds that are generally employed as fecal pollution indicators, coprostanol (5 β -cholestan-3 β -ol) and epicoprostanol (5 α -cholestan-3 β -ol), are the major components of C₂₇ sterols in most of the surface samples analyzed, with average values of 7.6 and 7.8 $\mu\text{g g}^{-1}$ EOM, respectively (Table IV). For both compounds, the highest concentrations were found at station ITA2 (11.1 $\mu\text{g g}^{-1}$ EOM and 12.0 $\mu\text{g g}^{-1}$ EOM for coprostanol and epicoprostanol, respectively). According to the literature, concentrations of coprostanol greater than 9 $\mu\text{g g}^{-1}$ are characteristic of sediments contaminated with sewage discharge (Nichols and Leeming 1991). However, it is important to mention that some authors have already emphasized that coprostanol can be

found in algae (Grimalt et al. 1990, Colombo et al. 1996) and also the reduction of Δ^5 sterols in sediments usually produces stanol mixtures where 5 α -epimers are dominant (Fernandes et al. 1999). Nonetheless, using the (5 β / 5 β + 5 α) stanol ratio proposed by Grimalt et al. (1990) for sewage pollution identification, we found ratios ranging from 0.30 to 0.58 (stations ITA4 and ITA1, respectively) for the sediments of Itacorubi (Table IV). Values for this ratio higher than 0.70 are considered to be indicative of significant sewage contamination (Grimalt et al. 1990, Carrera et al. 2001). In a previous study on the Itacorubi mangrove, we found values between 0.64 and 0.82 for river sediments receiving sewage discharges (Mater et al. 2004). These previous results suggest a higher level of contamination in the river sediments and, to a lesser extent, in the sediments of the mangrove located within the intertidal zone.

TABLE IV
Concentration and ratios of sterols.

Stations ($\mu\text{g g}^{-1}$ EOM) Compound	Mangrove Leaf ($\mu\text{g g}^{-1}$ DL)							
	ITA1	ITA2	ITA3	ITA4	Avic	Lagun	Spar	Rhiz
5 α -cholestan-3 β -ol	7.7	12.0	6.9	4.4	4.4	--	--	--
5 β -cholestan-3 β -ol	10.7	11.1	6.6	1.9	1.9	--	--	--
5-cholesten-3 β -ol	3.5	2.7	13.4	0.8	0.8	--	--	--
5 α -cholest-8-en-3-one	3.0	5.9	15.4	5.0	5.0	--	--	--
24-methyl-5,22-cholestadien-3 β -ol	26.0	15.7	10.7	12.0	12.0	--	--	2.1
24 α -ethyl-5,22-cholestadien-3 β -ol	4.1	6.7	10.7	1.0	1.0	5.1	5.8	5.0
24 β -ethyl-5,22-cholestadien-3 β -ol	20.2	7.4	9.6	4.5	4.5	5.4	1.0	4.5
24 β -ethyl-5 β -cholesten-3 β -ol	7.4	5.1	23.9	3.0	3.0	15.0	29.7	9.6
24 β -ethyl-5 β -cholestan-3 β -ol	11.3	16.6	2.2	5.5	5.5	--	--	--
5 β /(5 β + 5 α) ^a	0.58	0.48	0.49	0.30	0.30	--	--	--

^a ratio of cholestanols

The C₂₉ phytosterols were detected in higher abundance than the C₂₈ components. In fact, only 24-methylcholesta-5,22-dien-3 β -ol was determined, with very high concentration. A previous work with algae has shown that C₂₈ sterol can be found in diatoms (Barret et al. 1995). In the sediment samples, 24 β -ethylcholesta-5,22-dien-3 β -ol and 24 β -ethyl-5-cholesten-3 β -ol are the major C₂₉ compounds. These two compounds come mainly from terrigenous sources, but are not restricted to higher plants (Volkman 1986, Barret et al. 1995). Accordingly, in the extracts from the leaf samples only C₂₈ and C₂₉ were detected, with a strong predominance of 24 β -ethyl-5-cholesten-3 β -ol (β -sitosterol), varying from 9.6 to 33.5 $\mu\text{g g}^{-1}$ DL. It is interesting to note the presence of the stanol 24-ethyl-5 β -cholestan-3 β -ol in the sediment samples only. This compound has been previously detected in a marsh area of a mangrove ecosystem in northern Brazil (Koch et al. 2003). Stanols in sediments can originate from direct biogenetic inputs and possible mixtures of 5 α -stanols from marine organisms. Our results showed that no stanols were indentified in the leaf

samples, suggesting the occurrence of a direct input from diatoms as they are likely to be found in the flooding seawater (Soriano-Sierra and Ledo 1998), or a direct reduction from the C₂₉5-cholesten-3 β -ol by microbially-mediated diagenetic transformation reactions in the sediments (Takahashi et al. 1985, Emeis et al. 2000).

With regard to the triterpenols identified in the leaf extracts, seven compounds appear predominantly (Figure 4). Leaves of *A. schaueriana*, the dominant species in the mangrove, were characterized by high amounts of β -amyirin (olean-12-en-3 β -ol; 90.6 $\mu\text{g g}^{-1}$ DL) and lower contents of lupeol (lup-20(29)-en-3 β -ol), α -amyirin (ursen-12-en-3 β -ol), germanicol (olean-18-en-3 β -ol), betulin (lup-20(29)-en-3 β ,28 β -diol) and friedelin (friedelan-3-one). *Laguncularia racemosa*, the second most abundant adult species, was dominated by β -amyirin and lupeol, but with much lower amounts than those found in *A. schaueriana* (Figure 4). Among the triterpenes found in the sediments, β -amyirin showed the greatest decrease in the relative percentage found in the leaf extract. It seems to be the most labile molecule

among the triterpenols. However, a previous work with an incubation experiment (Koch et al. 2005) demonstrated that all leaf-derived triterpenols can be degraded. Working with sediment samples from a northern Brazilian mangrove, the authors reported that the relative biomarker signature in the sediment predominantly represented *R. mangle* leaf litter. In our study, a simple comparison between the relative abundance of seven triterpenols showed that betulin and α -amyrin are relatively higher in the sediments compared with the distribution in the leaf extracts. A very low amount of taraxerol ($1.9 \mu\text{g g}^{-1}$ DL) that has been used for identifying mangrove-derived organic matter in sediments (Koch et al. 2003, Jaffé et al. 2006, Volkman et al. 2007), was only found in the leaf extract of *Rhizophora mangle*. However, it is well preserved when compared to β -amyrin. Taraxerol can occur in many other higher plant species, and other sources may also contribute to its increase. However, the region is strongly dominated by mangrove species and other sources are unlikely. Interesting results were found for betulin, which was only detected in the leaf extract of *A. schaueriana* ($2.3 \mu\text{g g}^{-1}$ DL), but increased in the sediments. Betulin has been previously reported in sediments and is a good chemotaxonomic biomarker for mangrove species.




Figure 4 – Relative abundance of triterpenols in the sediments and leaf extracts (*Avicennia schaueriana*, *Laguncularia racemosa*, *Spartina alterniflora* and *Rhizophora mangle*).

The composition of lipids in the sediments and leaves showed that saturated fatty acids are the major compounds, accounting for over 35% of the total lipids identified (Figure 5). The leaves of *Rhizophora mangle* and *Spartina* are dominated by saturated fatty acids (mainly $n\text{C}_{16:0}$ and $n\text{C}_{18:0}$). A high percentage of saturated fatty acids (52%) was also detected in *L. racemosa*, but, in contrast to the other leaves, this species contains the highest percentage of unsaturated fatty acids ($n\text{C}_{18:3}$; 33%). The leaves of *A. schaueriana* showed the highest abundance of terpenes and sterols, comprising 33% and 17% of the total lipid concentration, respectively. Aliphatic hydrocarbons are the least abundant lipids and constitute less than 12% of the total lipids. Apart from saturated fatty acids, the sediments are dominated by sterols (20%), which showed an increase due to the presence of C_{27} compounds from other sources. Terpenes are the third most abundant components (13%), showing their importance as terrestrial biomarkers for mangrove-derived organic matter.




Figure 5 – Composition of lipids in the sediment samples (ss) and leaf extracts (sterols, triterpenols, unsaturated fatty acids, saturated fatty acids and aliphatic hydrocarbons).

CONCLUSIONS

The composition of lipids in the leaves revealed that sterols and triterpenols are two important groups of compounds that can be used to assess the contribution of the organic matter from mangrove leaves. Despite the changes in the relative abundance

from the leaf extract to the sediments, triterpenols, such as germanicol, α -amyrin, betulin and lupeol, are important biomarkers as they are well preserved in the sediments. In contrast to other terpenols, β -amirin was the most abundant triterpenol found in the plants, but it is rapidly degraded and less preserved in the sediments.

Saturated and unsaturated fatty acids cannot be exclusively correlated to mangrove input. In spite of being detected in the *L. racemosa* leaf, the unsaturated fatty acid ($nC_{18:3}$) was not found in the sediments. However the α -hydroxy acids (from α -OH C_{20} to C_{26}) found in leaves and sediments are good indicators of mangrove input.

ACKNOWLEDGMENTS

Scholarship for C. A. Silva was provided by Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES). Financial support was provided by Petrobras and Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq).

RESUMO

A distribuição relativa de vários compostos identificados em quatro amostras de sedimentos recentemente depositados no manguezal do Itacorubi, localizado na Ilha de Santa Catarina, no sul do Brasil, foi comparada a dos diversos compostos extraídos de folhas frescas de três espécies de mangues: *Avicennia schaueriana*, espécie dominante na região, *Rhizophora mangle*, *Laguncularia racemosa*, e a gramínea *Spartina alterniflora*. Terpenóis identificados previamente em espécies de mangues no norte do Brasil também foram encontrados. A espécie *A. schaueriana* contém principalmente β -amirina ($90,6 \mu\text{g g}^{-1}$ de material orgânico extraído); pequenas quantidades de friedelina, betulina e germanicol foram detectadas somente nas folhas dessa espécie. *R. mangle* também contém uma quantidade significativa de β -amirina e foi a única espécie na qual o taraxerol foi detectado. Contrariamente ao encontrado nas folhas, nas amostras de sedimentos predominaram o germanicol, α -amirina e campesterol. Apesar da maior labilidade, o composto

betulina também foi identificado. Duas séries homólogas de α e ω -hidroxi de ácidos graxos foram identificadas na fração ácida-alcalina. Apesar de serem relatados na literatura como componentes de plantas terrígenas, não foram identificados os ω -hidroxiácidos saturados. Nossos resultados indicam que embora os terpenóis possam ser usados como biomarcadores para material orgânico proveniente de manguezais, a distribuição relativa pode variar conforme a região.

Palavras-chave: biomarcadores, folha, manguezal, Santa Catarina, sedimento.

REFERENCES

- BALTZER F. 1982. La transition eau douce-eau salée dans les mangroves. Conséquences sédimentologiques et géochimiques. B Soc Geol Fr 144: 27-42.
- BARRET SM, VOLKMAN JK, DUNSTAN GA AND LEROI J-M. 1995. Sterols of 14 species of marine diatoms (Bacillariophyta). J Phycol 31: 360-369.
- BOUILLON S, DAHDOUN-GUEBAS F, RAO AVVS, KOEDAM N AND DEHAIRS F. 2003. Sources of organic carbon in mangrove sediments: variability and possible implications for ecosystem functioning. Hydrobiologia 495: 33-39.
- CARRERA R, WAGENER ALR, FILEMAN T AND READMAN JW. 2001. Distribuição de coprostanol (5β (H)-coletan-3 β -ol) em sedimentos superficiais da Baía de Guanabara: indicador da poluição recente por esgotos domésticos. Quim Nova 24: 37-42.
- COLOMBO JC, SILVERBERG N AND GEARING JN. 1996. Lipid biogeochemistry in the Laurentian Trough: I—fatty acids, sterols and aliphatic hydrocarbons in rapidly settling particles. Org Geochem 25: 211-225.
- DE LEEUW JW AND LARGEAU CA. 1993. A review of macromolecular organic compounds that comprise living organisms and their role in kerogen, coal and petroleum formation. In: Engel MH and Macko SA (Eds), Organic Geochemistry Principles and Applications, Plenum: New York, USA, p. 23-72.
- DE LEEUW JW, RIJSTRA WIC AND NIENHUIS PH. 1995. Free and bound fatty acids and hydroxyl fatty acids in the living and decomposing eelgrass *Zostera marina* L. Org Geochem 23: 721-728.
- DUTRA SJ. 1998. A bacia hidrográfica do Córrego Grande, Ilha de Santa Catarina, Brasil. In: Soriano-Sierra EJ and Ledo BS (Eds), Ecologia e Gerenciamento do Manguezal do Itacorubi, Florianópolis: NEMAR, CCB, UFSC, p. 31-35.
- EMEIS KC, STRUCK U, LEIPE T, POLLEHME F, KUNZENDORF H AND CHRISTIANSEN C. 2000. Changes in the C, N, P burial rates in some Baltic Sea sediments over the last 150 years—relevance to P regeneration rates and the phosphorus cycle. Mar Geol 167: 43-59.

- FARIAS CO, HAMACHER C, WAGENER ALR AND SCOFIELD AL. 2008. Origin and degradation of hydrocarbons in mangrove sediments (Rio de Janeiro, Brazil) contaminated by an oil spill. *Org Geochem* 39: 289-307.
- FERNANDES MB, SICRE MA, CARDOSO JN AND MACEDO SJ. 1999. Sedimentary 4-desmethyl sterols and *n*-alkanols in an eutrophic urban estuary, Capibaribe River, Brazil. *Sci Total Environ* 231: 1-16.
- GARCETTE-LEPEQC A, LARGEAU C, BOULOUBASSI I, DERENNE S, SALIOT A, LORRE A AND POINT V. 2004. Lipids and their modes of occurrence in two surface sediments from Danube delta and northwestern Black Sea: implications for sources and early diagenetic alteration I. Carboxylic acids. *Org Geochem* 35: 959-980.
- GOOSSENS H, DE LEEUW JW, RIJSTRA WIC, MEYBURG GJ AND SCHENCK PA. 1989a. Lipids and their mode of occurrence in a bacteria and sediments – I A methodological study of the lipids composition of *Acinetobacter calcoaceticus* LMD 79-41. *Org Geochem* 14: 15-25.
- GOOSSENS H, DÜREN RR, DE LEEUW JW AND SCHENCK PA. 1989b. Lipids and their mode of occurrence in a bacteria and sediments – II Lipids in the sediment of a stratified, freshwater lake. *Org Geochem* 14: 27-41.
- GRIMALT JO, FERNANDEZ P, BAYONA JM AND ALBAIGES J. 1990. Assessment of fecal sterols and ketones as indicator of urban sewage inputs to coastal waters. *Environ Sci Technol* 24: 357-363.
- HEFTMANN E. 1971. Functions of sterols in plants. *Lipids* 6: 128.
- HERNANDEZ ME, MEAD R, PERALBA MC AND JAFFÉ R. 2001. Origin and transport of *n*-alkane-2-ones in a subtropical estuary: potential biomarkers for seagrass-derived organic matter. *Org Geochem* 32: 21-32.
- JAFFÉ R, RUSHDI AI, MEDEIROS PM AND SIMONEIT BRT. 2006. Natural product biomarkers as indicators of sources and transport of sedimentary organic matter in a subtropical river. *Chemosphere* 64: 1870-1884.
- KEINÄNEN MM, KORHONEN LK, MARTIKAINEN PJ, VARTIAINEN T, MIETTINEN IT, LEHTOLA MJ, NENONEN K, PAJUNEN H AND KONTRO MH. 2003. Gas chromatographic – mass spectrometric detection of 2- and 3-hydroxy fatty acids as methyl esters from soil, sediment and biofilm. *J Chromatog B* 783: 443-451.
- KILLOPS S AND KILLOPS V. 2005. Introduction to Organic Geochemistry. Second Edition. Blackwell Publishing, UK.
- KOCH BP, HARDER J, LARA RJ AND KATTNER G. 2005. The effect of selective microbial degradation on the composition of mangrove derived pentacyclic triterpenols in surface sediments. *Org Geochem* 36: 273-285.
- KOCH BP, RULLKÖTTER J AND LARA RJ. 2003. Evaluation of triterpenols and sterols as organic matter biomarkers in a mangrove ecosystem in northern Brazil. *Wetl Ecol Manag* 11: 257-263.
- KOCH BP, SOUZA FILHO PWM, BEHLING H, COHEN MCL, KATTNER G, RULLKÖTTER J, SCHOLZ-BÖTTCHER B AND LARA RJ. 2011. Triterpenols in mangrove sediments as a proxy for organic matter derived from the red mangrove (*Rhizophora mangle*). *Org Geochem* 42: 62-73.
- LACERDA LD. 1998. Mangrove Ecosystems Occasional Papers No. 2. Vannucci M (Ed), ISME: Okinawa.
- LACERDA LD, ITTEKKOT V AND PATCHINEELAM SR. 1995. Biogeochemistry of mangrove soil organic matter: a comparison between *Rhizophora* and *Avicennia* soils in south-eastern Brazil. *Estuar Coast Shelf Sci* 40: 713-720.
- LAGO PF. 1997. Ilha de Santa Catarina: urbanização e manejo. Atas do Colóquio Franco-Brasileiro. In: Gestão de la zone littorale de l'île de Santa Catarina (Brésil).
- LALLIER-VERGÈS E, PERRUSSEL B, DISNAR JR AND BALTZER F. 1998. Relationships between environmental conditions and the diagenetic evolution of organic matter derived from higher plants in a modern mangrove swamp system (Guadeloupe, French West Indies). *Org Geochem* 29: 1663-1686.
- MADUREIRA LAS, VAN KREVELD SAV, EGLINTON G, CONTE MH, GANSEN G, HINTE JE AND OTTENS J. 1997. Late Quaternary High-Resolution biomarker and other sedimentary paleoclimate proxies in a Northeast Atlantic core. *Paleoceanography* 12: 255-269.
- MATER L, ALEXANDRE MR, HANSEL F AND MADUREIRA LAS. 2004. Assessment of lipid compounds and phosphorus in mangrove sediments of Santa Catarina Island, Brazil. *J Braz Chem Soc* 15: 725-734.
- NICHOLS PD AND LEEMING R. 1991. Tracing sewage in the marine environment. *Chem Australia*, July, p. 274-276.
- PEMPKOWIAK J, TYLMANN W, STANISZEWSKI A AND GOLEBIEWSKI R. 2006. Lignin depolymerization products as biomarkers of the organic matter sedimentary record in ²¹⁰Pb - ¹³⁷Cs-dated lake sediments. *Org Geochem* 37: 1452-1464.
- REZENDE CE, PFEIFFER WC, MARTINELLI LA, TSAMAKIS E, HEDGESII AND KEIL RG. 2010. Lignin phenols used to infer organic matter sources to Sepetiba Bay RJ, Brazil. *Estuar Coast Shelf Sci* 87: 479-486.
- RIELEY G, COLLIER RJ, JONES DM AND EGLINTON G. 1991. The biogeochemistry of Ellesmere Lake, U.K.-I: source correlation of leaf wax inputs to the sedimentary lipid record. *Org Geochem* 17: 901-912.
- RUIZ-FERNANDÉZ AC, HILLAIRES-MARCEL C, GHALEB B, SOTO-JIMÉNEZ M AND PAEZ-OSUNA F. 2002. Recent sedimentary history of anthropogenic impacts on the Culiacan River Estuary, northwestern Mexico: geochemical evidence from organic matter and nutrients. *Environ Pollut* 118: 365-377.
- SHEPARD JP. 1954. Nomenclature based on sand-silt-clay ratios. *J Sediment Petrol*, p. 151-158.
- SORIANO-SIERRA EJ AND LEDO BS. 1998. Ecologia e Gerenciamento do Manguezal do Itacorubi. Florianópolis: Série Fepema, Santa Catarina, 408 p.
- SUGUIO K. 1973. Introdução à sedimentologia. Universidade de São Paulo Press, São Paulo, 150 p.
- TAKAHASHI T, BROECKER WS AND LANGER S. 1985. Redfield ratio based on chemical-data from isopycnal surfaces. *J Geophys Res-Oceans* 90: 6907-6924.
- VENKATESAN MI AND KAPLAN IR. 1982. Distribution and transport of hydrocarbons in surface sediments of the Alaskan outer continental shelf. *Geochim Cosmochim Acta* 46: 2135-2149.

- VERSTEEGH GJM, BOSCH H-J AND DE LEEUW JW. 1997. Potential paleoenvironmental information of C₂₄ to C₃₆ mid-chain diols, keto-ols and mid-chain hydroxyl fatty acids; a critical review. *Org Geochem* 27: 1-13.
- VOGTS A, MOOSSEN H, ROMMERSKIRCHEN F AND RULLKÖTTER J. 2009. Distribution pattern and stable isotope composition of alkanes and alkan-1-ols from plant waxes of African rain forest and savanna C3 species. *Org Geochem* 40: 1037-1054.
- VOLKMAN JK. 1986. A review of sterol markers for marine and terrigenous organic matter. *Org Geochem* 9: 83-99.
- VOLKMAN JK, EGLINTON G AND CORNER EDS. 1980. Sterols and fatty acids of the marine diatom *biddulphia sinensis*. *Phytochemistry* 19: 1809-1813.
- VOLKMAN JK, REVILL AT, BONHAM PI AND CLEMENTSON LA. 2007. Sources of organic matter in sediments from the Ord River in tropical northern Australia. *Org Geochem* 38: 1039-1060.
- XU S, GAO X, LIU M AND CHEN Z. 2001. China's Yangtze estuary: II. Phosphorus and polycyclic aromatic hydrocarbons in tidal flat sediments. *Geomorphology* 41: 207-217.