



Anais da Academia Brasileira de Ciências

ISSN: 0001-3765

aabc@abc.org.br

Academia Brasileira de Ciências

Brasil

KLAUCK, VANDERLEI; PAZINATO, RAFAEL; LOPES, LEANDRO S.; CUCCO, DIEGO C.; DE LIMA, HORACIO L.; VOLPATO, ANDREIA; RADAPELLI, WILLIAN M.; STEFANI, LENITA C.M.; DA SILVA, ALEKSANDRO S.

Trichostrongylus and Haemonchus anthelmintic resistance in naturally infected sheep from southern Brazil

Anais da Academia Brasileira de Ciências, vol. 86, núm. 2, junio, 2014, pp. 777-784

Academia Brasileira de Ciências

Rio de Janeiro, Brasil

Available in: <http://www.redalyc.org/articulo.oa?id=32731288021>

- How to cite
- Complete issue
- More information about this article
- Journal's homepage in redalyc.org

redalyc.org

Scientific Information System

Network of Scientific Journals from Latin America, the Caribbean, Spain and Portugal

Non-profit academic project, developed under the open access initiative



***Trichostrongylus* and *Haemonchus* anthelmintic resistance in naturally infected sheep from southern Brazil**

VANDERLEI KLAUCK¹, RAFAEL PAZINATO¹, LEANDRO S. LOPES², DIEGO C. CUCCO²,
HORACIO L. DE LIMA¹, ANDREIA VOLPATO¹, WILLIAN M. RADAVELLI¹,
LENITA C.M. STEFANI^{2,3} and ALEKSANDRO S. DA SILVA²

¹Graduação de Zootecnia, Departamento de Zootecnia, Universidade do Estado de Santa Catarina (UDESC),
Rua Beloni Trombeta Zanin, 680E, Chapecó, 89815-630 Santa Catarina, SC, Brasil

²Departamento de Zootecnia, Universidade do Estado de Santa Catarina (UDESC),
Rua Beloni Trombeta Zanin, 680E, Chapecó, 89815-630 Santa Catarina, SC, Brasil

³Programa de Pós-Graduação em Ciência Animal, Universidade do Estado de Santa Catarina (UDESC),
Avenida Luiz de Camões, 2090, 88520-000 Lages, SC, Brasil

Manuscript received on February 19, 2013; accepted for publication on September 9, 2013

ABSTRACT

The anthelmintic resistance in small ruminants is a common problem and concern worldwide. The aim of this study was to verify anthelmintic treatment efficacy in naturally infected sheep. This study was conducted on nine herds that used the same anthelmintic management for over a year. In each farm, the animals were divided into two groups: untreated control group (n = 5) and treated (n = 10) according to the number of eggs per gram of feces (EPG). The treatment effect was checked based on EPG results and larval culture performed before treatment and 10 days after treatment. Significant differences were not observed ($P > 0.05$) on EPG results between untreated and treated groups. The coproculture showed that the animals were infected primarily by *Haemonchus* spp., *Trichostrongylus* spp., *Teladorsagia* spp., *Cooperia* spp. and *Oesophagostomum* spp. In all farms, anthelmintic resistance by genera *Haemonchus* and *Trichostrongylus* was found, but this resistance varied greatly between farms. *Haemonchus* spp. showed resistance to closantel, levamisole, and albendazole. *Trichostrongylus* spp. was shown to be resistant to closantel, levamisole, and albendazole. The drugs tested showed to be efficient against the genera *Teladorsagia*, *Cooperia*, and *Oesophagostomum*. Based on these results, we conclude that the anthelmintic resistance to the tested drugs is a problem present in the farms evaluated.

Key words: Anthelmintic resistance, albendazole, closantel, levamisole, helminths.

INTRODUCTION

Gastrointestinal parasites affect sheep production causing economic losses due to weight loss, decreased milk and wool production, and are the leading cause of death among infected untreated animals. Currently, Brazilian livestock has 17.3

million sheep, and the state of Santa Catarina has about 203.000 heads, mainly for meat production (IBGE 2010). Sheep raised under grazing conditions are more susceptible to various parasites and the primary control method is through the use of broad-spectrum anthelmintics (Waller 1997). However, the indiscriminate use of these drugs without appropriate association with other methods to fight parasite

Correspondence to: Aleksandro S. da Silva
E-mail: aleksandro_ss@yahoo.com.br

infection has favored the development of helminth resistance to various drugs in the gastrointestinal tract of small ruminants (Morales and Pino 2001).

The development of anthelmintic resistance has increased lately in small ruminants in the Southern Hemisphere, where sheep production takes place primarily in grazing systems (Waller 2003). The main sheep parasite in terms of prevalence and pathogenicity is *Haemonchus contortus*, but other species should be highlighted such as *Trichostrongylus colubriformis*. Mixed infections are commonly reported among these animals, mainly by *Cooperia* spp., *Teladorsagia* spp., *Oesophagostomum* spp. and *Strongyloides* spp. (Ramos et al. 2004). *H. contortus* is the most important nematode associated to the rapid development of resistance in small ruminants in southern Brazil (Farias et al. 1997), northeastern Brazil (Charles 1989), and in other countries in Latin America (Waller et al. 1996).

Anthelmintic resistance can be defined as a decrease in the efficiency of a drug against the parasite population that is generally susceptible to the drug (Sangster and Gill 1999). According to some researchers, various species of nematodes can develop resistance to a specific anthelmintic (multi-specific resistance) or a nematode species to various drugs (multidrug resistance) (Coles and Roush 1992). Resistant parasites are usually in a smaller number in population, but the selection pressure exerted by a given drug favors the increase of resistance among individuals in the same population originally susceptible to the anthelmintic treatment (Lejambre et al. 2000). Given the explanation above this study aims at verifying anthelmintic treatment efficacy in naturally infected sheep from Santa Catarina, southern Brazil.

MATERIALS AND METHODS

This study surveyed nine farms (farms A to I) located in the western part of the state of Santa Catarina, southern Brazil in a 30-day period, corresponding

to the month of May, 2012. First, an interview was conducted to find out about farm management regarding the use of anti-parasitic drugs as well as clinical cases of parasitism and animal mortality. Six of these farms with around 21 to 43 animals each, did not have sheep as their main source of income, but for family consumption only or for small sales purposes. Three other farms had a large herd (between 400 and 1000 sheep), of which two raised sheep for milk production, and one for meat. It was noted that all farms had other animal species; however, the sheep did not share the same pasture area. These farms were chosen because they were using the same drug for over a year, five of which were using the same drug for more than two years. Another requirement for the farm to be included in the study was that the sheep could not have received antiparasitic treatment in the last 30 days.

Among the tests used to detect anthelmintic resistance, the fecal egg count reduction test (FECRT) is the most widely employed. The FECRT is a practical and inexpensive test appropriate for all anthelmintic drugs and only requires technician's ability to count eggs per gram (EPG) of feces (Coles 2005, Verissimo et al. 2012). Fecal samples from 15 to 20 adult sheep were collected directly from the rectum, stored in identified plastic bags and transported under refrigeration in styrofoam boxes to the laboratory. Stool samples were processed within 48 hours of collection using the McMaster technique with supersaturated sugar solution in order to quantify the number of EPG (Gordon and Whitlock 1939), using 2 grams of feces from each animal. Sheep from each farm were divided into two groups: five animals were used as untreated control ($n = 5$) and ten animals that were treated with the same anthelmintic drug for more than one year. The groups were formed based on the results of EPG, aiming at homogeneous groups. A pool fecal sample from each group was also collected to perform coproculture under constant humidity and temperature (Roberts and O'Sullivan 1950).

After this initial analysis, the treatment was carried out with the anthelmintic dose recommended by the manufacturer, on each farm. Sheep from Farms A and B were treated with closantel (10mg kg^{-1}); animals from Farms C and D were treated with levamisole hydrochloride (5mg kg^{-1}); sheep from Farms E, F, G and H were treated with albendazole (4mg kg^{-1}); and Farm I used a combination of albendazole (4mg kg^{-1}) and levamisole (5mg kg^{-1}). All treatments were carried out orally in a single dose. After 10 days of treatment new samples were collected to perform EPG and larval culture as previously mentioned.

The data were tested for normality and transformed when necessary. EPG data were not normally distributed and were rank-transformed before statistical analysis. EPG mean \pm standard deviations were analyzed by Student's test (t test) among control and treated groups, and considered significant when $P < 0.05$. The anthelmintic efficiency in the FECRT was calculated by the RESO 2.0 program, where the mean arithmetic of EPG for the control group and the group treated by calculating the confidence interval of the mean was used. We considered the existence of anthelmintic resistance reduction when the EPG observed after treatment was less than 95% and the confidence interval below 90% as recommended by literature. In this study we did not aim to compare farms, since management, types of parasites and treatments were different.

RESULTS

Surveyed farms reported the same anthelmintic treatment for more than one year, treating animals at 1-4 month intervals, depending on the season and farm management. Only three farms that had sheep production as their main source of income had an anti-parasite program with treatments every 45 days during hot and rainy seasons. These farms also had reports of helminthiasis (pale mucous membranes and diarrhea) in young and adult animals and mortality in untreated young sheep.

The other farms treated their sheep only when clinical signs appeared, and these farms did not have mortality in adult sheep by nematodes, but there was lamb mortality in four of these farms between October and December 2011. All farms were using antiparasitic only in animals that showed clinical signs. None of the farms used their animals before treatment, and used the same dosage for all individuals, and only four farms were using rotational grazing. Due to the lack of nearby specialized laboratory assistance, parasitological stool examinations were not performed in the last year, even though they continued to use the same anthelmintic treatment as usual.

In the investigated farms a few different genera of helminths were identified, including *Haemonchus*, *Trichostrongylus*, *Teladorsagia*, *Cooperia* and *Oesophagostomum*, in which *Haemonchus* and *Trichostrongylus* were present in all farms and in greater prevalence (Table I). There was no significant difference ($P > 0.05$) in EPG counts between treated and untreated control groups in all farms tested. Parasitic treatments were unreliable according to highly variable EPG results from treated animals. FECRT values were below 95% in all farms (Figure 1), which characterizes anthelmintic resistance.

The main parasites that presented resistant to the anthelmintics tested were *Haemonchus* spp. and *Trichostrongylus* spp. as shown in Table I. *Trichostrongylus* resistance to closantel was observed in Farm B, and resistance to *Haemonchus* and *Trichostrongylus* was seen in Farm A. *Trichostrongylus* and *Haemonchus* resistance to levamisole hydrochloride was also observed in Farms C and D, respectively. All farms that used albendazole showed *Haemonchus* resistance (Farms E to H) as well as *Haemonchus* resistance associated with *Trichostrongylus* resistance in one farm (Farm F). A low efficiency against *Haemonchus* and anthelmintic resistance by *Trichostrongylus* was observed when the drugs albendazole and levamisole were used in combination (Farm I).

TABLE I

Average number of eggs per gram of feces (EPG) and percentage of helminthes (*Haemonchus* [HAE], *Trichostrongylus* [TRI], *Teladorsagia* [TEL], *Cooperia* [COO] and *Oesophagostomum* [OES]) present in fecal samples from sheep before and after anthelmintic treatment with closantel (Farms A and B), levamisole (Farms C and D), albendazole (Farms E, F, G and H), and combination of levamisole and albendazole (Farm I).

Farm	Treatment	Average of EPG/Group		Helminths per group (%)									
		Control	Treated	Control group					*Treated group				
				HAE	TRI	TEL	COO	OES	HAE	TRI	TEL	COO	OES
A	Before	1220±540	1310±835	58.0	38.0	0.0	3.0	1.0	54.0	42.0	0.0	2.0	2.0
	After	920±630	660±410	50.0	46.0	0.0	4.0	0.0	64.0	36.0	0.0	0.0	0.0
B	Before	850±245	760±335	10.0	71.0	19.0	0.0	0.0	6.0	80.0	14.0	0.0	0.0
	After	975±410	337±292	5.0	75.0	20.0	0.0	0.0	0.0	100.0	0.0	0.0	0.0
C	Before	2520±1045	2060±1235	34.0	66.0	0.0	0.0	0.0	28.0	72.0	0.0	0.0	0.0
	After	2725±2734	440±230	27.0	73.0	0.0	0.0	0.0	0.0	100.0	0.0	0.0	0.0
D	Before	870±385	950±504	89.0	11.0	0.0	0.0	0.0	94.0	6.0	0.0	0.0	0.0
	After	805±375	690±400	80.0	20.0	0.0	0.0	0.0	100.0	0.0	0.0	0.0	0.0
E	Before	210±165	268±220	53.0	47.0	0.0	0.0	0.0	40.0	60.0	0.0	0.0	0.0
	After	257±180	550±140	46.0	54.0	0.0	0.0	0.0	100.0	0.0	0.0	0.0	0.0
F	Before	1700±835	1550±1092	81.0	19.0	0.0	0.0	0.0	90.0	10.0	0.0	0.0	0.0
	After	1750±980	788±674	74.0	26.0	0.0	0.0	0.0	44.0	56.0	0.0	0.0	0.0
G	Before	200±100	325±283	60.0	30.0	0.0	1.0	9.0	68.0	20.0	0.0	4.0	8.0
	After	226±173	100±100	44.0	52.0	0.0	0.0	4.0	100.0	0.0	0.0	0.0	0.0
H	Before	440±267	452±405	50.0	42.0	8.0	0.0	0.0	54.0	44.0	2.0	0.0	0.0
	After	480±155	151±185	41.0	54.0	5.0	0.0	0.0	96.0	4.0	0.0	0.0	0.0
I	Before	8090±5750	7400±4840	67.0	28.0	5.0	0.0	0.0	72.0	25.0	2.0	1.0	0.0
	After	7705±6210	4205±3204	56.0	40.0	4.0	0.0	0.0	32.0	68.0	0.0	0.0	0.0

#There was no statistically significant difference ($P > 0.05$) in EPG between treated and untreated control group in each of the farms. *The sheep were treated with the dose recommended by the manufacturer, see label.

* In the last column "treated group" in **bold** presented resistant parasites (i.e. between 4-100%).




Figure 1 - The fecal egg count reduction test (FECRT) on farms in western Santa Catarina, when we tested the efficacy of anthelmintics to closantel (Farms A and B), levamisole (Farms C and D), albendazole (Farms E, F, G and H) and combination of levamisole and albendazole (Farm I). *FECRT less than 95% represents the occurrence of anthelmintic resistance.

DISCUSSION

Anthelmintic ineffectiveness is a global reality, where drugs that have no record of parasite resistance to helminths from the *Trichostrongylidae* family, are rare (Coles 2005, Monteiro 2010, Laurenson et al. 2012). In this study, we noted multidrug and multispecies resistance in sheep in the western part of the state of Santa Catarina, a region undiagnosed hitherto for helminthic resistance. In Brazil, the development of parasites is benefited by the tropical climate, which is conducive to the free-living stages of parasites such as eggs and larvae present in the environment (Monteiro 2010). In general, sheep helminths develop strong resistance to the anti-parasitic drugs commercially available. This resistance is a matter of evolution and adaptation of the parasite to survive. This process can occur in a short period of time or take a long time, depending primarily on the type of drug management. The everyday use, without technical guidance and incorrect anthelmintic treatment, is the main cause for the process of resistance, among other factors (Monteiro 2010, Elsheikha and Khan 2011). However, the control of nematode parasites cannot be viewed in isolation, because frequent feeding, age and physiological state are important factors which favor parasite control. Many authors argue that the main way to control helminthiasis is related to strategies designed to control the spread of parasites in the environment, but recent studies have shown that to attain this goal it is better to select resistant animals to parasites (Laurenson et al. 2012).

In this study, it was noted that most farms used the same anthelmintic drug on short intervals during a long period of time (1 to 4 years), without knowing its true effectiveness. All farms investigated in this study presented problems with resistance to one or more parasites, therefore it is a multispecies resistance with different degrees between farms varying accordingly to the type of drug used, similar to what was recently observed in the state of São Paulo, Brazil (Veríssimo et al. 2012).

In the study, we recorded the prevalence mainly by helminth from the *Trichostrongylidae* family, similar to that observed in other Brazilian states (Kawano et al. 2001, Ramos et al. 2004, Monteiro 2010, Veríssimo et al. 2012) and other countries (Elsheikha and Khan 2011, Papadopoulos et al. 2012). In the region studied, there was a predominance of parasites from the genus *Haemonchus* and *Trichostrongylus*, which were also parasites with diagnosis of resistance. In some farms, the genus *Teladorsagia* appeared prevalent, unlike the genera *Cooperia* and *Oesophagostomum*. This study found no resistance to the tested drugs by *Teladorsagia*, *Cooperia* and *Oesophagostomum*, different from what is reported in other Brazilian states (Sczesny-Moraes et al. 2010, Veríssimo et al. 2012).

The closantel belongs to the class of salicylanides that acts uncoupling oxidative phosphorylation in the parasite. The results observed for closantel should take into account that this anthelmintic is specifically targeted at *Haemonchus* sp. and it is not effective against many other helminths (Dash 1986). In this study, this drug was investigated in two farms, and in both farms drug resistance by *Trichostrongylus* sp. was detected, but only on those farms that had drug resistance by *Haemonchus* sp. However, closantel treatment for *Cooperia*, *Oesophagostomum* and *Teladorsagia* was effective in controlling these helminths, although this drug is not recommended to treat these worms (Dash 1986). Parasite resistance and/or low susceptibility of *Trichostrongylidae* family to closantel has been described in different continents, as well as Latin America (Nari et al. 1996, Ramos et al. 2002, Rosalinski-Moraes et al. 1992, Sczesny-Moraes et al. 2010, Veríssimo et al. 2012).

Another drug tested in our study was levamisole, an imidazothiazole, which is administered to control nematodes in ruminants (Gong et al. 2006). This anti-parasitic is widely used in Brazil for dairy cattle because the manufacturers claim that the product has no withdrawal period. However,

recent studies suggest that levamisole can leave residues in milk, even after the industrial process (Power et al. 2012). We mention this because one of the studied farms that used levamisole to control gastrointestinal nematodes in sheep is also a major milk and milk products producer. Two farms that used levamisole had nematode resistance, one of them by *Haemonchus* spp. and the other by *Trichostrongylus* spp., which is not a surprise (Nari et al. 1996, Ramos et al. 2002, Rosalinski-Moraes et al. 1992, Sczesny-Moraes et al. 2010, Veríssimo et al. 2012).

Albendazole, a broad-spectrum anti-parasitic drug, has been used in four farms, which acts by inhibiting the absorption of glucose by the parasite preventing the production of ATP (Elsheikha and Khan 2011). In all farms, anthelmintic resistance to the genera *Trichostrongylus* and *Haemonchus* was observed. The resistance of helminths in relation to albendazole has been registered for several years (Ramos et al. 2002, Molento 2004, Rosalinski-Moraes et al. 1992, Sczesny-Moraes et al. 2010, Veríssimo et al. 2012).

Due to individual resistance of some active principles, many industries have invested in the combination of drugs. Nevertheless, the outcome is not favorable as found in this study, where the combination of albendazole and levamisole had no effect on FECRT, and *Haemonchus* and *Trichostrongylus* resistance was observed. Multiple anthelmintic resistance in nematodes from small ruminants are widespread in some places, especially in tropical countries (Voigt et al. 2012). We believe that the use of drugs combined, uncontrollably and without orientation, can aggravate the problem of parasitic resistance.

According to some researchers, it is essential to understand the mechanisms involving drug resistance so that alternative treatment strategies can be developed (Waller 1997, James et al. 2009). Changes in genes or in gene expression in response to drugs, enable the organism to survive treatment

and might reflect evolution in a toxic environment in which drug resistance leads to 'survival of the fittest' (James et al. 2009). The drugs investigated in this study are often used and consequently presented a broad level of resistance by parasites from the *Trichostrongylidae* family. As already mentioned, the resistance is a difficult process to be avoided, being a matter of adaptation of the parasite and thus, surviving. However, errors related to the management and use of incorrect treatments can accelerate this process of resistance, which probably occurred in the analyzed farms which presented no differences between treated and untreated animals regarding FECRT results.

Based on our findings we conclude that farms in the western region of Santa Catarina have anthelmintic resistance to closantel, albendazole, and levamisole; i.e. the FECRT was less than 95% in all farms investigated. The anthelmintic resistance of the genera *Haemonchus* and *Trichostrongylus* was observed in the farms. In contrast, no resistance was observed to *Teladorsagia* spp., *Cooperia* spp. and *Oesophagostomum* spp. in this study.

ACKNOWLEDGMENTS

We thank Amanda B. Almeida, Julia Rossett, and Assis E.A. Bongoski from the Animal Science Department/UDESC for their assistance in collecting and processing all samples.

RESUMO

A resistência de anti-helmínticos em pequenos ruminantes é um problema comum e de interesse em todo o mundo. O objetivo deste estudo foi verificar a eficácia do tratamento anti-helmíntico em ovinos naturalmente infectados. Este estudo foi realizado em nove rebanhos que utilizaram o mesmo anti-helmíntico por mais de um ano. Em cada fazenda, os animais foram divididos em dois grupos: grupo controle, não tratado (n = 5) e tratados (n = 10) com base no número de ovos por grama nas fezes (OPG). O efeito do tratamento foi avaliado com base nos resultados de OPG e coprocultura

realizados antes do tratamento e 10 dias após. Não foi observada diferença significativa ($P > 0,05$) entre os grupos tratados e não tratados em relação aos resultados do OPG. A coprocultura demonstrou que os animais estavam infectados principalmente por *Haemonchus* spp., *Trichostrongylus* spp., *Teladorsagia* spp., *Cooperia* spp. e *Oesophagostomum* spp. Em todas as fazendas foi encontrada resistência anti-helmíntica dos helmintos *Haemonchus* spp. e *Trichostrongylus* spp., mas esta resistência variou muito entre fazendas. *Haemonchus* spp. demonstrou ser resistente ao closantel, levamisol e albendazol. *Trichostrongylus* spp. demonstrou ser resistente ao levamisole, closantel e albendazol. As drogas testadas mostraram-se eficientes contra *Teladorsagia* spp., *Cooperia* spp. e *Oesophagostomum* spp. Com base nesses resultados, podemos concluir que a resistência anti-helmíntica às drogas testadas é um problema presente nas fazendas avaliadas.

Palavras-chave: resistência anti-helmíntica, albendazol, closantel, levamisole, helmintos.

REFERENCES

- CHARLES TP. 1989. Seasonal prevalence of gastrointestinal nematodes of goats in Pernambuco State, Brazil. *Vet Parasitol* 30: 335-343.
- COLES GC. 2005. Anthelmintic resistance, looking to the future: A UK perspective. *Res Vet Sci* 78: 99-108.
- COLES GC AND ROUSH RT. 1992. Slowing the spread of anthelmintic resistant nematodes of sheep and goats in the United Kingdom. *Vet Rec* 130: 505-510.
- DASH KM. 1986. Control of helminthosis in lambs by strategic treatment with closantel and broad-spectrum anthelmintics. *Aust Vet J* 63: 47-48.
- ELSHEIKHA HM AND KHAN NA. 2011. Essentials of Veterinary Parasitology. Loughborough: Caister Academic Press, 221 p.
- FARIAS MT, BORDIN EL, FORBES AB AND NEWCOMB KA. 1997. A survey on resistance to anthelmintic in sheep studs farms of southern Brazil. *Vet Parasitol* 72: 209-214.
- GONG FC, WU DX, CAO Z AND HE XC. 2006. A fluorescence enhancement-based sensor using glycosylated metalloporphyrin as a recognition element for levamisole assay. *Bios Bioelect* 22: 423-428.
- GORDON HM AND WHITLOCK HV. 1939. A new technique for counting nematode eggs in sheep faeces. *J Sci Ind Res* 12: 50-52. A new technique for counting nematode eggs in sheep faeces.
- IBGE - INSTITUTO BRASILEIRO DE GEOGRAFIA E ESTATÍSTICA. 2010. Pesquisa Pecuária Municipal. Disponível em <<http://www.ibge.gov.br>> Accessed in July, 3rd, 2012.
- JAMES CE, HUDSON AL AND DAVEY MW. 2009. Drug resistance mechanisms in helminths: is it survival of the fittest? *Trends Parasitol* 25: 328-335.
- KAWANO EL, YAMAMURA MH AND RIBEIRO ELA. 2001. Efeito do tratamento com anti-helmíntico em cordeiros naturalmente infectados com helmintos gastrintestinais sobre os parâmetros hematológicos, ganho de peso e qualidade da carcaça. *Acta Vet Sci* 29: 113-121.
- LAURENSEN YCSM, KYRIAZAKIS I, FORBES AB AND BISHOP SC. 2012. Exploration of the epidemiological consequences of resistance to gastro-intestinal parasitism and grazing management of sheep through a mathematical model. *Vet Parasitol* 189: 238-249.
- LEJAMBRE L, GILL J, LENANE J AND BAKER P. 2000. Inheritance of ivermectin resistance in *Haemonchus contortus*. *Int J Parasitol* 30: 105-111.
- MOLENTO MB. 2004. Sustainable worm management. *Vet Rec* 171: 95-96.
- MONTEIRO SG. 2010. Parasitologia Veterinária. São Paulo: Roca, 356 p.
- MORALES G AND PINO LA. 2001. Drogas anti-helmínticas sobre estrongilos digestivos em ovinos estabulados. *Rev Cient Vet Trop* 26: 147-158.
- NARI A, SALLES J, GIL A, WALLER PJ AND HANSEN JW. 1996. The prevalence of anthelmintic resistance in nematode parasites of sheep in Southern Latin America: Uruguay. *Vet Parasitol* 62: 213-222.
- PAPADOPOULOS E, GALLIDIS E AND PTOCHOS S. 2012. Anthelmintic resistance in sheep in Europe: A selected review. *Vet Parasitol* 198: 85-88.
- POWER C, SAYERS R, O'BRIEN B, BLOEMHOFF Y, DANAHER M, FUREY A AND JORDAN K. 2012. Partitioning of nitroxylin, oxyclozanide and levamisole residues from milk to cream, skim milk and skim milk powder. *Int J Dairy Technol* 65: 1-4.
- RAMOS CI, BELLATO V, ÁVILA VS, COUTINHO GC AND SOUZA AP. 2002. Resistência de parasitos gastrintestinais de ovinos a alguns anti-helmínticos no Estado de Santa Catarina, Brasil. *Ciênc Rural* 32: 473-477.
- RAMOS CI, BELLATO V, SOUZA AP, ÁVILA VS, COUTINHO GC AND DALAGNOL CA. 2004. Epidemiologia das helmintoses gastrintestinais de ovinos no Planalto Catarinense. *Ciênc Rural* 34: 1889-1895.
- ROBERTS FHS AND O'SULLIVAN JP. 1950. Methods for egg counts and larval cultures for strongyles infesting the gastrointestinal tract of cattle. *Aust J Agric Res* 1: 99-102.
- ROSALINSKI-MORAES F, MORETTO LH, BRESOLIN WS, GABRIELLI I, KAER L, ZANCHET IK, COLES G AND ROUSH R. 1992. Slowing the spread of anthelmintic resistance nematodes of sheep and goats in the United Kingdom. *Vet Rec* 130: 505-510.
- SANGSTER NC AND GILL J. 1999. Pharmacology of anthelmintic resistance. *Parasitol Today* 15: 141-146.
- SCZESNY-MORAES EA, BIANCHIN I, DA SILVA KF, CATTO JBR AND PAIVA F. 2010. Anthelmintic resistance of gastrointestinal nematodes in sheep, Mato Grosso do Sul, Brazil. *Pesq Vet Bras* 30: 229-236.

- VERÍSSIMO CJ ET AL. 2012. Multidrug and multispecies resistance in sheep flocks from São Paulo State, Brazil. *Vet Parasitol* 187: 209-216.
- VOIGT AK, SCHEUERLEB M AND HAMELB D. 2012. Triple anthelmintic resistance in *Trichostrongylus* spp. in a German sheep flock. *Small Rumin Res* 106: 30-32.
- WALLER PJ. 1997. Anthelmintic Resistance. *Vet Parasitol* 72: 391-4127.
- WALLER PJ. 2003. Global perspectives on nematode parasite control in ruminant livestock: the need to adopt alternatives to chemotherapy, with emphasis on biological control. *Anim Health Res Rev* 4: 35-43.
- WALLER PJ, ECHEVARRIA F, EDDI C, MACIEL S, NARI A AND HANSEN JW. 1996. The prevalence of anthelmintic resistance in nematode parasites of sheep in Southern Latin America: General overview. *Vet Parasitol* 62: 181-187.