



Anais da Academia Brasileira de Ciências

ISSN: 0001-3765

aabc@abc.org.br

Academia Brasileira de Ciências

Brasil

LLIBRE, JAUME; NOVAES, DOUGLAS D.; TEIXEIRA, MARCO A.
Periodic solutions of Lienard differential equations via averaging theory of order two
Anais da Academia Brasileira de Ciências, vol. 87, núm. 4, octubre-diciembre, 2015, pp.
1905-1913
Academia Brasileira de Ciências
Rio de Janeiro, Brasil

Available in: <http://www.redalyc.org/articulo.oa?id=32743236002>

- How to cite
- Complete issue
- More information about this article
- Journal's homepage in redalyc.org

redalyc.org

Scientific Information System

Network of Scientific Journals from Latin America, the Caribbean, Spain and Portugal

Non-profit academic project, developed under the open access initiative



Periodic solutions of Lienard differential equations via averaging theory of order two

JAUME LLIBRE¹, DOUGLAS D. NOVAES² and MARCO A. TEIXEIRA²

¹Departament de Matemàtiques, Universitat Autònoma de Barcelona, 08193 Bellaterra, Barcelona, Catalonia, Spain

²Departamento de Matemática, Universidade Estadual de Campinas, Rua Sérgio Buarque de Holanda, 651, Cidade Universitária Zeferino Vaz, 13083-859 Campinas, SP, Brasil

Manuscript received on March 21, 2014; accepted for publication on February 20, 2015

ABSTRACT

For $\varepsilon \neq 0$ sufficiently small we provide sufficient conditions for the existence of periodic solutions for the Lienard differential equations of the form

$$x'' + f(x)x' + n^2x + g(x) = \varepsilon^2 p_1(t) + \varepsilon^3 p_2(t),$$

where n is a positive integer, $f : \mathbb{R} \rightarrow \mathbb{R}$ is a C^3 function, $g : \mathbb{R} \rightarrow \mathbb{R}$ is a C^4 function, and $p_i : \mathbb{R} \rightarrow \mathbb{R}$ for $i = 1, 2$ are continuous 2π -periodic function. The main tool used in this paper is the averaging theory of second order. We also provide one application of the main result obtained.

Key words: periodic solution, Lienard differential equation, averaging theory, bifurcation theory.

1 - INTRODUCTION AND STATEMENT OF THE MAIN RESULTS

In Ma and Wang (2013) it was studied the existence of periodic solutions for the class of Lienard differential equations of the form

$$x'' + f(x)x' + V'(x) + g(x) = p(t), \quad (1)$$

where $f, V, g, p : \mathbb{R} \rightarrow \mathbb{R}$ are continuous and p is 2π -periodic. They also assumed that g and V' are locally Lipschitz; and the function V is a $2\pi/n$ -isochronous potential, i.e. all nontrivial solutions of $x'' + V'(x) = 0$ are $2\pi/n$ -periodic, where n is a positive integer. The authors provided sufficient bounded conditions related with the functions involved in equation (1) to ensure the existence of periodic solutions for this equation. We shall study a particular subclass of equations (1) in which such bounded conditions are not necessary.

In this paper we consider the subclass

$$x'' + f(x)x' + n^2x + g(x) = \varepsilon^2 p_1(t) + \varepsilon^3 p_2(t), \quad (2)$$

of Lienard differential equations (1) where n is a positive integer, ε is a small parameter, $f : \mathbb{R} \rightarrow \mathbb{R}$ is a \mathcal{C}^3 function in a neighborhood of $x = 0$, $g : \mathbb{R} \rightarrow \mathbb{R}$ is a \mathcal{C}^4 function in a neighborhood of $x = 0$, and $p_i : \mathbb{R} \rightarrow \mathbb{R}$ for $i = 1, 2$ are continuous 2π -periodic functions. Note that here we are taking $V(x) = n^2 x^2 / 2$ which is a $2\pi/n$ -isochronous potential, already considered in Ma and Wang (2013).

The objective of this paper is to give sufficient conditions on the functions f , g and p_i to assure the existence of periodic solutions for the equation (2). Here the functions f and g do not need to satisfy the bounded conditions of Ma and Wang (2013).

In general, to obtain analytically periodic solutions of a differential system, is a very difficult problem, many times impossible. Here using the averaging theory this difficult problem for the differential equations (2) is reduced to find the zeros of a nonlinear system of two functions with two unknowns. We must mention that the averaging theory for finding periodic solutions, in general, does not provide all the periodic solutions of the system. More precisely, the main tool used will be the averaging theory of second order for computing periodic orbits, see Llibre (2002), Buică and Llibre (2004), Giné et al. (2013), Llibre et al. (2014). This theory provides a quantitative relation between the solutions of some non-autonomous periodic differential system and the solutions of the averaged differential system, which is autonomous. In this way a finite dimensional function f is computed, the simple zeros of this function correspond with the periodic orbits of the non-autonomous periodic differential system for values of a parameter $\varepsilon \neq 0$ sufficiently small. Here a simple zero a of a function f means that the Jacobian of f at a is not zero. For a general introduction to the averaging theory see, for instance, the book Sanders et al. (2007).

In order to present our results we need some preliminary definitions and notations. We define the constants

$$\alpha = \int_0^{2\pi} p_1(t) \sin(nt) dt \quad \text{and} \quad \beta = \int_0^{2\pi} p_1(t) \cos(nt) dt.$$

and the two functions

$$\begin{aligned} f_{21}(u, v) = & -\frac{n^2 u^2 + v^2}{24n^6} \left(3n^2(a_1 b_1 - a_2 n^2)u + (10b_1^2 + n^2(a_1^2 - 9b_2))v \right) \\ & - \frac{1}{2\pi n^3} \int_0^{2\pi} \sin(nt) \left(\left(\int_0^t p_1(s) \cos(ns) ds \right) \right. \\ & \cdot (-b_1 v + (n^2 a_1 u + b_1 v) \cos(2nt) + n(a_1 v - b_1 u) \sin(2nt)) \\ & + n \left(n p_2(t) + \left(\int_0^t \frac{p_1(s) \sin(ns)}{n} ds \right) \right. \\ & \left. \left. \cdot (n b_1 u + n(b_1 u - a_1 v) \cos(2nt) + (n^2 a_1 u + b_1 v) \sin(2nt)) \right) \right) dt, \end{aligned}$$

and

$$\begin{aligned} f_{22}(u, v) = & \frac{n^2 u^2 + v^2}{24n^4} \left((10b_1^2 + n^2(a_1^2 - 9b_2))u + (3n^2 a_2 - 3a_1 b_1)v \right) \\ & + \frac{1}{2\pi n^2} \int_0^{2\pi} \left(n^2 p_2(t) \cos(nt) + \cos(nt) \left(\int_0^t \frac{p_1(s) \sin(ns)}{n} ds \right) \right. \\ & \cdot n(b_1 u + n(b_1 u - a_1 v) \cos(2nt) + (n^2 a_1 u + b_1 v) \sin(2nt)) \\ & + \cos(nt) \left(\int_0^t p_1(s) \cos(ns) ds \right) \\ & \cdot (-b_1 v + (n^2 a_1 u + b_1 v) \cos(2nt) + n(a_1 v - b_1 u) \sin(2nt)) \Big) dt, \end{aligned}$$

where

$$a_1 = f'(0), \quad a_2 = \frac{1}{2}f''(0), \quad b_1 = \frac{1}{2}g''(0), \quad \text{and} \quad b_2 = \frac{1}{6}g'''(0).$$

Our main result is the following.

Theorem 1. Assume that the functions f and g of the Lienard differential equation (2) satisfy that f is a \mathcal{C}^3 function in a neighborhood of $x = 0$, g is a \mathcal{C}^4 function in a neighborhood of $x = 0$, and $f(0) = g(0) = g'(0) = 0$. Suppose also that the constants $\alpha = \beta = 0$. Then for $\varepsilon \neq 0$ sufficiently small and for every simple zero (u^*, v^*) of the system

$$f_{21}(u, v) = 0, \quad f_{22}(u, v) = 0,$$

there exists a periodic solution $x(t, \varepsilon)$ of the differential equation (2) such that $x(0, \varepsilon) \approx \varepsilon u^* + \mathcal{O}(\varepsilon^2)$ and $x'(0, \varepsilon) \approx \varepsilon v^* + \mathcal{O}(\varepsilon^2)$.

Theorem 1 is proved in section 2.

In the next corollary we apply Theorem 1 to a given Lienard differential equation (2) and we show that such an equation has two periodic solutions.

Corollary 2. Assume that

$$f(x) = a_1 x + a_2 x^2, \quad g(x) = b_1 x^2 + b_2 x^3, \quad p_1(t) = 1, \quad \text{and} \quad p_2(t) = \sin(nt),$$

where

$$\begin{aligned} a_1 &= \frac{-7n^5 + 2n^3 + n}{-6n^4 - 4n^2 + 2}, \\ a_2 &= \frac{n(7n^7 - 6n^6 - 2n^5 + 2n^4 - n^3 + 6n^2 - 2)}{(n^2 + 1)^2(3n^2 - 1)^2}, \\ b_1 &= \frac{2n^5}{3n^4 + 2n^2 - 1}, \quad \text{and} \\ b_2 &= \frac{n^2(49n^8 + 132n^6 - 144n^5 - 10n^4 - 96n^3 + 4n^2 + 48n + 1)}{36(n^2 + 1)^2(3n^2 - 1)^2}. \end{aligned}$$

Then, for $\varepsilon \neq 0$ sufficiently small the Lienard differential equation (2) has two periodic solutions $x_i(t, \varepsilon)$ for $i = 1, 2$ such that

$$x_1(0, \varepsilon) \approx \varepsilon + \mathcal{O}(\varepsilon^2) \quad \text{and} \quad x'_1(0, \varepsilon) \approx \varepsilon + \mathcal{O}(\varepsilon^2),$$

$$x_2(0, \varepsilon) \approx 2\varepsilon + \mathcal{O}(\varepsilon^2) \quad \text{and} \quad x'_2(0, \varepsilon) \approx \mathcal{O}(\varepsilon^2).$$

Corollary 2 is proved in section 2.

In Appendix we summarize the averaging theory of second order, for studying periodic solutions that we shall need for proving Theorem 1.

We remark that in Ma and Wang (2013) the results about periodic solutions of system (1) are given, among other hypotheses, under the assumption that the function

$$F(x) = \int_0^x f(s) ds$$

is bounded on \mathbb{R} or that

$$-\infty < \liminf_{|x| \rightarrow +\infty} \frac{F(x)}{|x|^{p-2}x} \leq \lim_{|x| \rightarrow +\infty} \frac{F(x)}{|x|^{p-2}x} < +\infty,$$

for some $1 \leq p < 2$. It is worth emphasizing that Theorem 1 assures the existence of periodic solutions of system (2) without making any bounding assumption. Consequently, the present result detects some periodic solutions which are not detected by the theorems given in Ma and Wang (2013). Indeed, concerning the Corollary 2, $F(x) = a_1 x^2/2 + a_2 x^3/3$. It is easy to see that the above conditions do not hold in this case.

2 - PROOF OF THE RESULTS

In this section we shall prove Theorem 1 and Corollary 2.

Proof of Theorem 1. First we shall write the Lienard differential equations (2) in the normal form for applying the averaging theory, see Theorem 3 of the Appendix.

We change the variable x by a new variable z doing the rescaling $x = \varepsilon z$. Then equation (3) becomes

$$z'' + f(\varepsilon z)z' + n^2 z + \frac{g(\varepsilon z)}{\varepsilon} = \varepsilon p_1(t) + \varepsilon^2 p_2(t). \quad (3)$$

Since f is a \mathcal{C}^3 function in a neighborhood of $x = 0$, g is a \mathcal{C}^4 function in a neighborhood of $x = 0$, and $f(0) = g(0) = g'(0) = 0$, we can write

$$f(\varepsilon z) = \varepsilon f'(0)z + \varepsilon^2 \frac{1}{2} f''(0)z^2 + \mathcal{O}(\varepsilon^3) = \varepsilon a_1 z + \varepsilon^2 a_2 z^2 + \mathcal{O}(\varepsilon^3),$$

$$g(\varepsilon z) = \varepsilon^2 \frac{1}{2} g''(0)z^2 + \varepsilon^3 \frac{1}{6} g'''(0)z^3 + \mathcal{O}(\varepsilon^4) = \varepsilon^2 b_1 z^2 + \varepsilon^3 b_2 z^3 + \mathcal{O}(\varepsilon^4).$$

Thus

$$f(\varepsilon z)z' + \frac{g(\varepsilon z)}{\varepsilon} = \varepsilon(a_1 z z' + b_1 z^2) + \varepsilon^2(a_2 z^2 z' + b_2 z^3) + \mathcal{O}(\varepsilon^3).$$

We introduce a new variable $w = z'$. Then the differential equation (3) can be written as the differential system

$$\begin{aligned} z' &= w, \\ w' &= -n^2 z + \varepsilon(p_1(t) - a_1 z w - b_1 z^2) + \varepsilon^2(p_2(t) - a_2 z^2 w - b_2 z^3) \\ &\quad + \mathcal{O}(\varepsilon^3). \end{aligned} \quad (4)$$

Now we change the variables (z, w) by the new variables (u, v) defined through the equality

$$\begin{pmatrix} z \\ w \end{pmatrix} = \begin{pmatrix} \cos(nt) & \frac{\sin(nt)}{n} \\ -n \sin(nt) & \cos(nt) \end{pmatrix} \begin{pmatrix} u \\ v \end{pmatrix}. \quad (5)$$

We do these changes so that the differential system in the new variables (u, v) starts with terms of order $\mathcal{O}(\varepsilon)$ and we can apply the averaging theory described in the Appendix. Thus the differential system (4) in the new variables becomes

$$\begin{aligned} u' &= \varepsilon \frac{1}{n^3} G_1 \sin(nt) + \varepsilon^2 \frac{1}{n^4} G_2 \sin(nt) + \mathcal{O}(\varepsilon^3) \\ &= \varepsilon F_{11}(t, u, v) + \varepsilon^2 F_{21}(t, u, v) + \mathcal{O}(\varepsilon^3), \\ v' &= \varepsilon \frac{1}{n^3} G_1 \cos(nt) + \varepsilon^2 \frac{1}{n^4} G_2 \cos(nt) + \mathcal{O}(\varepsilon^3) \\ &= \varepsilon F_{12}(t, u, v) + \varepsilon^2 F_{22}(t, u, v) + \mathcal{O}(\varepsilon^3), \end{aligned} \quad (6)$$

where

$$\begin{aligned} G_1 &= -n^2 p_1(t) + A(B_1 \cos(nt) + C_1 \sin(nt)), \\ G_2 &= -n^3 p_2(t) + A^2(B_2 \cos(nt) + C_2 \sin(nt)), \\ A &= nu \cos(nt) + v \sin(nt), \\ B_i &= n(b_i u + a_i v), \\ C_i &= -n^2 a_i u + b_i v, \end{aligned}$$

In short, the differential system (6) is in the normal form for applying the averaging theory described in Theorem 3 of the Appendix. Using the notation of the Appendix we have $T = 2\pi$.

Let $F_i(t, u, v) = (F_{i1}(t, u, v), F_{i2}(t, u, v))$ for $i = 1, 2$. Now we compute the function $f_1(u, v)$ defined in the Appendix and we get, from the assumptions, that

$$\begin{aligned} f_1(u, v) &= \frac{1}{2\pi} \left(\int_0^{2\pi} F_{11}(t, u, v) dt, \int_0^{2\pi} F_{12}(t, u, v) dt \right) \\ &= \frac{1}{2\pi} \left(-\frac{\alpha}{n}, \beta \right) = (0, 0). \end{aligned}$$

Since the function $f_1(u, v) = (0, 0)$ we shall apply the averaging theory of second order. So we first compute

$$\begin{aligned} \int_0^t F_{11}(s, u, v) ds &= - \int_0^t \frac{p_1(s) \sin(ns)}{n} ds \\ &\quad - \frac{3}{12n^4} \left((b_1 u^2 n^2 - 2a_1 u v n^2 + 3b_1 v^2) \cos(nt) \right. \\ &\quad \left. + (b_1 u^2 n^2 + 2a_1 u v n^2 - b_1 v^2) \cos(3nt) \right. \\ &\quad \left. - 4((-a_1 u^2 n^3 + a_1 v^2 n + 2b_1 u v n) \sin^3(nt) \right. \\ &\quad \left. + b_1 n^2 u^2 + 2b_1 v^2 - a_1 n^2 u v) \right), \\ \int_0^t F_{21}(s, u, v) ds &= \int_0^t p_1(s) \cos(ns) ds \\ &\quad + \frac{1}{12n^3} \left(-n(a_1 n^2 u^2 - 2b_1 v u - a_1 v^2) \right. \\ &\quad \cdot (3 \cos(nt) + \cos(3nt) - 4) \\ &\quad - 3(2a_1 u v n^2 + b_1(3n^2 u^2 + v^2)) \sin(nt) \\ &\quad \left. + (b_1(v^2 - n^2 u^2) - 2a_1 n^2 u v) \sin(3nt) \right), \\ D_{(u,v)}(F_1(t, u, v)) &= \begin{pmatrix} \frac{\partial F_{11}(t, u, v)}{\partial u} & \frac{\partial F_{11}(t, u, v)}{\partial v} \\ \frac{\partial F_{21}(t, u, v)}{\partial u} & \frac{\partial F_{21}(t, u, v)}{\partial v} \end{pmatrix} \end{aligned}$$

Now we are ready to compute the function

$$f_2(u, v) = (f_{21}(u, v), f_{22}(u, v)), \quad (7)$$

defined in the Appendix, and we get the functions $f_{21}(u, v)$ and $f_{22}(u, v)$ which appear in the statement of the theorem.

Now, from Theorem 3, we obtain that for every $\varepsilon \neq 0$ sufficiently small and for every simple zero (u^*, v^*) of the system (7), i.e. satisfying that

$$\det \begin{pmatrix} \frac{\partial f_{21}(u, v)}{\partial u} & \frac{\partial f_{21}(u, v)}{\partial v} \\ \frac{\partial f_{22}(u, v)}{\partial u} & \frac{\partial f_{22}(u, v)}{\partial v} \end{pmatrix} \bigg|_{(u, v) = (u^*, v^*)} \neq 0,$$

there exists a periodic solution $(u(t, \varepsilon), v(t, \varepsilon))$ of the differential system (6) such that

$$(u(0, \varepsilon), v(0, \varepsilon)) \rightarrow (u^*, v^*) \quad \text{when } \varepsilon \rightarrow 0.$$

Going back through the change of variables (5) the periodic solution $(u(t, \varepsilon), v(t, \varepsilon))$ of the differential system (6) becomes the periodic solution

$$\begin{aligned} z(t, \varepsilon) &= \cos(nt)u(t, \varepsilon) + \frac{1}{n} \sin(nt)v(t, \varepsilon), \\ w(t, \varepsilon) &= -n \sin(nt)u(t, \varepsilon) + \cos(nt)v(t, \varepsilon), \end{aligned}$$

of the differential system (4) such that

$$(z(0, \varepsilon), w(0, \varepsilon)) \rightarrow (u^*, v^*) \quad \text{when } \varepsilon \rightarrow 0.$$

Finally, since $x = \varepsilon z$ the periodic solution $(z(t, \varepsilon), w(t, \varepsilon))$ of the differential system (4) provides the periodic solution

$$x(t, \varepsilon) = \varepsilon \left(\cos(nt)u(t, \varepsilon) + \frac{1}{n} \sin(nt)v(t, \varepsilon) \right),$$

of the Lienard differential equation (2) such that

$$x(0, \varepsilon) \approx \varepsilon u^* + \mathcal{O}(\varepsilon^2).$$

Moreover, since $x' = \varepsilon w$ the theorem follows. \square

Proof of Corollary 2. We shall apply the results of Theorem 1 to the Lienard differential equation of the statement of Corollary 2. So we compute the functions $f_{21}(u, v)$ and $f_{22}(u, v)$ defined just before the statement of Theorem 1, and we obtain

$$\begin{aligned} & \frac{-(u^3 - 7u + 6)n^4 + (u^3 - 2vu^2 - (v^2 + 2)u + 8v - 4)n^2 - 2v^3 + u(v^2 - 1) + 2}{4n(3n^4 + 2n^2 - 1)}, \\ & \frac{(7v + u(2u^2 - vu - 8))n^4 + v(u^2 + 2vu - v^2 - 2)n^2 + v^3 - v}{4n(3n^4 + 2n^2 - 1)}, \end{aligned}$$

respectively.

Doing the resultant of the functions f_{21} and f_{22} with respect to the variable v we obtain a cubic polynomial in the variable u which has the following three roots

$$u_1 = 1, \quad u_2 = 2, \quad u_3 = \frac{-3n^6 + 31n^4 - 25n^2 + 5}{(n^2 + 1)^3}.$$

In a similar way doing the resultant of the functions f_{21} and f_{22} with respect to the variable u we obtain another cubic polynomial in the variable v which has the following three roots

$$v_1 = 1, \quad v_2 = 0, \quad v_3 = \frac{-15n^6 + 35n^4 - 13n^2 + 1}{(n^2 + 1)^3}.$$

From the properties of the resultants it follows that all the solutions (u^*, v^*) of the system $f_{21}(u, v) = 0$, $f_{22}(u, v) = 0$ are of the form (u_i, v_j) being u_i and v_j some of the above roots. Trying the nine possible solutions, we obtain only two solutions for the system $f_{21}(u, v) = 0$, $f_{22}(u, v) = 0$, namely

$$(u_1^*, v_1^*) = (1, 1) \quad \text{and} \quad (u_2^*, v_2^*) = (2, 0).$$

Then applying Theorem 1 the corollary follows. □

For more information about the resultants see, for instance, Oliver (1999).

ACKNOWLEDGMENTS

The first author is partially supported by Ministerio de Economía y Competitividad (MiNECO) grant number MTM2013-40998-P, by Agência de Gestió d'Ajuts Universitaris i de Recerca (AGAUR) grant number 2013SGR-568, and by Marie Curie International Research Staff Exchange Scheme (IRSES) grants numbers FP7-PEOPLE-2012-IRSES 318999 and 316338. The second author is partially supported by Fundação de Amparo à Pesquisa do Estado de São Paulo (FAPESP) grant number 2015/02517-6. The third author is partially supported by FAPESP grant number 2012/18780-0. The three authors are also supported by Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES) grant number 88881.030454/2013-01 from the program CSF-PVE. We thank to the referees for their helpful comments and suggestions.

RESUMO

Para $\varepsilon \neq 0$ suficientemente pequeno encontramos condições suficientes que garantem a existência de soluções periódicas para sistemas diferenciais de Lienard da forma

$$x'' + f(x)x' + n^2x + g(x) = \varepsilon^2 p_1(t) + \varepsilon^3 p_2(t),$$

onde n é um inteiro positivo, $f : \mathbb{R} \rightarrow \mathbb{R}$ é uma função C^3 , $g : \mathbb{R} \rightarrow \mathbb{R}$ é uma função C^4 , e $p_i : \mathbb{R} \rightarrow \mathbb{R}$ para $i = 1, 2$ são funções contínuas 2π -periódicas. A principal ferramenta usada neste artigo é a teoria "averaging" de segunda ordem. Uma aplicação do resultado principal é feita.

Palavras-chave: solução periódica, equação diferencial de Lienard, teoria da média, teoria de bifurcação.

REFERENCES

- BUICĂ A AND LLIBRE J. 2004. Averaging methods for finding periodic orbits via brouwer degree. *Bull Sci Math* 128: 7-22.
- GINÉ J, GRAU M AND LLIBRE J. 2013. Averaging theory at any order for computing periodic orbits. *Phys D* 250: 58-65.
- LLIBRE J. 2002. Averaging theory and limit cycles for quadratic systems. *Radovi Matematički* 11: 215-228.
- LLIBRE J, NOVAES DD AND TEIXEIRA MA. 2014. Higher order averaging theory for finding periodic solutions via brouwer degree. *Nonlinearity* 27: 563-583.
- MA T AND WANG Z. 2013. Periodic solutions of lienard equations with resonant isochronous potentials. *Disc Cont Dyn Sys* 33: 1563-1581.

OLIVER P. 1999. Classical invariant theory, volume 44 of London Math. Soc. Student Texts Cambridge Univ. Press, New York, 2nd edition.

SANDERS JA, VERHULST F AND MURDOCK J. 2007. Averaging methods in nonlinear dynamical systems, volume 59 of Applied Mathematical Sciences Springer, New York, 2nd edition.

APPENDIX THE AVERAGING THEORY OF SECOND ORDER

In this section we recall the averaging theory of second order to find periodic orbits.

Theorem 3. *Consider the differential system*

$$\dot{x}(t) = \varepsilon F_1(t, x) + \varepsilon^2 F_2(t, x) + \varepsilon^3 R(t, x, \varepsilon), \quad (8)$$

where $F_1, F_2 : \mathbb{R} \times D \rightarrow \mathbb{R}^n$, $R : \mathbb{R} \times D \times (-\varepsilon_f, \varepsilon_f) \rightarrow \mathbb{R}^n$ are continuous functions, T -periodic in the first variable, and D is an open subset of \mathbb{R}^n . Assume that the following hypothesis (i) and (ii) hold.

- (i) $F_1(t, \cdot) \in C^1(D)$ for all $t \in \mathbb{R}$, F_1, F_2, R and $D_x F_1$ are locally Lipschitz with respect to x . We define $f_1, f_2 : D \rightarrow \mathbb{R}^n$ as

$$f_1(z) = \frac{1}{2\pi} \int_0^T F_1(s, z) ds,$$

$$f_2(z) = \frac{1}{2\pi} \int_0^T [D_z F_1(s, z) \int_0^s F_1(t, z) dt + F_2(s, z)] ds.$$

- (ii) For $V \subset D$ an open and bounded set and for each $\varepsilon \in (-\varepsilon_f, \varepsilon_f) \setminus \{0\}$, there exist $a \in V$ such that $f_1(a) + \varepsilon f_2(a) = 0$ and $d_B(f_1 + \varepsilon f_2, V, 0) \neq 0$ (see its definition later on).

Then for $|\varepsilon| > 0$ sufficiently small, there exists a T -periodic solution $\varphi(\cdot, \varepsilon)$ of the system such that $\varphi(0, \varepsilon) \rightarrow a$ when $\varepsilon \rightarrow 0$.

As usual we have denoted by $d_B(f_1 + \varepsilon f_2, V, 0)$, the Brouwer degree of the function $f_1 + \varepsilon f_2 : V \rightarrow \mathbb{R}^n$ at its fixed point a . A sufficient condition for showing that the Brouwer degree of a function f at its fixed point a is non-zero, is that the Jacobian of the function f at a (when it is defined) is non-zero.

If the function f_1 is not identically zero, then the zeros of $f_1 + \varepsilon f_2$ are mainly the zeros of f_1 for ε sufficiently small. In this case Theorem 3 provides the so-called *averaging theory of first order*.

If the function f_1 is identically zero and f_2 is not identically zero, then the zeros of $f_1 + \varepsilon f_2$ are the zeros of f_2 . In this case Theorem 3 provides the so-called *averaging theory of second order*.