



Anais da Academia Brasileira de Ciências

ISSN: 0001-3765

[aabc@abc.org.br](mailto:aabc@abc.org.br)

Academia Brasileira de Ciências  
Brasil

de Oliveira, Débora R.; Nepomuceno, Delci D.; Castro, Rosane N.; Braz Filho, Raimundo;  
de Carvalho, Mário G.

Special metabolites isolated from *Urochloa humidicola* (Poaceae)

Anais da Academia Brasileira de Ciências, vol. 89, núm. 2, abril-junio, 2017, pp. 789-797

Academia Brasileira de Ciências

Rio de Janeiro, Brasil

Available in: <http://www.redalyc.org/articulo.oa?id=32751197002>

- How to cite
- Complete issue
- More information about this article
- Journal's homepage in [redalyc.org](http://redalyc.org)

[redalyc.org](http://redalyc.org)

Scientific Information System

Network of Scientific Journals from Latin America, the Caribbean, Spain and Portugal

Non-profit academic project, developed under the open access initiative



## Special metabolites isolated from *Urochloa humidicola* (Poaceae)

DÉBORA R. DE OLIVEIRA<sup>1</sup>, DELCI D. NEPOMUCENO<sup>2</sup>, ROSANE N. CASTRO<sup>1</sup>,  
RAIMUNDO BRAZ FILHO<sup>1,3</sup> and MÁRIO G. DE CARVALHO<sup>1</sup>

<sup>1</sup>Universidade Federal Rural do Rio de Janeiro, Instituto de Ciências Exatas, Departamento de Química, Rodovia BR 465, Km 07, 23890-000 Seropédica, RJ, Brazil

<sup>2</sup>Universidade Federal Rural do Rio de Janeiro, Instituto de Zootecnia, Departamento de Nutrição Animal e Pastagens, Rodovia BR 465, Km 07, 23890-000 Seropédica, RJ, Brazil

<sup>3</sup>Setor de Química de Produtos Naturais, Universidade Estadual do Norte Fluminense Darcy Ribeiro, 28013-600 Campos dos Goytacazes, RJ, Brazil

Manuscript received on March 10, 2016; accepted for publication on November 12, 2016

### ABSTRACT

This study aims to identify special metabolites in polar extracts from *Urochloa humidicola* (synonym *Brachiaria humidicola*) that have allelopathic effects and induce secondary photosensitization in ruminants. The compounds were isolated and identified via chromatographic and spectroscopic techniques. The compounds 4-hydroxy-3-methoxy-benzoic acid, *trans*-4-hydroxycinnamic acid, and *p*-hydroxy-benzoic acid; the flavonols isorhamnetin-3-*O*- $\beta$ -D-glucopyranoside and methyl quercetin-3-*O*- $\beta$ -D-glucuronate; and kaempferitrin, quercetin-3-*O*- $\alpha$ -L-rhamnopyranoside, and tricin were identified in the extract from the leaves of *Urochloa humidicola*. Two furostanic saponins, namely, dioscin and 3-*O*- $\alpha$ -L-rhamnopyranosyl-(1-4)-[ $\alpha$ -L-rhamnopyranosyl-(1-2)]- $\beta$ -D-glucopyranosyl-penogenin, as well as catechin-7-*O*- $\beta$ -D-glucopyranoside were identified in the methanolic extract obtained from the roots of this plant. This species features a range of metabolites that may be toxic for animals if used in food and may interfere with the growth medium, thereby inhibiting the development of other species.

**Key words:** *Brachiaria humidicola*, flavonoids, steroidal saponins, *Urochloa*.

### INTRODUCTION

The *Urochloa* genus belongs to the Poaceae family, Paniceae tribe, also known as *Brachiaria* (Morrone and Zuloaga 1992, Veldkamp 1996). The species of this genus have adapted to different soil types (Lapointe 1993) and are used as dead matter for protecting the agricultural soil system. Among *Urochloa* species, *U. decumbens*, *U. brizantha*,

and *U. humidicola* are the most frequently used as animal feed in Brazil.

The frequent use of these species in animal production need a detailed understanding of their potential dangers. In particular, their secondary metabolites include compounds that adversely affect animal health, such as photosensitization in ruminants and horses (Tokarnia et al. 2012). These problems are attributed to the steroidal saponins synthesized by fodder, especially protodioscin (25-

Correspondence to: Débora Ramos de Oliveira  
E-mail: [deboraolv@gmail.com](mailto:deboraolv@gmail.com)

R and 25-S) (Brum et al. 2009). An antinutritional effect has also been reported, which is attributed to the reduction in food intake and digestibility of animals due to the presence of metabolites such as terpenoids and flavonoids (Silva et al. 2012).

In terms of allelopathic effects, these species also have adverse effects on cropping systems; for example, the cinnamic acid derivative metabolized by *Brachiaria* species inhibits seed germination in *Euphorbia heterophylla* and *Bidens pilosa* (Oliveira et al. 2014). However, there may be both negative as well as positive allelopathic effects of different species depending upon crops consortiums. In this context, Rodrigues et al. (2012) cited negative allelopathic effects of the extracts from *U. brizantha* and *U. decumbens* on the seed germination of *Stylosanthes guianensis* and of *U. decumbens* on the germination of *Stylosanthes capitata*; they also reported a positive allelopathic effect of *U. brizantha* on the seeds of *Stylosanthes macrocephala*. The exudates of *U. humidicola* roots inhibit nitrification by *Nitrosomonas europaea* bacteria because of the action of two compounds identified in a phytochemical study of this exudate: *p*-coumaric and ferulic acids. These acids can permeate the cell membrane of these bacteria and inhibit the action of enzymes responsible for their nitrification (Gopalakrishnan et al. 2007, Subbarao et al. 2006). To the best of our knowledge, no study has investigated the isolation and identification of metabolites in the extracts from *U. humidicola*.

Terpenoids and flavonoids in fodder are known to have qualitative effects, such as antinutritional effects resulting from a reduction in consumption and digestibility (Silva et al. 2012). However, flavonoids such as tricetin, quercetin-3-*O*- $\alpha$ -L-rhamnoside, and isorhamnetin 3-*O*- $\beta$ -D-glucoside are cited as beneficial, as they have anti-inflammatory and antioxidant activities (Luyenn et al. 2014, 2015, Riethmüller et al. 2015, Kim et al. 2009) as well as antifungal activity, which are associated with increased feed efficiency

(Aderogba et al. 2013). The saponin dioscin has also been reported to promote beneficial effects, particularly against liver fibrosis (Zhang et al. 2015), beyond its anti-inflammatory (Wu et al. 2015), antitumoral (Kaskiw et al. 2009), and antioxidant (Gao et al. 2012) activities. Additionally, Jayanegara et al. (2014) reported beneficial effects of secondary plant metabolites, particularly in the case of saponins in livestock systems due to the reduction of ruminal methane, which increases animal productivity and provides environmental benefits. These benefits have been yet not explored in animal production, nor has the possibility that metabolites other than cinnamic acid derivatives or a combination of other compounds are responsible for the allelopathic activities. To address this gap, this manuscript presents the first phytochemical study of the extracts from this plant, entailing the isolation of the metabolites in the leaves and roots of *U. humidicola*, to clarify the possible positive and negative effects of its use in animal feed and crop production.

## EXPERIMENTAL

### PLANT MATERIAL

The roots and leaves of *U. humidicola* were collected in April 2013 in an area already established at the Goat Sector of the Animal Science Institute of the Universidade Federal Rural do Rio de Janeiro (UFRRJ) in the municipality of Seropédica-RJ (latitude: 22°44'38"S, longitude: 43°42'27"W; altitude: 26 m). It was identified by Dr. Delci de Deus Nepomuceno, Animal Science Institute, and a voucher specimen (N° RBR-38719) is deposited at the Herbarium of Biology Institute, UFRRJ.

### EQUIPMENTS

One- and two-dimensional  $^1\text{H}$  and  $^{13}\text{C}$  nuclear magnetic resonance (NMR) spectra were obtained on Bruker NMR spectrometer [400 and 500 MHz ( $^1\text{H}$ ), 100 and 125 MHz ( $^{13}\text{C}$ )] using tetramethylsilane

(TMS) as an internal standard for chemical shift reference. Some compounds were identified using a GC-MS unit (model GC-MS- QP-2010 Plus, from Shimadzu, Japan) equipped with HP-5 fused silica capillary column (30mm x 0.25mm i.d x film thickness 0.25  $\mu$ m) with a quadrupole mass analyzer and electron impact ionization at 70 eV. Electrospray ionization–high-resolution mass spectrometry (EI–HRMS) spectra were recorded on a quadrupole/time-of-flight instrument (microOTOF II, Bruker Daltonics, Billerica, MA). High-performance liquid chromatography (HPLC) analyses were performed using an instrument equipped with pump LC-10AS, SPD-10A detector, CBM-20A-Communications Module (Shimadzu), and Rheodyne injector with loop of 500  $\mu$ L. All other equipment commonly used for the preparation and fractionation of extracts belongs to the Laboratory of Natural Products Chemistry at UFRRJ.

#### EXTRACTION AND ISOLATION

The botanical material was dried at room temperature without exposure to sunlight. The 2.1 kg of leaves and 1.81 kg of roots obtained were then ground separately in a hammer mill at the Laboratory of Animal Nutrition Science, Animal Science Institute, UFRRJ. The milled material was subjected to extraction by maceration using hexane, methanol, and methanol/water (8:2) as solvents. The extractions were performed thoroughly at 7-day intervals, and each solution was concentrated on a rotary evaporator under vacuum. This process yielded six extracts: hexane leaves (UHFH), methanol leaves (UHFM), methanol: water leaves (UHFMH<sub>2</sub>O), hexane roots (UHRH), methanol roots (UHRM), and methanol: water roots (UHRMH<sub>2</sub>O). The hydromethanolic and methanolic extracts were solubilized in methanol: water (8:2) subjected to liquid–liquid partitioning by organic solvents of increasing polarity: hexane, dichloromethane, ethyl acetate, and butanol. The

ethyl acetate fractions obtained from the methanol (UHFM-Ac) and hydromethanolic (UHFMH<sub>2</sub>O-Ac) extract from the leaves and the methanol (UHRM-Ac) extracts from the roots were selected for chromatographic fractionation.

First, 6.079 g of UHFMH<sub>2</sub>O-Ac was subjected to chromatographic fractionation on 70–230 mesh silica gel (224.00 g) using dichloromethane, ethyl acetate, and methanol as the mobile phase, in a gradient of increasing polarity. In total, 320 fraction of 100 ml were collected. The fractions were pooled according to their thin-layer chromatography (TLC) profiles. The group containing fractions 51–56 (0.1342 g), which were obtained with dichloromethane:ethyl acetate (8:2) as the mobile phase, was analyzed by <sup>13</sup>C and <sup>1</sup>H NMR and GC–MS. These analyses allowed the identification of 4-hydroxy-3-methoxy-benzoic acid (**1**), *trans*-4-hydroxycinnamic acid (**2**), and 4-hydroxy-benzoic acid (**3**). The group containing fractions 80–86 (0.1024 g), which were obtained with dichloromethane: ethyl acetate (6: 4), was purified on a silica gel column. The purified fractions were then analyzed by <sup>13</sup>C and <sup>1</sup>H NMR to identify isorhamnetin-3-*O*- $\beta$ -D-glucopyranoside (**4**) and methyl-quercetin-3-*O*- $\beta$ -D-glucuronate (**5**). The group containing fractions 133–136 (0.4668 g), which were obtained with ethyl acetate: methanol (8:2), were combined and chromatographed on a Sephadex LH-20 column with methanol elution. This process produced 34 fractions, and kaempferitrin (**6**) was obtained.

The ethyl acetate fraction UHFM-Ac (1.59 g) was chromatographed on 70–230 mesh silica gel column (54.50 g) with dichloromethane, ethyl acetate, and methanol elution in a gradient of increasing polarity. In total, 152 fractions of 25 ml were collected. Fractions were combined on the basis of TLC analysis. The group of fractions 22–26 (0.0821 g), obtained with ethyl acetate as eluent, was eluted on Sephadex LH-20 column using methanol as eluent. The analysis

of the fractions led to the identification of tricetin (**8**). Fractions 97–99 (0.1240 g), obtained with ethyl acetate and methanol (4:6), were subjected to additional chromatographic fractionation on a silica gel column, and the analysis of the fraction led to the identification of quercetin-3-*O*- $\alpha$ -L-rhamnopyranoside (**7**).


The fraction UHRM-Ac (5.74 g) was subjected to fractionation on 70–230 mesh silica gel column (190.00 g) by elution with ethyl acetate and methanol in a gradient of increasing polarity. In total, 227 fractions of 100 ml were collected. The fractions were pooled according to their chromatographic profiles observed by TLC analysis. The group of fractions 120–124 (0.3082 g), obtained with ethyl acetate: methanol (6:4), was subjected to fractionation by HPLC. This chromatographic fractionations were composed of water as solvent (A) and acetonitrile as solvent (B) in a 4:6 ratio. The mobile phase was filtered before use and delivered isocratically at a flow rate of 4 ml min<sup>-1</sup>. The analysis was conducted at room temperature using a Phenomenex C18 semi-preparative column (250 mm  $\times$  10 mm i.d., 5  $\mu$ m), and the analytes were monitored at 205 nm. In addition to some unidentified components, saponins represented by peaks at 3.084 and 5.801 min were isolated, analyzed by <sup>1</sup>H and <sup>13</sup>C NMR and HRMS, and identified as the steroidal saponins dioscin (**9**) and 3-*O*- $\alpha$ -L-rhamnopyranosyl (1-4)-[ $\alpha$ -L-rhamnopyranosyl-(1-2)]- $\beta$ -D-glucopyranosides penogenin (**10**), respectively. The group of fractions 129–133 (0.5340 g), eluted with ethyl acetate and methanol (6:4), was subjected to fractionation on a Sephadex LH-20 column using methanol as the eluent. This fractionation produced 47 fractions, among which fractions 36–40 (0.0197 g) were recombined and filtered on (230–400 mesh) flash silica column using ethyl acetate:methanol (8:2) as the eluent. The NMR analysis of fractions 1–3 (0.025 g) then enabled the identification of catechin-7-*O*- $\beta$ -D-glucopyranoside (**11**).

## RESULTS AND DISCUSSION

The identification of compounds **1**, **2**, and **3** (Figure 1) in mixture was accomplished by interpretation of the mass spectra and the <sup>1</sup>H and <sup>13</sup>C NMR spectra, including such two-dimensional experiments as heteronuclear single-quantum coherence (HSQC), heteronuclear multiple bond correlation (HMBC), and <sup>1</sup>H–<sup>1</sup>H correlation spectroscopy (COSY). Analysis of the <sup>1</sup>H NMR and <sup>1</sup>H–<sup>1</sup>H COSY spectra allowed the identification of signals compatible with *trans*-carboxylic acid (**2**), *para*-substituted aromatic rings in **2** and **3**, and an ABC system in **1**. The doublets at  $\delta_H$  7.48, and 6.31 ( $J = 16$  Hz), represent hydrogens 7 and 8, respectively, of compound **2**. The singlet at  $\delta_H$  7.52, and the doublets at  $\delta_H$  6.80 ( $J = 8.5$  Hz) of the hydrogens assigned to the 2, 6 and 3, 5 positions in compound **2**; the signals at  $\delta_H$  6.84 and 7.43 assigned to the hydrogens of the *p*-substituted aromatic ring of **3**; and the remaining signals at  $\delta_H$  7.78 (d,  $J = 8.5$  Hz), 7.43 (s), and 6.82 (d,  $J = 8.5$  Hz) represent the hydrogens of the aromatic ring of compound **1**. Analysis of the <sup>13</sup>C-NMR and two-dimensional (HSQC and HMBC) spectra and comparison with the literature data confirmed the proposed structures of 4-hydroxy-3-methoxy-benzoic acid for **1** (Pouchert and Behnke 1993a), *trans*-4-hydroxycinnamic acid for **2** (Wang et al. 2011), and *p*-hydroxybenzoic acid for **3** (Pouchert and Behnke 1993b). GC–MS analysis of the fraction containing these compounds confirmed the proposed structures and even the presence of a methoxyl group in **1**.

Compounds **4–8** (Figure 1) were identified as flavonoids on the basis of their <sup>1</sup>H and <sup>13</sup>C NMR spectra, which are consistent with the molecular skeleton of flavonols. Compounds **4** and **5** were identified in mixture from the <sup>1</sup>H NMR spectrum, which presented signals at  $\delta_H$  6.21 (d,  $J = 2.2$  Hz), corresponding to H-6 (both **4** and **5**), and  $\delta_H$  6.42 (d,  $J = 2.2$  Hz) and 6.45 (d,  $J = 2.2$  Hz), corresponding to H-8 of **4** and **5**, respectively. Additional signals





**Figure 1** - Compounds identified in the extracts from *Urochloa humidicola*.

at  $\delta_{\text{H}}$  7.51 (d,  $J = 2.2$  Hz, H-2'), 7.57 (dd,  $J = 8.5$  Hz and 2.2 Hz, H-6'), and 6.85 (d,  $J = 8.5$  Hz, H-5') were attributed to the protons of ring B of compound **4**. The signals at  $\delta_{\text{H}}$  7.52 (d,  $J = 2.0$  Hz), 6.93 (d,  $J = 8.5$  Hz), and 7.59 (dd,  $J = 8.0$  and 2.0 Hz) were proposed to correspond to the protons H-2', H-5', and H-6' of ring B in **5**, respectively. Additional analysis of the  $^{13}\text{C}$  and 2D NMR spectra confirmed the proposed flavonoid skeleton. These analyses allowed the identification of the signals of two sugar moieties that were linked in each flavonol. Two doublets at  $\delta_{\text{H}}$  5.59 ( $J = 7.25$  Hz) and 5.49 ( $J = 7.25$  Hz) were assigned to the hydrogens at the anomeric carbons of the  $\beta$ -D-glycopyranosyl unit. These proposed assignments were confirmed by additional signals observed in the HSQC spectrum.

The location of two methoxyl groups and the sugar moieties were determined from the HMBC and nuclear Overhauser effect (NOE) spectra. These analyses allowed the location of the sugar unit at C-3 of each flavonoid, a methoxyl group at C-3' of **4**, and a methyl ester in the sugar moiety of **5**. These assignments were confirmed by the absence of  $\text{CH}_2$ -6'' and the connection of this group to a carboxyl group. The  $^1\text{H}$  and  $^{13}\text{C}$  chemical shifts were in agreement with the literature data reported for isorhamnetin-3-*O*- $\beta$ -D-glucopyranoside (**4**) (Yuan et al. 2013) and methyl-quercetin-3-*O*- $\beta$ -D-glucuronate (**5**) (Hilbert et al. 2015).

The  $^1\text{H}$  NMR spectrum of flavonoid **7** presented signals at  $\delta_{\text{H}}$  7.85 (d,  $J = 8.5$  Hz) corresponding to H-2',6' of ring B, and  $\delta_{\text{H}}$  6.95 (d,  $J = 8.5$  Hz),

corresponding to H-3',5' of ring B. The doublets at  $\delta_{\text{H}}$  6.47 (d,  $J=1.8$  Hz) and 6.72 (d,  $J=1.85$  Hz) were assigned to hydrogens H-6 and H-8, respectively. The presence of the sugar units was confirmed by broad singlets at  $\delta_{\text{H}}$  5.58 (brs) and 5.40 (brs), which are characteristic of hydrogen at the anomeric carbon of rhamnose. This assignment was confirmed by the doublets at  $\delta_{\text{H}}$  1.27 ppm ( $J=5.95$  Hz) and 0.96 ppm ( $J=5.65$  Hz). The location of the sugars in the structure was determined by HMBC analysis, which revealed long-range coupling between  $\delta_{\text{H}}$  and  $\delta_{\text{C}}$ , 5.58/162.58 (H-1"/C-7) and 5.40/135.47 (H-1"/C-3). The mass spectrum recorded in negative mode showed a signal at  $m/z$  577.15 [M-H], consistent with the molecular formula  $\text{C}_{27}\text{H}_{29}\text{O}_{14}$ , as well as signals at  $m/z$  431.09 [(M-H)-146] and 285.03 [(M-H)-292]. These data allowed the identification of compound **6** as kaempferitrin (Pizzolatti et al. 2003).

The  $^1\text{H}$  NMR spectrum showed signals corresponding to five hydrogens in aromatic systems, which were attributed to the A and B rings of quercetin as the basic skeleton of compound **7**. The chemical shifts observed in the  $^{13}\text{C}$  NMR spectrum confirmed this assignment. The signal at  $\delta_{\text{H}}$  5.37 (s), characteristic of a proton at an anomeric carbon, along with the observed signal at  $\delta_{\text{H}}$  0.97 (d,  $J=7.0$  Hz) in the  $^1\text{H}$  NMR spectrum led us to propose that a rhamnose unit was part of the structure. The position of the sugar was determined from the HMBC spectrum, which revealed a coupling between the hydrogen represented by the singlet at  $\delta_{\text{H}}$  5.37 (H-1") and C-3 ( $\delta_{\text{C}}$  134.82). Further analysis of the  $^1\text{H}$  and  $^{13}\text{C}$  NMR spectra and comparison with the literature led to the identification of compound **7** as quercetin 3-*O*- $\beta$ -D-rhamnoside (Ozgem et al. 2010). Compound **8** was obtained as a yellow crystalline solid, soluble in methanol. Its  $^1\text{H}$  NMR spectrum showed only five signals: one singlet integrating to two hydrogens at  $\delta_{\text{H}}$  7.40, which was attributed to the H-2',6'; two doublets at  $\delta_{\text{H}}$  6.27 (d,  $J=2.2$  Hz) and 6.57 (d,  $J=2.2$  Hz) for the H-6 and H-8 of ring

A of a flavonoid; one singlet at  $\delta_{\text{H}}$  3.98 integrating to six hydrogens related to two methoxyls; and a singlet at  $\delta_{\text{H}}$  6.75, which was assigned to the H-3 of a flavonoid. The 2D NMR spectra allowed the location of the methoxyl at C-3' and C-5'. These analyses and comparison with the literature led to the identification of compound **8** as triclin (Zielinska et al. 2008).

Compounds **9** and **10** (Figure 1) were identified as the furostanoic steroidal saponins. Both presented three carbohydrate units, glucose linked directly to C-3 of the aglycone and two rhamnoses linked to the glucose at C-2' and C-4'. These saponins differ only in a hydroxyl group found at C-17 in compound **9**. The positive-ion-mode HRMS spectrum of **9** showed peaks at  $m/z$  891.4695 [M + Na], 869.4893 [M + H], 723.4309 [M-146], 577.3731 [M-146-146], and 415.3181 [M-146-146-162], and that of **10** showed peaks at  $m/z$  907.4653 [M + Na], 885.4843 [M + H], 739.6032 [M-146], 593.3676 [M-146-146], and 431.3143 [M-146-146-162]. This difference of 16 units indicates the presence of an additional hydroxyl in **10**.

The  $^{13}\text{C}$  NMR spectra of **9** and **10** both with signals of 45 carbons, including 4 and 5 quaternary carbons in compounds **9** and **10**, respectively; 24 and 23 methyne carbons in compounds **9** and **10**, respectively; and 11 methylene and 6 methyl carbons in both compounds. The chemical shift of the quaternary carbon at  $\delta_{\text{C}}$  110 of a spiro system is characteristic of the tetrahydrofuran and pyran rings in the structure. The relative stereochemistry of C-25 was identified as (*R*) by comparison of the chemical shifts of methyl-27 and  $\text{CH}_2$ -23 (Pires et al. 2002, Espejo et al. 1982). The presence of three methyne carbons with chemical shifts near 100 ppm was compatible with the presence of pyranosyl units, on what differences in the mass spectra confirmed the existence of two rhamnose and one glucose units in each compound.

The presence of a hydroxyl linked to the C-17 carbon in compound **10** affects the C-16 and C-17

chemical shifts in the  $^{13}\text{C}$  NMR spectra; these signals shift downfield to approximately 90 ppm. Detailed analysis of the  $^1\text{H}$  and  $^{13}\text{C}$  NMR spectra besides comparison with the literature data led to the identification of **9** as dioscin (Pires et al. 2002), and **10** as 3-*O*- $\alpha$ -L-rhamnopyranosyl-(1-4)-[ $\alpha$ -L-rhamnopyranosyl-(1-2)]- $\beta$ -D-glucopyranosides penogenin (Espejo et al. 1982, Feng et al. 2007).

The  $^1\text{H}$  and  $^{13}\text{C}$  NMR spectra of compound **11** showed characteristic signals of a flavanol, with saturation in the C ring. Analysis of the 1D and 2D  $^1\text{H}$  and  $^{13}\text{C}$  NMR spectra and comparison with the values for catechin led to identifying it as flavane. The additional signals were compatible with a structure containing glucose, and HMBC analysis indicated that it is located at C-7. These analyses led to the identification of compound **11** (Figure 1) as catechin-7-*O*- $\beta$ -D-glucopyranoside (Benavides et al. 2006, Agrawal 1989).

### CONCLUSIONS

Twelve substances, including flavonoids, have been reported in this species for the first time in this work. The saponin diosgenin, which was identified in the extracts from the roots of *U. humidicola*, has been isolated from the leaves *U. decumbens*, synonym *Brachiaria decumbens* (Pires et al. 2002). This is the first identification of the steroidal saponin penogenin in the *Urochloa* genus.

The identification of *p*-coumaric acid and other benzoil acid derivatives is consistent with the literature which reports that phenolic compounds, such as those present in *U. humidicola*, promote allelopathic action, and prevent pasture invasion plants in monoculture systems, independent of organic acids (Souza Filho et al. 2005, Kobayashi and Noguchi 2015). However, for pasture system consortiums, the presence of these substances has disadvantages, as the resulting inefficiency of the systems hinders the establishment of different plant species (Rodrigues et al. 2012). The identified

constituents improve our understanding of the diversity of metabolites produced by these species. In the literature, metabolites responsible only for the photosensitization and/or allelopathic effects of *Urochloa* genus have been reported. In contrast, the saponins and flavonoids identified in this study have not been associated with these effects, which merit careful consideration. The identified flavonoids have biological activities and can therefore endow ruminants with beneficial functions, increasing their nutritional value.

### ACKNOWLEDGMENTS

The authors thank to Fundação Carlos Chagas Filho de Amparo a Pesquisa do Estado do Rio de Janeiro (FAPERJ), to Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq), and to Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES) for scholarship and financial support, Dr. Norberto Peoporini Lopes (NPPNS- Núcleo de Pesquisa em Produtos Naturais e Sintéticos) for the HRMS.

### REFERENCES

- ADEROGBA M, NDHLALAA, RENGASAMY K AND VAN STADEN J. 2013. Antimicrobial and Selected In Vitro Enzyme Inhibitory Effects of Leaf Extracts, Flavonols and Indole Alkaloids Isolated from *Croton menyharthii*. Mol 18(10): 12633-12644.
- AGRAWAL PK. 1989. Carbon-13 MNR of Flavonoids. Central Institute of Medicinal and Aromatic Plants, Lucknow- India: Elsevier: 474-478.
- BENAVIDES A, MONTORO P, BASSARELLO C, PIACENTE S AND PIZZA C. 2006 Catechin derivatives in *Jatropha macrantha* stems: Characterisation and LC/ESI/MS/MS qualitative-quantitative analysis. J Pharm Biomed Anal 40: 639-647.
- BRUM KB, HARAGUCHI M, GARUTTI MB, NÓBREGA FN AND ROSA B. 2009. Fioravanti MCS. Steroidal saponin concentrations in *Brachiaria decumbens* and *B. brizantha* at different developmental stages. Cienc Rural 39(1): 279-281.
- ESPEJO O, LAVOT JC, JUNG H AND GIRAL F. 1982. Spirostanic diosgenin precursors from *Dioscorea composita* tubers. Phytochemistry 21(2): 413-416.



- FENG B, KANG L, MA B, QUAN B, ZHOU W, WANG Y, ZHAO Y, LIU Y AND WANG S. 2007. The substrate specificity of a glucoamylase with steroidal saponin-rhamnoside activity from *Curvularia lunata*. *Tetrahedron* 63: 6796-6812.
- GAO L, LI F, YAO S AND QIN S. 2012. The Anti-oxidative and Anti-inflammatory Effects of Dioscin from the Roots of *Paris polyphylla* Smith. *International Symposium on Information Technology in Medicine and Education*, p. 786-789.
- GOPALAKRISHNAN S, SUBBARAO GV, NAKAHARA K, YOSHIHASHI T, ITO O, MAEDA I, ONO H AND YOSHIDA M. 2007. Nitrification Inhibitors from the root tissues of *Brachiaria humidicola*, a tropical grass. *J Agric Food Chem* 55(4): 1385-1388.
- HILBERT G ET AL. 2015. Flavonol profiles in berries of wild *Vitis* accessions using liquid chromatography coupled to mass spectrometry and nuclear magnetic resonance spectrometry. *Food Chem* 169: 49-58.
- JAYANEGARA A, WINA E, TAKAHASHI J. 2014. Meta-analysis on Methane Mitigating Properties of Saponin-rich Sources in the Rumen: Influence of Addition Levels and Plant Sources. *Asian Australas J Anim Sci* 27(10): 1426-1435.
- KASKIW MJ, TASSOTTO ML, MOK M, TOKAR SL, PYCKO R, TH'NG J and JIANG Z-H. 2009. Structural analogues of diosgenyl saponins: Synthesis and anticancer activity. *Bioorg Med Chem* 17(22): 7670-7679.
- KIM YA, KONG C, UM YR, LIM S, YEA SS AND SEO Y. 2009. Evaluation of *Salicornia herbacea* as a Potential Antioxidant and Anti-Inflammatory Agent. *J Med Food* 12(3): 661-668.
- KOBAYASHI A AND NOGUCHI HK. 2015. The seasonal variations of allelopathic activity and allelopathic substances in *Brachiaria brizantha*. *Bot Stud* 56: 25.
- LAPOINTE SL. 1993. Manejo de los plagas clave para forrajes de las sabanas neotropicales. *Past Trop* 15(3): 1-9.
- LUYENN BTT, LEE YM, LEE SH, JANG HD AND KIM YH. 2015. The Anti-Osteoporosis and Antioxidant Activities of Chemical Constituents from *Chrysanthemum indicum* Flowers. *Phytother Res* 29: 540-548.
- LUYENN BTT, TAI BH, THAO NP, CHA JY, LEE HY, LEE YM AND KIM YH. 2014. Anti-inflammatory components of *Chrysanthemum indicum* flowers. *Bioorg Med Chem Lett* 25(2): 266-269.
- MORRONE O AND ZULOAGA FO. 1992. Revision de las especies sudamericanas nativas e introducidas de los géneros *Brachiaria* y *Urochloa* (Poaceae: Panicoideae: Paniceae). *Darwiniana* 4(1/4): 43-109.
- OLIVEIRA RS, RIOS FA, CONSTANTIN J, ISHII-IWAMOTO EL, GEMELLI A AND MARTINI PE. 2014. Grass straw mulching to suppress emergence and early growth weeds. *Planta Daninha* 32: 11-17.
- OZGEM U, SEVINDIK H, KAZAZ C, YIGIT D, KANDEMIR A, SECEN H AND CALIS I. 2010. A new sulfated  $\alpha$ -ionone glycoside from *Sonchus erzincanicus*. *Mol* 15: 2593-2599.
- PIRES VS, TAKETA ATC, GOSMANN G AND SCHENKEL EP. 2002. Saponins and Sapogenins from *Brachiaria decumbens* Stapf. *J Braz Chem Soc* 13(2): 135-139.
- PIZZOLATTI MG, JUNIOR AC, SZPOGANICZ B, SOUZA E, BRAZ-FILHO R AND SCHRIPEMA J. 2003. Flavonoides glicosilados das folhas e flores de *Bauhinia fortificata* (Leguminosae). *Quim Nova* 26: 466-469.
- POUCHERT CJ AND BEHNKE J. 1993a. The Aldrich Library of  $^{13}\text{C}$  and  $^1\text{H}$  FT NMR spectra, 1<sup>st</sup> ed., Milwaukee: Aldrich Chemical Company, 1115 p.
- POUCHERT CJ AND BEHNKE J. 1993b. The Aldrich Library of  $^{13}\text{C}$  and  $^1\text{H}$  FT NMR spectra, 1<sup>st</sup> ed., Milwaukee: Aldrich Chemical Company, 1084 p.
- RIETHMÜLLER E, TÓTH G, ALBERTI Á, VÉGH K, BURLINI I, KÖNCZÖL Á, BALOGH, GT AND KÉRY Á. 2015. First characterisation of flavonoid- and diarylheptanoid-type antioxidant phenolics in *Corylus maxima* by HPLC-DAD-ESI-MS. *J Pharm Biomed Anal* 107: 159-167.
- RODRIGUES APDC, LAURA VA, PEREIRA SR AND DEISS C. 2012. Alelopatia de duas espécies de braquiária em sementes de três espécies de estilosantes. *Cienc Rural* 42(10): 1758-1763.
- SILVA NDS, SILVA HDS, MARCKS E, ANDRADE G AND SOUSA DE JR. 2012. Fatores antinutricionais em plantas forrageiras. *Antinutritional factors in forages*. *RVADS* 7(4): 1-7.
- SOUZA FILHO APS, PEREIRA AAG AND BAYMA JCA. 2005. Aleloquímico produzido pela gramínea forrageira *Brachiaria humidicola*. *Planta Daninha* 23: 25-32.
- SUBBARAO GV, ISHIKAWA T, ITO O, NAKAHARA K, WANG HY AND BERRY WL. 2006. A bioluminescence assay to detect nitrification inhibitors released from plant roots: a case study with *Brachiaria humidicola*. *Plant Soil* 288: 101-112.
- TOKARNIA CH, BRITO MF, BARBOSA JD, PEIXOTO PV AND DOBEREINER J. 2012. Plantas Tóxicas do Brasil para Animais de Produção. 2<sup>a</sup> ed., Editora Helianthus, Rio de Janeiro, Brasil, p. 323-333.
- VELDKAMP JF. 1996. *Brachiaria*, *Urochloa* (Gramineae-Paniceae) in Malesia. *Blumea – Biodiversity, Evolution and Biogeography of Plants* 41: 413-437.
- WANG XX, HE JM, WANG CL, ZHANG RP, HE WY, GUO SX, SUN RX AND ABLIZ Z. 2011. Simultaneous Structural Identification of Natural Products in Fractions of Crude Extract of the Rare Endangered Plant *Anoectochilus roxburghii* Using  $^1\text{H}$  NMR/RLC-MS Parallel Dynamic Spectroscopy. *Int J Mol Sci* 12: 2556-2571.

- WU S, XU H, PENG J, WANG C, JIN Y, LIU K, SUN H AND QIN J. 2015. Potent anti-inflammatory effect of dioscin mediated by suppression of TNF- $\alpha$ -induced VCAM-1, ICAM-1 and EL expression via the NF- $\kappa$ B pathway. *Biochimie* 110: 62-72.
- YUAN H, ZHOU X, MENG L, QUIN F AND ZHOU G. 2013. Chemical constituents from *Commelina communis*. *China J of Chin Mater Med* 38: 3304-3308.
- ZHANG X, HAN X, YIN L, XU L, QI Y, XU Y, SUN H, LIN Y, LIU K AND PENG J. 2015. Potent effects of dioscin against liver fibrosis. *Sci Rep* 5: 9713.
- ZIELINSKA A, PARADOWSKA K, JAKOWSKI J AND WAWER I. 2008.  $^{13}\text{C}$  CP MAS NMR and GIAO-CHF/DFT calculation of flavonoids: Morin, kaempferol, tricetin, genistein, formononetin and 3,7-dihydroxyflavone. *J Mol Struct* 873: 109-116.