



Ciência Rural

ISSN: 0103-8478

cienciarural@mail.ufsm.br

Universidade Federal de Santa Maria  
Brasil

Streck, Nereu Augusto; Bellé, Rogério Antonio; Kersting da Rocha, Edileusa; Schuh, Mariângela  
Estimating leaf appearance rate and phyllochron in safflower (*Carthamus tinctorius* L.)

Ciência Rural, vol. 35, núm. 6, novembro-dezembro, 2005, pp. 1448-1450

Universidade Federal de Santa Maria

Santa Maria, Brasil

Available in: <http://www.redalyc.org/articulo.oa?id=33135636>

- How to cite
- Complete issue
- More information about this article
- Journal's homepage in redalyc.org

redalyc.org

Scientific Information System

Network of Scientific Journals from Latin America, the Caribbean, Spain and Portugal

Non-profit academic project, developed under the open access initiative

## Estimating leaf appearance rate and phyllochron in safflower (*Carthamus tinctorius* L.)

### Estimativa da taxa de aparecimento de folhas e do filocrono em cártamo (*Carthamus tinctorius* L.)

Nereu Augusto Streck<sup>1</sup> Rogério Antonio Bellé<sup>1</sup> Edileusa Kersting da Rocha<sup>2</sup>  
Mariângela Schuh<sup>3</sup>

#### - NOTE -

#### ABSTRACT

Safflower may be an interesting option for the flower market, either as fresh or dried cut flower. Estimating the leaf appearance rate and the phyllochron (the time interval between the appearance of successive leaves) is important for calculating the number of emerged leaves (NL) on the plant, which is an excellent measure of plant development. The objective of this study was to estimate the leaf appearance rate and the phyllochron in safflower (*Carthamus tinctorius* L.). An experiment was conducted in Santa Maria, RS, Brazil, inside an 8 x 15m plastic greenhouse. Sowing was on 03 October 2003 and emergence was on 08 October 2003. The experimental design was a randomized complete block design with three replications. The main stem NL was measured twice a week from 24 October 2003 to 15 November 2003 in four plants per replication. Daily growing degree days above a base temperature (5°C) and accumulated thermal time (TT) were calculated. The NL was linearly regressed against TT. The angular coefficient of the linear regression is the LAR (leaves/°C day) and the phyllochron (°C days leaf<sup>-1</sup>) was estimated by the inverse of the angular coefficient of the linear regression. The LAR was  $0.0467 \pm 0.0203$  leaves °C<sup>-1</sup> day and the phyllochron was  $25.5 \pm 14.6$ °C days leaf<sup>-1</sup>.

**Key words:** leaf emergence, thermal time, safflower.

#### RESUMO

Cártamo pode ser uma alternativa para o mercado de flores, podendo ser comercializada como flor

fresca ou seca. A estimativa da taxa de aparecimento de folhas (LAR) e do filocrono (tempo necessário para o aparecimento de folhas sucessivas) é importante no cálculo do número de folhas emergidas (NL) na planta, o qual é uma excelente medida de tempo vegetal. O objetivo deste trabalho foi estimar a taxa de aparecimento de folhas e o filocrono em cártamo (*Carthamus tinctorius* L.). Foi realizado um experimento em Santa Maria, RS, em estufa plástica de 8 x 15m. A semeadura foi em 03 de outubro de 2003 e a emergência foi em 08 de outubro de 2003. O delineamento experimental foi o de blocos ao acaso com três repetições. O NL na haste principal foi medido durante o período de 24 de outubro a 15 de novembro de 2003. Foi calculado o número de graus dia diário acima de uma temperatura base (5°C) e a soma térmica acumulada (TT). Foi realizada uma análise de regressão linear entre NL e TT. O coeficiente angular da regressão linear é a LAR (folhas°C<sup>-1</sup> dia) e o filocrono (°C dia folha<sup>-1</sup>) foi estimado pelo inverso do coeficiente angular da regressão linear. A LAR foi  $0,0467 \pm 0,0203$  folhas °C<sup>-1</sup> dia e o filocrono foi  $25,5 \pm 14,6$ °C dia folha<sup>-1</sup>.

**Palavras-chave:** emergência de folhas, soma térmica, cártamo.

Safflower (*Carthamus tinctorius* L.), Asteraceae family, an annual crop, is native of Asia, where it is widely grown for textile dyeing and for an edible oil, which is extracted from the seeds (POLUNIN, 1991). Safflower has high ornamental value in Europe

<sup>1</sup>Departamento de Fitotecnia, Centro de Ciências Rurais (CCR), Universidade Federal de Santa Maria (UFSM), 97105-900, Santa Maria, RS, Brazil. E-mail: nstreck1@smail.ufsm.br. Corresponding author.

<sup>2</sup>Programa de Pós-graduação em Agronomia, CCR, UFSM, Santa Maria, RS, Brazil.

<sup>3</sup>Curso de Agronomia, CCR, UFSM, Santa Maria, RS, Brazil.

(MULLER FLOWERSEEDS, 1998). Major markets for safflower include fresh and dried cut flowers (STEVENSON, 1998). In North America and Latin America, safflower is used mainly for oil (GAYETTO et al., 1999; OELKE et al., 2004). It is estimated that about 0.85 million ha are cultivated with safflower world wide (FAOSTAT, 2004). In Brazil, safflower is not used for oil and it is little known for ornamental purpose.

Flowers are an increasing market in Brazil. Brazilian flower production is concentrated in the São Paulo State, with about 70% of the national flower production. Rio Grande do Sul State is an emerging center in area and number of flower growers (BATISTA, 2000). Safflower may be an interesting flower crop for this emerging center, either as fresh or dried cut flower.

Leaf appearance rate (LAR) is the number of leaves that become visible on a stem per unit time (STRECK et al., 2003). The calculation of LAR is an important component of many crop simulation models (e.g. AMIR & SINCLAIR, 1991). The integration of LAR on the main stem over time gives the number of emerged leaves (NL) on the plant main stem, which is an excellent measure of plant development. The main stem NL is related to tiller appearance (KLEPPER et al., 1982; RICKMAN & KLEPPER, 1991; McMASTER et al., 1991) and to the timing of certain key plant developmental stages (BROOKING et al., 1995; CALDERINI et al., 1996; ROBERTSON et al., 1996; JAMIESON et al., 1998). Accurately calculating the appearance of individual leaves and the rate of leaf area expansion also has an impact on calculating light interception and absorption by the canopy, canopy photosynthesis, and therefore, accumulation of dry matter and yield (AMIR & SINCLAIR, 1991; HODGES & RITCHIE, 1991; McMASTER et al., 1991).

Another approach to calculate NL is through the concept of the phyllochron, which is the time interval between the appearance of successive leaves (KLEPPER et al., 1982; KIRBY, 1995). Time can be expressed in thermal time (TT), measured in units of degree-days ( $^{\circ}\text{C day}$ ), which is a better plant time descriptor than day of the year and number of days after sowing (GILMORE & ROGERS, 1958; RUSSELE et al., 1984; McMASTER & SMIKA, 1988).

A literature search yielded no previous studies aiming to quantify LAR and the phyllochron in safflower, which constituted the rationale for this effort. The objective of this study was to estimate the leaf appearance rate and the phyllochron in safflower.

An experiment was conducted at Santa Maria, RS, Brazil, (latitude:  $29^{\circ}43'S$ , longitude:  $53^{\circ}42'S$  and altitude: 95m) inside an 8 x 15m plastic greenhouse. Sowing was on 03 October 2003 and emergence was

on 08 October 2003. Plants were well fertilized and irrigated as needed to avoid stress due to low soil nutrient and water.

The experimental design was a randomized complete block design with three replications. Each replication had  $1\text{m}^2$  and was composed by eight rows of plants and six plants/row ( $48\text{ plants m}^{-2}$ ). Plant spacing was 0.125m among rows and 0.165m among plants. Four plants located in the two center rows were randomly selected and tagged with colored wires after emergence. These plants were used to measure the visible main stem NL twice a week from 24 October 2003 to 15 November 2003.

Daily minimum and maximum air temperatures were recorded inside the plastic greenhouse throughout the experiment with a mercury-in-glass thermometer at 1.5m height. Daily growing degree-days (GDD,  $^{\circ}\text{C day}$ ) were calculated as (GILMORE & ROGERS, 1958; ARNOLD, 1960):

$$\text{GDD} = (\text{Tmean} - \text{Tb}) \cdot 1 \text{ day} \quad \{^{\circ}\text{C day}\} \quad (1)$$

where Tmean is the mean air temperature calculated as the average of daily minimum and maximum air temperatures, and Tb is the base temperature, assumed  $5^{\circ}\text{C}$ . Two reasons contributed to adopt a  $\text{Tb}=5^{\circ}\text{C}$  for safflower. The first reason is that a literature search yielded no value of Tb for this species. The second reason is that previous experience with growing this species in Santa Maria showed that it does not grow well in winter time and in summer time, but grows well in the spring (R.A. BELLÉ, personal communication). Winter crops (e.g. wheat) have a Tb around  $0^{\circ}\text{C}$  (STRECK et al., 2003) and summer crops (e.g. maize, sorghum, and soybean) have a Tb around  $10^{\circ}\text{C}$  (WARRINGTON & KANEMASU, 1983; MAJOR et al., 1990; SINCLAIR, 1986).

Accumulated thermal time (TT) from 24 October 2003 was calculated by:

$$\text{TT} = \sum \text{GDD} \quad \{^{\circ}\text{C day}\} \quad (2)$$

The NL was linearly regressed against TT. The angular coefficient of the linear regression is the LAR (leaves  $^{\circ}\text{C}^{-1} \text{ day}$ ) and the phyllochron ( $^{\circ}\text{C days/leaf}$ ) was estimated by the inverse of the angular coefficient of the linear regression, i.e.  $1/\text{LAR}$  (KLEPPER et al., 1982; KIRBY, 1995).

There was a strong relationship ( $R^2=0.95$  or greater) between NL and TT. The average LAR was  $0.0467 \pm 0.0203$  leaves  $^{\circ}\text{C}^{-1} \text{ day}$  and the average phyllochron was  $25.5 \pm 14.6^{\circ}\text{C days leaf}^{-1}$ . The LAR and the phyllochron estimated in block I was at least two-fold lower or greater, respectively, than the estimates in blocks II and III. We were not able to come up with a reason for these differences, as soil conditions and plant emergence rate were similar among blocks.

Because no previous reports on LAR and phyllochron in safflower was found in the literature, we can only make comparisons of the values obtained in this study with other species. Moreover, comparisons of growth and developmental parameters based on thermal time need to be done at the same Tb. In summer field crops with a Tb=10°C, the phyllochron in maize, sorghum, and soybean were 45.2°C days leaf<sup>-1</sup> (WARRINGTON & KANEMASU, 1983), 51.7°C days leaf<sup>-1</sup> (MAJOR et al., 1990), and 55.5°C days leaf<sup>-1</sup> (SINCLAIR, 1986), respectively. Using a Tb=10°C in safflower, the phyllochron in this study is 15.7°C days leaf<sup>-1</sup>, indicating that leaves in safflower appear faster than in the field crops.

The estimation of LAR and phyllochron is a valuable step towards the development of a model for predicting leaf number and leaf area in safflower. We are currently working this research goal.

#### ACKNOWLEDGEMENTS

Rocha, sponsored by Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES). Schuh, sponsored by FIPE, UFSM.

#### REFERENCES

- AMIR, J.; SINCLAIR, T.R. A model of water limitation on spring wheat growth and yield. **Field Crops Research**, Amsterdam, v.29, n.1, p.59-96, 1991.
- ARNOLD, C.Y. Maximum-minimum temperatures as a basis for computing heat units. **Proceedings of the American Society for Horticultural Sciences**, Boston, v.76, n.1, p.682-692, 1960.
- BATISTA, M.P. **Estudo exploratório do mercado de flores e plantas ornamentais**. 2000. 54f. Monografia (Especialização em Educação Ambiental) - Universidade Federal de Santa Maria.
- BROOKING, I.R. et al. The influence of daylength on final leaf number in spring wheat. **Field Crops Research**, Amsterdam, v.41, p.155-165, 1995.
- CALDERINI, D.F. et al. Appearance and growth of individual leaves as affected by semidwarfism in isogenic lines of wheat. **Annals of Botany**, London, v.77, p.583-589, 1996.
- FAOSTAT. **Safflower**. Rome: FAO, 2004. Available at: [www.fao.org/ag/agl/aglw/cropwater/safflower](http://www.fao.org/ag/agl/aglw/cropwater/safflower). On-line. Captured on 15 July 2004.
- GAYETTO, O. et al. Comportamento de cultivares de cartamo (*Carthamus tinctorius* L.) em la region de Rio Cuarto, Córdoba (Argentina). **Revista de Investigación Agraria, Produccion y Proteccion Vegetales**, v.14, p.1-2, 1999.
- GILMORE, E.C. Jr.; ROGERS, J.S. Heat units as a method of measuring maturity in corn. **Agronomy Journal**, Madison, v.50, n.10, p.611-615, 1958.
- HODGES, T.; RITCHIE, J.T. The CERES-Wheat phenological model. In: HODGES, T. **Predicting crop phenology**. Boston: CRC, 1991. p.133-143.
- JAMIESON, P.D. et al. Making sense of wheat development: a critique of methodology. **Field Crops Research**, Amsterdam, v.55, p.117-127, 1998.
- KIRBY, E.J.M. Environmental factors influencing the phyllochron. **Crop Science**, Madison, v.35, n.1, p.11-19, 1995.
- KLEPPER, B. et al. Quantitative characterization of vegetative development in small cereal grains. **Agronomy Journal**, Madison, v.7, p.780-792, 1982.
- McMASTER, G.S.; SMIKA, D.E. Estimation and evaluation of winter wheat phenology in the central Great Plains. **Agricultural and Forest Meteorology**, Amsterdam, v.43, n.1, p.1-18, 1988.
- MAJOR, D.J. et al. Temperature and photoperiod effects mediated by the sorghum maturity genes. **Crop Science**, Madison, v.30, p.305-310, 1990.
- McMASTER, G.S. et al. Simulation of shoot vegetative development and growth of unstressed winter wheat. **Ecological Modelling**, Amsterdam, v.53, p.189-204, 1991.
- MULLER FLOWERSEEDS. **Variety and price list for professional flower growers**. Holland: Lisse, 1998. 59p.
- OELKE, E. A. et al. **Safflower**. Alternative field crops Manual. Wisconsin: Cooperative Extension. Available at <http://www.hort.purdue.edu/newcrop/afcm/safflower.html>. On-line. Captured on 15 July 2004.
- POLUNIN, O. **Guía de campo de las flores de Europa**. Barcelona: Omega, 1991. p.538-541.
- RICKMAN, R.W.; KLEPPER, B.L. Tillering in wheat. In: HODGES T. (ed). **Predicting crop phenology**. Boston: CRC, 1991. p.73-83.
- ROBERTSON, M.J. et al. Temperature response to vernalization in wheat: modelling the effect on the final number of main stem leaves. **Annals of Botany**, London, v.78, p.371-381, 1996.
- RUSSELE, M.P. et al. Growth analysis based on degree days. **Crop Science**, Madison, v.24, n.1, p.28-32, 1984.
- SINCLAIR, T.R. Water and nitrogen limitations in soybean grain production. I: Model development. **Field Crops Research**, v.15, p.125-141, 1986.
- STEVENS, A.B. **Field grown cut flowers: a practical guide and sourcebook; commercial field grown; fresh and dried cut flower production**. 2.ed. New York: Avatar's World, 1998. 48p.
- STRECK, N.A. et al. Incorporating a chronology response function into the prediction of leaf appearance rate in winter wheat. **Annals of Botany**, London, v.92, n.2, p.181-190, 2003.
- WARRINGTON, I.J.; KANEMASU, E.T. Corn growth response to temperature and photoperiod. II: Leaf initiation and leaf appearance rate. **Agronomy Journal**, Madison, v.75, n.5, p.755-761, 1983.