



Ciência Rural

ISSN: 0103-8478

cienciarural@mail.ufsm.br

Universidade Federal de Santa Maria  
Brasil

Trevisan Gressler, Lucas; Quinto Pereira, Larissa; Brustolin, Joice Magali; Lovato, Maristela; Gonzalez Monteiro, Sílvia  
Occurrence of *Amblyomma longirostre* in *Cyanocompsa brissonii* in Rio Grande do Sul, Brazil  
Ciência Rural, vol. 46, núm. 4, abril, 2016, pp. 660-662  
Universidade Federal de Santa Maria  
Santa Maria, Brasil

Available in: <http://www.redalyc.org/articulo.oa?id=33144324015>

- How to cite
- Complete issue
- More information about this article
- Journal's homepage in redalyc.org

redalyc.org

Scientific Information System

Network of Scientific Journals from Latin America, the Caribbean, Spain and Portugal

Non-profit academic project, developed under the open access initiative

## Occurrence of *Amblyomma longirostre* in *Cyanocompsa brissonii* in Rio Grande do Sul, Brazil

### Ocorrência de *Amblyomma longirostre* em *Cyanocompsa brissonii* no Rio Grande do Sul, Brasil

Lucas Trevisan Gressler<sup>I</sup> Larissa Quinto Pereira<sup>II</sup>  
Joice Magali Brustolin<sup>III</sup> Maristela Lovato<sup>IV</sup> Sílvia Gonzalez Monteiro<sup>V</sup>

#### – NOTE –

#### ABSTRACT

Ticks are arthropods that are highly competent in transmitting pathogens to animals and humans. Among these, the genus *Amblyomma* is the most representative within the Neotropics. *Amblyomma longirostre* ticks are naturally distributed in countries of South, Central and North America. Their immature stages preferentially parasitize birds (Passeriformes), while adult stages are usually found on rodents. Therefore, reports of this tick species on wild hosts is epidemiologically relevant, especially because of these ticks' potential for transmitting pathogens to other wild and domestic animals, and also to humans. Thus, the aim of this study was to report infestation by *Amblyomma longirostre* on *Cyanocompsa brissonii* in southern Brazil.

**Key words:** ticks, *Ixodidae*, parasitism, birds.

#### RESUMO

Os carrapatos são artrópodes que apresentam elevada competência na transmissão de patógenos para animais e humanos. Entre esses, o gênero *Amblyomma* é o mais representativo dentro da região neotropical. Ixodídeos como *Amblyomma longirostre* estão distribuídos em países da América do Sul, Central e do Norte. Os estágios imaturos desta espécie parasitam preferencialmente aves (Passeriformes) e estágios adultos são encontrados principalmente em roedores. Logo, o registro de espécies de carrapatos em hospedeiros silvestres é epidemiologicamente relevante, devido ao potencial de transmissão dos patógenos a outros animais silvestres, domésticos e ao homem. Assim, o objetivo deste estudo foi relatar o parasitismo por *Amblyomma longirostre* em *Cyanocompsa brissonii* no sul do Brasil.

**Palavras-chave:** carrapatos, *Ixodidae*, parasitismo, aves.

The genus *Amblyomma* (Koch, 1844) is of medical and veterinarian importance as a transmitter of pathogens. It is considered the most representative genus within the Neotropical region and at least 64 species have been described, including *Amblyomma longirostre* (NAVA et al., 2014). The larvae of this species are found exclusively in birds, while nymphs are generally reported parasitizing passerines but on rare occasions have been reported on mammals. Conversely, the adult stages are preferentially found on rodent hosts of the genera *Coendou*, *Chaetomys* and *Sphiggurus* (*Erethizontidae*) (NAVA et al. 2010).

The ultramarine grosbeak (*Cyanocompsa brissonii*) is a passerine in the *Cardinalidae* family. It is reported in at least seven countries in South America. It occurs very commonly and since 1988 it has been classified by the International Union for Conservation of Nature (IUCN) in the category of "least concern" (BIRDLIFE, 2014). In the state of Rio Grande do Sul, Brazil, it inhabits regions of native grassland and areas with trees and bushes. It can also be reported on the edges of agricultural production zones (BENCKE, 2001).

After a storm, an adult male specimen of *C. brissonii* was found inside a house in the district of Boca do Monte, in the city of Santa Maria, RS (29°38'32.25" S, 53°55'44.62" W). This region is

<sup>I</sup>Programa de Pós-graduação em Medicina Veterinária (PPGMV), Universidade Federal de Santa Maria (UFSM), Santa Maria, RS, Brasil.

<sup>II</sup>Programa de Pós-graduação em Medicina Veterinária (PPGMV), Laboratório de Ornitopatologia, Universidade Estadual Paulista (UNESP), Botucatu, SP, Brasil.

<sup>III</sup>Programa de Pós-graduação em Veterinária, Universidade Federal de Pelotas (UFPEL), Pelotas, RS, Brasil.

<sup>IV</sup>Laboratório Central de Diagnóstico em Patologias Aviárias, Universidade Federal de Santa Maria (UFSM), Santa Maria, RS, Brasil.

<sup>V</sup>Departamento de Microbiologia e Parasitologia, Universidade Federal de Santa Maria (UFSM), 97105-900, Santa Maria, RS, Brasil.

E-mail: sgmonteiro@uol.com.br. Corresponding author.

characterized as forming part of the Atlantic Forest biome, with the presence of semi-deciduous forest. For this bird to be taken out of the house, it was caught with the aid of towels and physically restrained. On making a visual inspection, the presence of a parasitic tick was observed. This was removed with the aid of cotton wool soaked in 70° alcohol. The tick was identified with the aid of an Olympus stereoscopic microscope (series CX40) and the dichotomous key of MARTINS et al. (2010). The specimen presented absence of a genital opening, triangular base of the capitulum, lanceolate hypostome and elongated scutum with unornamented rugose surface, and was thus identified and classified as a nymph of *A. longirostre*. According to OGRZEWSKA et al. (2010), nymphs of *A. longirostre* can easily be identified because of their sharp-pointed lance-shaped hypostome, along with their elongated scutum. The tick collected in this study was deposited in the National Tick Collection (CNC) of the School of Veterinary Medicine and Zootechnics of the Universidade de São Paulo, SP, Brazil (access number: CNC-3054).

In a study conducted by NAVA et al. (2010), larval, nymph and adult stages of *A. longirostre* from different hosts such as birds and mammals (including humans) were catalogued. According to epidemiological studies, birds tend to be hosts for transporting *A. longirostre*, thereby helping to disperse the immature phases (larvae and nymphs), but only rarely the adult form (STORNI et al., 2005). Among the immature stages of *A. longirostre*, although larvae may often infest birds, reports of parasitism at this stage are less frequent because of difficulty in identifying the larvae. ARZUA et al. (2005) reported on a nymph stage of *A. longirostre* in *C. brissonii* in the state of Minas Gerais, Brazil. These authors also described bird species parasitized by ticks of the species *A. longirostre* in the state of Rio Grande do Sul (Table 1), and this has been one of the most significant studies so far. These findings reiterate that birds are important hosts for the immature stages of this species and have high potential for dispersing ticks and the pathogens that they transmit, such as bacteria of the genus *Rickettsia*.

According to LABRUNA et al. (2004), two adult specimens of *A. longirostre* that were collected from *Coendu prehensilis* (Linnaeus, 1758) in the state of Rondônia, Brazil, were positive for a strain of *Rickettsia* belonging to the macular fever group. This was then named “aranha” (spider) strain. OGRZEWSKA et al. (2008) reported that immature stages (nymphs) of *A. longirostre* removed from birds were infected with *Rickettsia*. Their isolate

Table 1 - Hosts of *A. longirostre* in the state of Rio Grande do Sul.

Host	----Stage----			Reference
	L	N	A	
-----Mammals-----				
<i>Coendou villosus</i>			2	Oliveira et al., 1997 apud SOARES et al., 2009; BRUM et al., 2003
-----Birds-----				
<i>Lathrotriccus euleri</i>	1			ARZUA et al., 2005
<i>Pipraeidea melanonota</i>	1			ARZUA et al., 2005
<i>Saltator similis</i>	3			ARZUA et al., 2005
<i>Synallaxis spixi</i>	1			ARZUA et al., 2005
<i>Tachyphonus coronatus</i>	1			ARZUA et al., 2005
<i>Turdus amaurocalinus</i>	1			ARZUA et al., 2005
<i>Turdus subalaris</i>	1			ARZUA et al., 2005
<i>Cyanocompsa brissonii</i>	1			Present study
<i>Ramphastos dicolorus</i>	NS			SOARES et al., 2009

\*NS = not stated.

was named the AL strain. According to MARTINS et al. (2004), identifying and recording tick species in wild hosts is epidemiologically important because of their potential for transmitting pathogens to other wild, and domestic animals and humans.

According to TORGA et al. (2013), records of tick species have recently been made among forest birds in the Araucaria, Atlantic Forest, Amazon and Cerrado regions and in northeastern Brazil. However, there is a scarcity of studies relating to characterization of parasitism among birds or among mammals, caused by *A. longirostre* or even by other species of *Amblyomma* in the ecosystems of the state of Rio Grande do Sul. The geographical distribution of *A. longirostre* includes countries in North, Central and South America. In Brazil, it occurs in all five macroregions (north, northeast, center-west, southeast and south) (SOARES et al., 2009). Table 1 shows the records of this species and its respective hosts in the state of Rio Grande do Sul up to the present date.

Factors such as constant changes to and desecration of ecosystems, undertaken in order to introduce livestock-rearing and, more intensively, agriculture and forestry, have reduced the size of natural areas. This process has increased the degree of contact between humans and wild animals and consequently their contact with very many parasites, including ticks (TORGA et al., 2013). This finding highlights the widespread distribution of species of *Amblyomma* and their parasitism in a great

diversity of wild hosts. This is the first report of *A. longirostre* in a bird of the family *Cardinalidae* in the state of Rio Grande do Sul. The present study is of importance for future epidemiological and ecological investigations and because of the implications of parasitism by this species among members of this family or even in other passerines.

## ACKNOWLEDGEMENT

Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq).

To Dr. Marcelo Bahia Labruna for assistance in identifying the tick.

## REFERENCES

- ARZUA, M.A. et al. Catalogue of the tick collection (Acari: *Ixodidae*) of the Museu de História Natural Capão da Imbuia, Curitiba, Paraná, Brazil. *Revista Brasileira de Zoologia*, v.22, n.3, p.623-632, 2005. Available from: <<http://www.scielo.br/pdf/rbzool/v22n3/26178.pdf>>. Accessed: Nov. 30, 2014. doi: 10.1590/S0101-81752005000300015.
- BENCKE, G.A. **Lista de referência das aves do Rio Grande do Sul**. Porto Alegre: Fundação Zoobotânica do Rio Grande do Sul, 2001. 104p.
- BirdLife International (2014) Species factsheet: *Cyanocompsa brissonii*. Available from: <<http://www.birdlife.org>>. Accessed: Nov. 30, 2014.
- LABRUNA, M.B. et al. Molecular evidence for a spotted fever group *Rickettsia* species in the tick *Amblyomma longirostre* in Brazil. *Journal of Medical Entomology*, v.41, n.3, p.533-537, 2004. Available from: <<http://www.bioone.org/doi/full/10.1603/0022-2585-41.3.533>>. Accessed: Nov. 30, 2014. doi: 10.1603/0022-2585-41.3.533.
- MARTINS, J.R. et al. Occurrence of ticks on giant anteater (*Myrmecophaga tridactyla*) and collared anteater (*Tamandua tetradactyla*) in the pantanal region of Mato Grosso do Sul State, Brazil. *Ciência Rural*, v.34, n.1, p.293-295, 2004. Available from: <[http://www.scielo.br/scielo.php?script=sci\\_arttext&pid=S0103-84782004000100048](http://www.scielo.br/scielo.php?script=sci_arttext&pid=S0103-84782004000100048)>. Accessed: Nov. 30, 2014. doi: 10.1590/S0103-84782004000100048.
- MARTINS, T.F. et al. Nymphs of the genus *Amblyomma* (Acari: *Ixodidae*) of Brazil: descriptions, redescrptions, and identification key. *Ticks and Tick-borne Diseases*, v.1, n.2, p.75-99, 2010. Available from: <<http://dx.doi.org/10.1016/j.ttbdis.2010.03.002>>. Accessed: Dec. 01, 2014. doi: 10.1016/j.ttbdis.2010.03.002.
- NAVA, S. et al. *Amblyomma hadanii* n. sp. (Acari: *Ixodidae*), a tick from northwestern Argentina previously confused with *Amblyomma coelebs* Neumann, 1899. *Systematic Parasitology*, v.88, n.3, p.261-272, 2014. Available from: <<http://www.ncbi.nlm.nih.gov/pubmed/24935128>>. Accessed: Nov. 30, 2014. doi: 10.1007/s11230-014-9500-9.
- NAVA, S. et al. First record of *Amblyomma longirostre* (Koch, 1844) (Acari: *Ixodidae*) from Peru, with a review of this tick's host relationships. *Systematic & Applied Acarology*, v.15, p.21-30, 2010. Available from: <[https://www.academia.edu/385947/The\\_first\\_record\\_of\\_Amblyomma\\_longirostre\\_Koch\\_144\\_Acari\\_Ixodidae\\_from\\_Peru\\_with\\_a\\_revision\\_of\\_its\\_host\\_range](https://www.academia.edu/385947/The_first_record_of_Amblyomma_longirostre_Koch_144_Acari_Ixodidae_from_Peru_with_a_revision_of_its_host_range)>. Accessed: Nov. 30, 2014.
- OGRZEWSKA, M. et al. Ticks (Acari: *Ixodidae*) infesting wild birds in the eastern Amazon, northern Brazil, with notes on rickettsial infection in ticks. *Parasitology Research*, v.106, p.809-816, 2010. Available from: <<http://www.ncbi.nlm.nih.gov/pubmed/20140452>>. Accessed: Apr. 05, 2015. doi: 10.1007/s00436-010-1733-1.
- OGRZEWSKA, M. et al. Ticks (Acari: *Ixodidae*) infesting wild birds in an Atlantic Forest area in the state of São Paulo, Brazil, with isolation of *Rickettsia* from the tick *Amblyomma longirostre*. *Journal of Medical Entomology*, v.45, n.4, p.770-774, 2008. Available from: <<http://www.bioone.org/doi/full/10.1603/00222585%282008%2945%5B770%3ATAIWB%5D2.0.CO%3B2>>. Accessed: Nov. 30, 2014. doi: 10.1603/0022-2585(2008)45[770:TAIWB]2.0.CO;2.
- SOARES, J.F. et al. Occurrence of *Amblyomma longirostre* in *Ramphastos dicolorus* in Southern Brazil. *Ciência Rural*, v.39, n.3, p.930-932, 2009. Available from: <[http://www.scielo.br/scielo.php?pid=S0103-84782009000300048&script=sci\\_arttext](http://www.scielo.br/scielo.php?pid=S0103-84782009000300048&script=sci_arttext)>. Accessed: Nov. 30, 2014. doi: 10.1590/S0103-84782009000300048.
- STORNI, A. et al. Feather mites and ticks (Acari) associated to *Turdus albicollis* Vieillot (Aves, *Muscicapidae*) in an area of Atlantic Forest at Ilha Grande, Rio de Janeiro, Brasil. *Revista Brasileira de Zoologia*, v.22, n.2, p.419-423, 2005. Available from: <[http://www.scielo.br/scielo.php?script=sci\\_arttext&pid=S0101-81752005000200017](http://www.scielo.br/scielo.php?script=sci_arttext&pid=S0101-81752005000200017)>. Accessed: Nov. 30, 2014. doi: 10.1590/S0101-81752005000200017.
- TORGA, K. et al. Ticks on birds from Cerrado forest patches along the Uberabinha river in the Triângulo Mineiro region of Minas Gerais, Brazil. *Ciência Rural*, v.43, n.10, p.1852-1857, 2013. Available from: <[http://www.scielo.br/scielo.php?pid=S0103-84782013001000019&script=sci\\_arttext](http://www.scielo.br/scielo.php?pid=S0103-84782013001000019&script=sci_arttext)>. Accessed: Dec. 01, 2014. doi: 10.1590/S0103-84782013005000121.