



Dementia & Neuropsychologia

ISSN: 1980-5764

demneuropsychy@uol.com.br

Associação Neurologia Cognitiva e do
Comportamento
Brasil

Moschetta, André Luiz; Volkweis Silveira, Carine; Rigo Dalacorte, Roberta; Herberto Schneider,
Rodolfo; da Silva Filho, Irênio Gomes

Time of delirium onset and prognosis amongst Southern Brazilian hospitalized elderly patients

Dementia & Neuropsychologia, vol. 3, núm. 4, octubre-diciembre, 2009, pp. 303-307

Associação Neurologia Cognitiva e do Comportamento

São Paulo, Brasil

Available in: <http://www.redalyc.org/articulo.oa?id=339529015007>

- How to cite
- Complete issue
- More information about this article
- Journal's homepage in redalyc.org

redalyc.org

Scientific Information System

Network of Scientific Journals from Latin America, the Caribbean, Spain and Portugal

Non-profit academic project, developed under the open access initiative

Time of *delirium* onset and prognosis amongst Southern Brazilian hospitalized elderly patients

André Luiz Moschetta¹, Carine Volkweis Silveira¹, Roberta Rigo Dalacorte²,
Rodolfo Herberto Schneider², Irênio Gomes da Silva Filho²

Abstract – The prognostic significance of *delirium* in hospitalized elderly has not yet been fully clarified. **Objectives:** The present study was designed to evaluate the relationship between prevalent *delirium* (PrD), incident *delirium* (InD) and final outcome. **Methods:** A historical cohort of 261 patients was selected. *delirium* was diagnosed using the Confusion Assessment Method. **Results:** The total frequency of *delirium* detected was 42.5%-31.4% PrD and 16.2% InD. Among patients with InD, the average length of hospital stay was 9.1 days longer than for patients without *delirium* ($p=0.002$), and the hospital mortality associated with InD was 48% versus 2.7% for those without *delirium* ($p<0.001$). However, no difference was observed between patients with PrD and those without *delirium*. **Conclusions:** These results suggest that, when investigating *delirium* and prognosis amongst hospitalized elderly, it is fundamental to differentiate in terms of time of onset. Furthermore, the absence of *delirium* seems to be an important protective factor.

Key words: *delirium*, acute confusional state, elderly, prognostic, hospitalization.

Momento de início do *delirium* e prognóstico em pacientes idosos hospitalizados no sul do Brasil

Resumo – O significado do prognóstico de *delirium* em idosos hospitalizados ainda não está completamente elucidado. **Objetivos:** O presente estudo foi designado para avaliar a relação entre *delirium* prevalente (DeP), *delirium* incidente (DeI) e o desfecho final. **Métodos:** Uma coorte histórica de 261 pacientes foi selecionada. *delirium* foi diagnosticado pelo *Confusion Assessment Method*. **Resultados:** A frequência total de *delirium* foi de 42.5%, DeP 31.4%, e DeI 16.2%. Para pacientes com DeI, a média de duração de hospitalização foi 9.1 dias maior do que aqueles sem *delirium* ($p=0.002$), e a mortalidade hospitalar associada a DeI foi de 48.3% contra 2.7% dos livres de *delirium* ($p<0.001$). Contudo, não houve diferença entre pacientes com DeP e sem *delirium*. **Conclusões:** Os resultados sugerem que, ao estudar *delirium* e prognóstico entre idosos hospitalizados, é fundamental diferenciar o problema quanto ao seu momento de início. Além disso, não apresentar *delirium* parece constituir-se num fator protetor importante.

Palavras-chave: *delirium*, estado confusional agudo, idoso, prognóstico, hospitalização.

Individuals aged 60 years or older make up approximately 12% of the population in the southern region of Brazil¹ and represent about 20% of all patients admitted to the regions' general hospitals.²⁻⁴ This patient group uses the hospital services more intensively than those of other age groups because their treatment and recuperation takes longer and costs more.⁴ *delirium*, or acute confusional state, is an organic mental syndrome which is acute, transient and fluctuating, that occurs frequently in hospitalized elderly

people,^{5,6} and is considered the most common hospital complication in this population.⁷⁻⁹

Irrespective of the reason for hospitalization, it is estimated that 15 to 20% of the elderly will be in an acute confusional state on admission, and 25 to 55% will develop the condition during their hospital stay.^{7,10} This frequency depends on the characteristics of the patient and their diseases, on the care dispensed and the sensitivity of the method used for diagnosis.¹¹ The link between *delirium*

¹MD, Institute of Geriatrics and Gerontology, Pontifical Catholic University, Porto Alegre RS, Brazil. ²MD, PhD, Institute of Geriatrics and Gerontology, Pontifical Catholic University, Porto Alegre RS, Brazil.

André L. Moschetta – Instituto de Geriatria e Gerontologia - Av. Ipiranga 6690 / 3º andar - 90610-000 Porto Alegre RS - Brazil. E-mail: andremoschetta@gmail.com

Disclosure: The authors report no conflicts of interest.

Received September 29, 2009. Accepted in final form November 16, 2009.

and severity of acute morbid states is well known.¹² Studies on patients hospitalized in intensive care units, or those suffering from terminal diseases, have demonstrated incidences of the syndrome up to 80%.^{10,12,13}

The biological mechanisms involved in the genesis of *delirium* are not yet fully understood. It is known that a complex relationship exists between individual vulnerability and aggressor factors, often multiple, making it more difficult to determine the exact role *delirium* plays in the disease process. Consequently, the data in the literature on the prognostic implications of *delirium* remain controversial. Although various investigative studies, apparently well conducted, have demonstrated an association between this acute confusional state and hospital mortality,^{11,14} prolonged hospital stay,^{15,16} functional decline and greater rates of discharge to places other than the home,^{6,7,17,18} other studies of the same quality have found no such association.¹⁹⁻²³ It is possible that one of the principal reasons for this difference in results is the failure to distinguish between prevalent *delirium* (PrD) and incident *delirium* (InD) where this likely leads to the different prognostics.^{24,25} In fact, none of the studies cited above evaluated the differences between the types of *delirium* with respect to the time of onset.

An intensive bibliographic search of the MEDLINE and SCIELO databases, and the respective reference textbooks identified only one paper in which the prognosis of the two presentations of *delirium* were studied separately. In a study of a cohort of 359 patients, McCusker et al. compared the duration of hospital stay amongst elderly patients with PrD versus those with InD, and those without either *delirium*. They found an association of the duration of hospital stay and *delirium* only in the group of patients with InD.²⁴ This finding reinforces the hypothesis that PrD and InD contribute differently to prognosis.

The present study was conducted in view of the lack of consensus over the effect on prognosis of *delirium* in elderly patients, the scarcity of data differentiating the time of onset, and the scant information available on its occurrence in our population. The objective was to describe and compare the duration of hospital stay and mortality in elderly patients with PrD versus those with InD, and without either *delirium*, who had been hospitalized in the geriatric ward of a general hospital in the south of Brazil.

Methods

The study was conducted in a geriatric ward with 16 beds – 8 for use by the Brazilian public health system (*Sistema Único de Saúde - SUS*) and 8 for private patients – in a university general hospital in the south of Brazil. A historical cohort was constructed based on the hospital patient records of all those aged 60 years or older who

had been hospitalized consecutively during 2007. We excluded all individuals who had stayed more than 24 hours in another hospital sector (n=102), those that had been discharged for transfer or non-medical reasons (n=3), and those that had been discharged after less than 24 hours of hospitalization (n=2).

These patient files are completed by the medical team of the Geriatric Service, all of whom receive frequent training in how to detect *delirium*. The diagnoses were reached using the Confusion Assessment Method (CAM) which is accurate, precise and valid for this purpose.^{26,27} The criteria used for definition of the cases were: a report of “*delirium*”, “acute confusional state” or “acute confusional state with organic origin”, or the presence of other information meeting the CAM criteria. Prevalent *delirium* was defined by a diagnosis of *delirium* in the first 24 hours of admission, and InD when *delirium* was detected after this period. Data was also collected on the demographic details (sex, age and health plan), the cause and characteristics of hospitalization (acute or elective; duration of hospital stay) and on the basic cause of the hospital obituary.

The mean hospital stay and the occurrence of death in the groups with PrD, and without *delirium* on admission, as well as the groups with InD and without *delirium*, were compared. To study the duration of hospital stay, patients who died were excluded from the analysis because we believe the relationship between hospital stay and mortality in these cases may be inverted, if the more seriously ill patients died more rapidly.

The respective research project for this study was approved by the Research Ethics Committee of the São Lucas Hospital at the Pontifical Catholic University of Rio Grande do Sul State. All regulations concerning patient privacy and the handling of the manuscripts were fully respected. In view of the fact that this research was conducted entirely from the patients' hospital files, informed consent forms were deemed unnecessary.

To compare the frequencies of PrD and InD according to demographic and clinical variations, and mortality, the basic chi-squared test was used, while for the age groups the chi-squared trend test was applied. In the analysis of mortality, Fisher's exact test was used whenever an expected value of less than five was found. The comparison of mean hospitalization times of the groups with and without *delirium*, either prevalent or incident, was performed using Student's *t* test. The significance level was assumed as 5%. The data was analyzed with the SPSS software package version 11.5.

Results

Table 1 lists the demographic and clinical data of the

261 patients analyzed in this study. Patient ages ranged from 60 to 104 years, having a mean of 79.9 years (SD=8.4). The predominant characteristics were female gender (58.4%), more than 80 years old (54.8%), public health system patients (53.3%), and suffering from acute disease (72.5%). The most frequent reasons for hospitalization were elective clinical indication (22.2%) followed by respiratory infections (12.3%). The mean length of hospital stay was 13.2 days (SD=11.2), and 32 patients (12.3%) evolved to hospital death.

The total incidence of *delirium* was 42.5%, with PrD occurring in 31.4%, and InD in 16.2%. The incidence of PrD differed according to age with a tendency to rise with increasing age and to be higher in the group hospitalized in the private patients' beds. (36.9% v. 26.6%, $p=0.075$). The proportion of PrD in patients hospitalized for acute conditions was higher (42.6% v. 1.4%, $p<0.001$). With regard to the reason for hospitalization, PrD was more frequent

in cases of urinary infection (73.9%), collateral medication effect (71.4%), acute abdominal surgery (57.1%) and respiratory infection (53.6%). No statistically significant difference in InD frequency was found that could be traced to the demographic and/or clinical characteristics.

The duration of hospital stay and mortality rate was the same in patients with and without PrD (Table 2). In patients with InD, however, the mean length of hospital stay was 9.1 days, and mortality was 48.3%, significantly greater than in the individuals without *delirium* who presented a mean hospital stay of 1.2 days ($p=0.002$) and a mortality rate of 2.7% ($p<0.001$).

Discussion

Research on *delirium* is subject to innumerable methodological problems due to the lack of clarity surrounding its physiopathology and the lack of consensus with respect to the criteria for diagnosis. Notwithstanding these issues, the

Table 1. Time of *delirium* onset, demographic and clinical characteristics.

	N (%)	Prevalent <i>delirium</i> *			Incident <i>delirium</i>		
		N	%	p [†]	N	%	p
Sex				0.202			0.681
Male	106 (40.6)	38	35.8		12	17.6	
Female	155 (59.4)	44	28.4		17	15.3	
Age group				0.020			0.369
60-69	28 (10.7)	4	14.3		4	16.7	
70-79	90 (34.5)	26	28.9		7	10.9	
80 or older	143 (54.8)	52	36.4		18	19.8	
Health Plan				0.075			0.532
Public	139 (53.3)	37	26.6		15	14.7	
Private	122 (46.7)	45	36.9		14	18.2	
Type of hospitalization				<0.001			0.577
Acute	190 (72.8)	81	42.6		19	17.4	
Elective	71 (27.2)	1	1.4		10	14.3	
Reason for hospitalization				<0.001			0.318
Elective clinical [‡]	58 (22.2)	1	1.7		7	12.3	
Respiratory infection	56 (21.5)	30	53.6		5	19.2	
Urinary infection	23 (8.8)	17	73.9		3	50.0	
Other infection	19 (7.3)	3	15.8		4	25.0	
Acute renal insufficiency	15 (5.7)	7	46.7		1	12.5	
Heart disease [§]	13 (5.0)	2	15.4		0	0.0	
Elective surgery	13 (5.0)	0	0.0		3	23.1	
Collateral medication effect	7 (2.7)	5	71.4		0	0.0	
Acute abdominal	7 (2.7)	4	57.1		0	0.0	
Other	50 (19.2)	13	26.0		6	16.2	
Total	261 (100)	82	31.4	--	29	16.2	--

*Prevalent *delirium*: proportion of *delirium* diagnosed within the first 24 hours of Hospital admission. Incident *delirium*: proportion of *delirium* diagnosed during hospital stay in patients who were *delirium*-free on arrival; [†]Values obtained from the χ^2 test. For the variable age, the χ^2 trend test was used; [‡]Includes: non-surgical diagnostic investigation, therapeutic organization, or family exhaustion; [§]Includes: acute decompensated heart failure, cardiac arrhythmia, acute coronary syndrome, or stroke.

Table 2. Time of *delirium* onset, duration of hospital stay and mortality.

	Prevalent <i>delirium</i>		p
	Yes	No	
N	82	179	
Duration of hospital stay in days, of elderly patients who were discharged (mean±SD)	12.3±9.9	13.1±11.0	0.602*
Mortality N (%)	14 (17.1)	18 (10.1)	0.109†
	Incident <i>delirium</i>		p
	Yes	No	
N	29	150	
Duration of hospital stay in days, of elderly patients who were discharged (mean±SD)	21.3±11.0	12.2±10.7	0.002*
Mortality N (%)	14 (48.3)	4 (2.7)	<0.001‡

*t test; †Chi-squared test; ‡Fisher's exact test.

results of the present study as regards the frequency of the incidence of *delirium*, and the association of PrD with older age, acute care hospitalization, and the reasons for hospitalization all agree with the data in the literature.^{7,15,17,22,28} The greater prevalence of *delirium* found in patients with urinary infections and collateral medication effects can be explained by the fact that the signs and symptoms involved with these conditions are less apparent amongst the elderly, delaying diagnosis and increasing the chances of developing acute confusion states before treatment is initiated.

We found no statistically significant association between PrD and prognosis in spite of the greater mortality in the group that presented *delirium* on admission. It is possible that our sample was too small to demonstrate any difference. Also, we excluded those patients who had been spent more than 24 hours outside the geriatric ward where these patients would generally be in a more serious condition, and would probably have presented with *delirium* on admission and have a poorer diagnosis. *delirium* that initiated during hospitalization, however, was strongly associated with both the duration of hospital stay and mortality. This difference between PrD and InD was first described by Mc Cusker et al., who found longer periods of hospitalization in patients with InD and no association of PrD with prognosis.²⁴

The difference found in the prognostic values of PrD and InD suggests that PrD is not necessarily an indicator of serious illness, and may be more closely related to previous vulnerable states (for instance: advanced age, cognitive impairment, institutionalization, fragility) which facilitate the onset of a confusional state. This neurological fragility likely influences the outcome but to a lesser extent than the degree of severity of acute disease. On the other hand, patients that developed InD, but were admitted to the hospital problem-free generally demonstrated less vulner-

ability. The occurrence of InD in these patients would indicate, therefore, negative evolution of the conditions that led to hospitalization in the first place, or occurrence of a hospital-acquired complication.

Clinical, epidemiological and basic scientific evidence supports the hypothesis that *delirium* is merely an indicator of prognosis and is not itself a cause of a poor prognosis.^{11,18,29} The significance of *delirium* may lie in its ability to reveal a deleterious interaction between the individual and the aggressor factor. In some cases, increased individual vulnerability could lead to a greater occurrence of PrD, less relevant for the prognosis. In other cases, because of more aggressive disease, which is generally the cause of the appearance of InD, the prognosis is worse. Considering the greater prevalence of PrD (31.4% v. 16.2%), it seems that increased individual vulnerability is more frequent in elderly patients with acute disease. This is clearly evident in cases of *delirium* due to benign problems, such as fecal impaction or mild dehydration which, in the majority of cases, would not justify hospitalization, much less lead to a prolonged hospital stay or a greater rate of mortality.

The mortality among the *delirium*-free group was much lower in comparison to patients who presented *delirium* (2.7% v. 25%) at some point, and this result is in line with findings of other studies.^{11,17} Therefore, the absence of *delirium* appears to indicate some degree of prognostic protection rather than the opposite. This finding may help to reach decisions in complicated situations where there is doubt over indicating invasive interventions in hospitalized elderly patients. The absence of acute confusional state in the patient's file would thus indicate a good prognosis, or less individual vulnerability, and consequently, lower risk of unfavorable evolution.

First recognizing that amongst elderly hospitalized in

geriatric wards, PrD is more related to an increased state of individual vulnerability (which has little effect on the prognosis), and that secondly, the occurrence of InD is most likely due to a serious situation (indicating evolution to worsened state), can help to clarify the role played by *delirium* as an indicator in prognosis – although this matter remains controversial. We believe, however, that it is fundamental to differentiate the time of *delirium* onset. In addition, the absence of *delirium* seems to be an important protective factor that could assist decision making regarding diagnostic and therapeutic procedures in hospitalized elderly patients. However, this hypothesis requires further elucidation.

Acknowledgments – In closing, we would like to thank Professor Dr. José Luiz da Costa Vieira for his support during the conducting of this study. The authors declare that they have no potential conflicts of interest pertaining to the present study.

References

1. Projeções e estimativas demográficas (2001-2006). Instituto Brasileiro de Geografia e Estatística (IBGE) <http://www.ibge.gov.br> Access date April 2009.
2. Castro M. A utilização das internações hospitalares no Brasil: fatores associados, grandes usuários, reinternações e efeito da oferta de serviços sobre o uso. Rio de Janeiro: Escola Nacional de Saúde Pública da Fundação Oswaldo Cruz; 2004:101.
3. Meneghell S, Armani T, Rosa R et al. Internações hospitalares no Rio Grande do Sul. Informe Epidemiológico do SUS 1997;6:49-59.
4. Loyola Filho A, Matos D, Giatti L, et al. Causas de internações hospitalares entre idosos brasileiros no âmbito do Sistema Único de Saúde. Epidemiologia e Serviços de Saúde 2004;13:229-238.
5. Lipowski ZJ. Delirium (acute confusional states). JAMA 1987; 258:1789-1792.
6. Fabbri R. Delirium. In: Freitas E, Py L, Cançado F, Doll J, Gorzoni M, editors. Tratado de Geriatria e Gerontologia, 2nd. Rio de Janeiro: Guanabara Koogan; 2006:325-332.
7. Agostini J, Inouye S. Delirium. In: Hazzard W, Blass J, Halter J, Ouslander J, Tinetti M, editors. Principles of Geriatric Medicine and Gerontology, 5th. New York: McGraw-Hill; 2003: 1503-1515.
8. Gillick MR, Serrell NA, Gillick LS. Adverse consequences of hospitalization in the elderly. Soc Sci Med 1982;16:1033-1038.
9. Inouye SK, Schlesinger MJ, Lydon TJ. Delirium: a symptom of how hospital care is failing older persons and a window to improve quality of hospital care. Am J Med 1999;106:565-573.
10. Inouye SK. Delirium in older persons. N Engl J Med 2006;354: 1157-1165.
11. Francis J, Martin D, Kapoor WN. A prospective study of delirium in hospitalized elderly. JAMA 1990;263:1097-1101.
12. Lawlor PG, Gagnon B, Mancini IL et al. Occurrence, causes, and outcome of delirium in patients with advanced cancer: a prospective study. Arch Intern Med 2000;160: 786-794.
13. Pisani MA, McNicoll L, Inouye SK. Cognitive impairment in the intensive care unit. Clin Chest Med 2003;24:727-737.
14. Kakuma R, du Fort GG, Arsenault L et al. Delirium in older emergency department patients discharged home: effect on survival. J Am Geriatr Soc 2003;51:443-450.
15. Inouye SK, Bogardus ST, Jr. Charpentier PA et al. A multicomponent intervention to prevent delirium in hospitalized older patients. N Engl J Med 1999;340:669-676.
16. Cole MG, Primeau FJ. Prognosis of delirium in elderly hospital patients. CMAJ 1993;149:41-46.
17. Pompei P, Foreman M, Rudberg MA, et al. Delirium in hospitalized older persons: outcomes and predictors. J Am Geriatr Soc 1994;42:809-815.
18. O'Keeffe S, Lavan J. The prognostic significance of delirium in older hospital patients. J Am Geriatr Soc 1997;45:174-178.
19. Adamis D, Treloar A, Darwiche FZ, et al. Associations of delirium with in-hospital and in 6-months mortality in elderly medical inpatients. Age Ageing 2007;36:644-649.
20. Siddiqi N, House AO, Holmes JD. Occurrence and outcome of delirium in medical in-patients: a systematic literature review. Age Ageing 2006;35:350-364.
21. Adamis D, Treloar A, Martin FC, et al. Recovery and outcome of delirium in elderly medical inpatients. Arch Gerontol Geriatr 2006;43:289-298.
22. Formiga F, Marcos E, Sole A, et al. Acute confusional syndrome in elderly patients hospitalized due to medical condition. Rev Clin Esp 2005;205:484-488.
23. Bellelli G, Frisoni GB, Turco R, et al. Delirium superimposed on dementia predicts 12-month survival in elderly patients discharged from a postacute rehabilitation facility. J Gerontol A Biol Sci Med Sci 2007;62:1306-1309.
24. McCusker J, Cole MG, Dendukuri N, et al. Does delirium increase hospital stay? J Am Geriatr Soc 2003;51:1539-1546.
25. Inouye SK. The dilemma of delirium: clinical and research controversies regarding diagnosis and evaluation of delirium in hospitalized elderly medical patients. Am J Med 1994;97: 278-288.
26. Fabbri RM, Moreira MA, Garrido R, et al. Validity and reliability of the Portuguese version of the Confusion Assessment Method (CAM) for the detection of delirium in the elderly. Arq Neuropsiquiatr 2001;59:175-179.
27. Inouye SK, van Dyck CH, Alessi CA, et al. Clarifying confusion: the confusion assessment method. A new method for detection of delirium. Ann Intern Med 1990;113:941-948.
28. Edlund A, Lundstrom M, Karlsson S, et al. Delirium in older patients admitted to general internal medicine. J Geriatr Psychiatry Neurol 2006;19:83-90.
29. Bellelli G, Morghen S, Turco R, et al. Delirium in older people: an epiphenomenon of incipient death or a separate biological process? Age Ageing 2008;37:353-354; author reply 355.