



Revista Chilena de Historia Natural

ISSN: 0716-078X

editorial@revchilhistnat.com

Sociedad de Biología de Chile

Chile

AKEMI NAKANO, MÁRCIA; OLIVEIRA DE MIRANDA, VITOR F.; RODRIGUES DE  
SOUZA, DÉBORA; FEITOSA, RODRIGO M.; MORINI, M. SANTINA C.  
Occurrence and natural history of *Myrmelachista Roger* (Formicidae: Formicinae) in the  
Atlantic Forest of southeastern Brazil  
Revista Chilena de Historia Natural, vol. 86, núm. 2, 2013, pp. 169-179  
Sociedad de Biología de Chile  
Santiago, Chile

Available in: <http://www.redalyc.org/articulo.oa?id=369944185006>

- How to cite
- Complete issue
- More information about this article
- Journal's homepage in redalyc.org

redalyc.org

Scientific Information System

Network of Scientific Journals from Latin America, the Caribbean, Spain and Portugal

Non-profit academic project, developed under the open access initiative



## RESEARCH ARTICLE

## Occurrence and natural history of *Myrmelachista* Roger (Formicidae: Formicinae) in the Atlantic Forest of southeastern Brazil

### Ocorrência e história natural de *Myrmelachista* Roger (Formicidae: Formicinae) en la Selva Atlántica del sureste brasileño

MÁRCIA AKEMI NAKANO<sup>1</sup>, VITOR F. OLIVEIRA DE MIRANDA<sup>2</sup>, DÉBORA RODRIGUES DE SOUZA<sup>1</sup>, RODRIGO M. FEITOSA<sup>3</sup> & M. SANTINA C. MORINI<sup>1,\*</sup>

<sup>1</sup>Laboratório de Mirmecologia, Universidade de Mogi das Cruzes, Postal Code 08701-970, Mogi das Cruzes, SP, Brazil

<sup>2</sup>Laboratório de Sistemática Vegetal, Universidade Estadual Paulista, Postal Code 14884-900, Jaboticabal, SP, Brazil

<sup>3</sup>Laboratório de Zoologia, Universidade Federal do Paraná, Postal Code 81531-980, Curitiba, Pr, Brazil

\*Corresponding author: mscmorini@gmail.com

#### ABSTRACT

Information on the biology of the species in *Myrmelachista* remains notably scarce. The objective of this study was to report the occurrence, analyze the influence of canopy cover on the number of nests in twigs in the litter describe behavioral characteristics of this genus. Hand sampling was performed on trees and twigs deposited in the litter. A total of 84 nests and seven species were found. *Myrmelachista ruskii* and *M. catharinae* were the most frequent and abundant species. In trees, most of the nests were located at the tips of dry twigs, and these nests were probably the source of the nests found in the litter of fragments of Atlantic Forest of southeastern Brazil. Canopy cover is associated with the number of nests that were in twigs.

**Key words:** canopy, dense ombrophilous forest, polydomy, reproduction, tree species.

#### RESUMEN

*Myrmelachista* es un género neotropical de hormigas arborícolas que nidifican en cavidades de troncos y ramas. El conocimiento sobre la biología de sus especies es escaso. Este trabajo tuvo como objetivo registrar la ocurrencia, analizar la influencia del dosel sobre el número de nidos en ramas en la cama (mezcla de hojas, ramas y tierra) que cubre la selva, y describir las características de comportamiento de *Myrmelachista*. Se realizaron recolectas manuales de arbóreas y en ramas depositadas en la mezcla anterior. Se encontraron 84 nidos y siete especies *Myrmelachista ruskii* y *M. catharinae* fueron las más frecuentes y abundantes. En las especies arbóreas la mayoría de los nidos se localizan en las extremidades secas de las ramas estos nidos fueron probablemente el origen de los nidos que se encuentran en las camas de los fragmentos de Floresta Atlántica de la región sureste brasileña. El número de nidos en ramas estaría asociado con la cobertura arbórea.

**Palabras clave:** bosque ombrófilo denso, dosel, especies arbóreas, polidomia, reproducción.

#### INTRODUCTION

The genus *Myrmelachista* is typically neotropical with 69 species described from southern Mexico to northern Argentina (Kempf 1972, Fernández & Sendoya 2004, Bolton et al. 2006, Longino 2006). The species of this genus forage on trees, nesting inside live stems (Longino 2006) and may develop mutualistic interactions with host plants, specialized domatia or feeding on extrafloral nectaries (Frederickson 2005, McNett et al. 2010) and/

or coccids (Stout 1979, Ketterl et al. 2003). Data regarding the ecology and behavior of species in the genus have been reported by Renner & Ricklefs (1998), Frederickson (2005, 2006), Frederickson & Gordon (2009), Edwards et al. (2009) and McNett et al. (2010). Information regarding the biology of certain species, specifically from Costa Rica, is summarized in the review by Longino (2006).

Despite the information mentioned above, the biology of the genus *Myrmelachista* is still considered to be poorly known in comparative

terms (Longino 2006). The objective of this study was to (1) report the occurrence of *Myrmelachista* species in Atlantic Forest remnants of southeastern Brazil, (2) analyze the association between canopy cover and the number of nests in twigs in the litter and (3) describe behavioral characteristics observed during the fieldwork.

## METHODS

### *Study sites and sampling of the species in vegetation*

Colonies were recorded and collected from the vegetation and twigs found in the litter of Atlantic Forest remnants located in southeastern São Paulo state, Brazil (Fig. 1).

### *Sampling of the species in vegetation*

The sampling was performed on a weekly basis from September 2008 to September 2009, covering all seasons of the region (Minuzzi et al. 2007), along the sunny trails. The colonies were collected from three areas of the Brazilian Atlantic forest. The sampling effort was kept constant, in terms of the number of field expeditions, length of stay ( $\geq 4$  h) and number of collectors (3). Worker ants that were exiting holes in tree trunks, or foraging until five centimeters from the hole, and tips of dry twigs on alive trees were collected by hand and were transferred to vials with 70 % ethanol. The presence of winged reproductive was recorded.

### *Sampling of the nests in twigs and canopy cover*

The colonies were collected from six areas of the Brazilian Atlantic forest. The sampling was performed on a weekly basis from September to December 2010 in areas without trails and not sunny. In these months, rainfall is more intense (Minuzzi et al. 2007), and results in higher deposit twigs in the leaf litter. In each area, six 16 m<sup>2</sup> sampling points, which were separated from

one another by 50 m, were delineated. All of the twigs of 10-30 cm in length were opened; those with colonies that were found on the litter surface were individually placed in plastic bags for subsequent identification. Twigs with *Myrmelachista* nests, workers, winged and immature individuals were counted and transferred to vials with 70 % ethanol.

The canopy cover was recorded at the center and corners of each 16 m<sup>2</sup> plot using a Nikon D80 digital camera with a fisheye lens. The images were analyzed using the Gap Light Analyzer (GLA) software, version 2.0 (Frazer et al. 1999).

### *Species identification*

The identification of the ant species was performed through comparison with specimens deposited in the reference collection of the Museum of Zoology, University of São Paulo (Museu de Zoologia da Universidade de São Paulo, MZUSP). Type specimens were deposited in the collection of the Laboratory of Myrmecology, University of Mogi das Cruzes (Universidade de Mogi das Cruzes, UMC) and in the MZUSP.

The identification of the plants where the *Myrmelachista* workers were found was performed through comparison with the collection of the Upper Tietê flora in the Herbarium Mogiense (Herbário Mogiense, HUMC) at the University of Mogi das Cruzes. For that purpose, fertile or sterile branches were collected. Type specimens were deposited in the HUMC.

### *Data analysis*

The abundances of adults and immature individuals in the twigs were compared using Mann-Whitney test. The association between canopy cover and the number of nests was determined using scatter plots and Spearman correlation. The level of significance was  $\alpha = 0.05$ .

## RESULTS

Seven *Myrmelachista* species were recorded in a total of 41 nests in 20 species of live trees




Fig. 1: Sampling sites. Fragments of the Atlantic Rainforest where the study was conducted.

Sitios de muestreo. Fragmentos de la Mata Atlántica donde se realizó el estudio.

TABLE 1

Number of nests of *Myrmelachista* species present in the different species of trees.Número de nidos de especies de *Myrmelachista* presentes en diferentes especies arbóreas.

Taxon	<i>M. ruszkii</i>	<i>M. catharinae</i>	<i>M. nodigera</i>	<i>M. arthuri</i>	<i>M. gallicola</i>	<i>Myrmelachista</i> sp. 4	<i>Myrmelachista</i> sp. 7
Anacardiaceae							
<i>Schinus terebinthifolius</i> Raddi					3		
Arecaceae							
Arecaceae sp. 1				1			
Asteraceae							
<i>Piptocarpha macropoda</i> (DC. Baker)	1						
Asteraceae sp. 1							1
Asteraceae sp. 2						1	
Cannabaceae							
<i>Trema micrantha</i> (L.) Blume	2					1	
Euphorbiaceae							
<i>Alchornea sidifolia</i> Muell. Arg.	1						
Fabaceae							
Fabaceae sp. 1	1						
Fabaceae sp. 2		1					
Lauraceae							
<i>Aniba</i> sp.1		1					
Melastomataceae							
<i>Tibouchina mutabilis</i> Cogn. (Vell.) Cogn. Cogn.		1	2	4			
Melastomataceae sp. 1		1					
Myrsinaceae							
<i>Rapanea</i> sp. 1	1	3	1		2		
Myrtaceae							
Myrtaceae sp. 1		1					
Piperaceae							
Piperaceae sp. 1	1			1			
Rubiaceae							
<i>Psychotria</i> sp. 1	1		1				
<i>Psychotria suterella</i> Muell. Arg.		1					
<i>Psychotria vellosiana</i> Benth	1						
Solanaceae							
<i>Solanum</i> sp. 1		1	1				
Malvaceae							
Malvaceae sp. 1			1				2
Total number of nests	9	10	6	6	5	2	3

TABLE 2

Reproductive winged forms of *Myrmelachista* observed over a year in the live trees. \*Data from one nest kept in the laboratory.

Formas aladas reproductivas de *Myrmelachista* observadas a lo largo de un año en árboles vivos. \*Datos de un nido mantenido en el laboratorio.

Month	<i>M. gallicola</i>	<i>M. arthuri</i> *	<i>M. sp. 7</i>		<i>M. ruskii</i>	<i>M. catharinae</i>
	♂	♂	♀	♂	♂	♂
January						
February						
March						
April			x	x		
May						
June						
July						
August		x				
September	x	x	x	x	x	
October			x	x	x	x
November		x				x
December						

(Table 2). *Myrmelachista ruskii* Forel and *M. catharinae* Mayr were found on the greatest number of tree species followed by *M. nodigera* Mayr and *M. arthuri* Forel (Fig. 2). Only *M. arthuri* nests (n = 6; Table 1) were recorded in live twigs of live trees. Nests of the other species were found only in the tips of dry twigs in live trees (nests = 35; Table 1). Surprisingly, *M. gallicola* nests were found in 14 twigs of a single *Schinus terebinthifolius* Raddi plant with dry tips. These colonies consisted of a queen, workers and immature individuals. Each tree nest observed only one species of *Myrmelachista*, but different individuals of the same tree species may be occupied by different *Myrmelachista* species, as was the case of *Rapanea* sp.1 (with four *Myrmelachista* species), *Tibouchina mutabilis* (Vell.) Cogn (three), *Trema micrantha* (L.) Blume (two), Piperaceae sp.1 (two), *Psychotria* sp.1 (two), *Solanum* sp.1 (two) and Malvaceae sp.1 (two) (Table 2).

Nests were also found in small dead trees (diameter at breast height < 15 cm). These nests belonged to the species *M. ruskii* (n = 3), *M. catharinae* (n = 3), *M. gallicola* (n = 2) and *M. nodigera* (n = 1). Two *M. arthuri* nests

were found in dead trees fallen in the trail. One nest was located in a 30 cm diameter *T. mutabilis* (Melastomataceae) trunk, and the length of the trunk region that was occupied by the colony was approximately 50 cm. This trunk was maintained in the laboratory (28 ± 2 °C; 70 ± 10 %) to determine the time of emergence of winged individuals (Table 2; Fig. 3). The other nest remained in the forest, and the tree species in which it was located could not be identified. Moreover, the presence of a single *M. nodigera* nest in a partially dry *Psychotria* sp.1 (Rubiaceae) trunk fallen in the litter was also recorded. Approximately 300 workers, several immature individuals and a single queen were collected in this nest.

The presence of winged individuals in live tree was recorded for some *Myrmelachista* species, and the reproductive period mainly occurred from August to December (Table 3). The number of winged individuals could only be counted for *Myrmelachista* sp. 7; 109 unfertilized queens and 11 males were found in one nest, while 113 unfertilized queens and only one male were found in the other nest.

A total of 33 nests, 19 of *M. ruskii* and 14 of *M. catharinae*, were collected in twigs in




Fig. 2: Total number of tree species associated with each studied *Myrmelachista* species in the Atlantic Forest of southeastern Brazil.

Número total de especies arbóreas asociadas a cada especie estudiada de *Myrmelachista* en la Floresta Atlántica del sudeste de Brasil.




Fig. 3: Breeders of (♂) de *Myrmelachista arthuri*. Scale: A y C = 1 cm, B = 1mm.

Formas reproductoras aladas (♂) de *Myrmelachista arthuri*. Escala: A y C = 1 cm, B = 1mm.



TABLE 3

Total number of nests, workers and immature individuals of *Myrmelachista* in dry twigs. The data are reported as medians with the range in parentheses.

Número total de nidos, obreras e individuos inmaduros de *Myrmelachista* en ramas secas. Los datos corresponden a las medianas, los intervalos se muestran entre paréntesis.

	<i>M. ruskii</i>	<i>M. catharinae</i>	test	P
Nests	19	14		
Workers	28 (9 - 458)	58.5 (13 - 349)	U = 112	0.44
Immature	58 (5 - 330)	55 (6 - 537)	U = 122.5	0.70

TABLE 4

Angiosperm families and species observed to associate with *Myrmelachista* species according to the literature.

Familias y especies de angiosperma observadas y asociadas con especies de *Myrmelachista* de acuerdo con la literatura.

Family	Species	Reference
Araucariaceae	<i>Araucaria angustifolia</i> (Bert.) O. Ktze	Ketterl et al. (2003)
Bignoniaceae	<i>Tabebuia</i> sp.	Kusnezov (1951)
Boraginaceae	<i>Cordia alliodora</i> (Ruiz & Pavon) Oken	Longino (2006)
	<i>Cordia hispidissima</i> A. de Candolle	Wheeler (1934)
	<i>Cordia nodosa</i> Lamark	Davidson & McKey (1993)
		Edwards et al. (2009)
		Frederickson & Gordon (2009)
Bromeliaceae		Goitía & Jaffé (2009)
		Frederickson & Gordon (2009)
	<i>Canistrum</i> sp.	Rosumek et al. (2008)
	<i>Nidularium</i> sp.	Rosumek et al. (2008)
	<i>Tillandsia</i> sp.	Benzing (1970)
	<i>Tillandsia balbisiana</i> Schultes	Wheeler (1934)
	<i>Tillandsia circinnata</i> Schlecht	Wheeler (1934)
Cecropiaceae	<i>Tillandsia valenzuelana</i> A. Richard	Wheeler (1934)
	<i>Cecropia angustifolia</i> Trécul.	Longino (2006)
	<i>Cecropia insignis</i> Liebm.	Longino (2006)
	<i>Cecropia obtusifolia</i> Bertol.	Longino (2006)
Chrysobalanaceae	<i>Licania pittieri</i> Prince	Goitía & Jaffé (2009)
Clusiaceae	<i>Calophyllum brasiliense</i> Camb.	Longino (2006)
	<i>Clusia alata</i> Planch.	Longino (2006)
	<i>Vismia</i> sp.	Longino (2006)
Elaeocarpaceae	<i>Sloanea</i> sp.	Longino (2006)
Euphorbiaceae	<i>Croton floribundus</i> (L.) Spreng.	Morini et al. (2006)
	<i>Euphorbia pulcherrima</i> Willdenow	Haber et al. (1981)

TABLE 4. Continuation

Family	Species	Reference
Fabaceae	<i>Sapium oligoneurum</i> K. Schum.	Longino (2006)
	<i>Acacia aroma</i> Gill. Ex Hook et Arn	Quirán & Martínez (2006)
	<i>Acacia caven</i> (Mol.) Mol.	Quirán & Martínez (2006)
	<i>Anadenanthera macrocarpa</i> Benth	Campos et al. (2006)
	<i>Erythrina</i> sp.	Longino (2006)
	<i>Inga</i> sp.	Goitia & Jaffé (2009)
		Longino (2006)
	<i>Lonchocarpus</i> sp.	Longino (2006)
	<i>Prosopis caldenia</i> Burkart	Quirán & Martínez (2006)
	<i>Prosopis nigra</i> (Griseb.)	Quirán & Martínez (2006)
Fagaceae	<i>Prosopis ruscifolia</i> (Vinal)	Kusnezov (1951)
	<i>Nothofagus antarctica</i> (G.Forster) Oerst	Kusnezov (1951)
Lauraceae	<i>Licaria</i> sp.	Hanson & Longino (2006)
		Longino (2006)
	<i>Ocotea</i> spp.	Stout (1979)
		Hammel (1986)
		Longino (2006)
		McNett et al. (2010)
	<i>Ocotea austinii</i> CK Allen	Longino (2006)
	<i>Pleurothyrium</i> sp.	Davidson et al. (1989)
Malvaceae	<i>Hampea appendiculata</i> (Donn. Smith) Standl	Longino (2006)
Melastomataceae	<i>Clidemia heterophylla</i> (Desr.) Gleason	Renner & Ricklefs (1998)
	<i>Conostegia</i> sp.	Longino (2006)
	<i>Miconia</i> sp.	Hanson & Longino (2006)
	<i>Pterocladon spruce</i> Hook. F.	Wheeler (1934)
	<i>Tibouchina mutabilis</i> (Vell.) Cogn.	Morini et al. (2006)
	<i>Tococa guianensis</i> Aubl.	Morawetz et al. (1992)
		Renner & Ricklefs (1998)
Meliaceae	<i>Carapa guianensis</i> Aubl.	Longino (2006)
	<i>Guarea</i> sp.	Hanson & Longino (2006)
	<i>Guarea rhopalocarpa</i> Radlk.	Longino (2006)
Moraceae	<i>Ficus</i> sp.	Longino (2006)
Myristicaceae	<i>Virola</i> sp.	Edwards et al. (2009)
Rubiaceae	<i>Borreria verticillata</i> (L.) G. Mey	Wheeler (1934)
	<i>Coffea arabica</i> L.	Armbrecht & Gallego (2007)
		Philpott et al. (2006)
	<i>Duroia hirsute</i> (Poeppig & Endl) Schumann.	Davidson & McKey (1993)
		Frederickson et al. (2005)
Tiliaceae		Frederickson & Gordon (2009)
	<i>Palicourea</i> sp.	Longino (2006)
	<i>Hasseltia floribunda</i> Kunth.	Haber et al. (1981)
Urticaceae	<i>Pourouma</i> sp.	Edwards et al. (2009)



the litter (Table 4), all of which were with a single queen. Winged ants were recorded only in four colonies, three of *M. ruzskii* and one of *M. catharinae*. The populations of workers and immature individuals in the *M. ruzskii* and *M. catharinae* colonies were not significantly different (Table 3). The canopy openness was associated with the number of nests in the leaf litter ( $r_s = -0.4151$ ;  $P = 0.02$ ) (Fig. 4)

#### DISCUSSION

The seven *Myrmelachista* species recorded in the studied Atlantic Forest remnants complement the existing data on ant biodiversity in the region (Suguituru et al. 2011, 2013) and the tree species in which nests of this ant genus can be found (Table 4).

The species richness that was found is relevant for biodiversity studies because workers are rarely observed foraging on the surfaces of plants (Longino 2006); as a

consequence, species records are rare or even absent in studies on arboreal ants (Silvestre et al. 2003, Morini et al. 2006, Delabie et al. 2007).

Most of the nests found in the vegetation were located in the tips of dry twigs of live trees or in dead trunks of trees; few nests were found in live stems. Nevertheless, most *Myrmelachista* species nest in cavities or live stems of trees (Stout 1979, Brown 2000, Longino 2006, Edwards et al. 2009).

*Myrmelachista* nests are also found in dry twigs fallen in the litter but are more abundant in areas with open canopies. For other ant species recorded in dry twigs in the Atlantic Forest, the abundance of nests increases with canopy cover (Fernandes et al. 2012), as there is a greater supply of resources (Philpott & Foster 2005). Most nests in dry twigs recorded by Nakano et al. (2012) in sunny trials belonged to *M. catharinae* and *M. ruzskii*, which corroborates the present work. In addition, five species that occur at the tips of dry twigs of live




Fig. 4: Association between canopy cover and the number of nests in twigs in the Atlantic Forest of southeastern Brazil.

Asociación entre cobertura del dosel y el número de nidos en ramas pequeñas en la Floresta Atlántica del sudeste de Brasil

trees are found in dry twigs fallen on the litter (Nakano et al. 2012). Exception is made for *M. arthuri* that was recorded in this work only in live stems, but not in fallen twigs on the litter (Nakano et al. 2012).

Thus, it is likely that the some *Myrmelachista* species found in this study have a preference for nesting in drier microhabitats and that the twigs inhabited by ants in the litter originated from the trees listed in Table 1, which housed the colony with the queen and winged individuals at the time at which the twigs fell to the ground. This fact is supported (1) by the observation made in this study for *M. gallicola* and also by Ketterl et al. (2003), who found the presence of three *Myrmelachista* species in *Araucaria angustifolia* (Bertol.) Kuntze, both in trees and nearby fallen twigs, and (2) because the most frequently observed *Myrmelachista* species in tips of dry twigs of live trees are the same as those found nesting in twigs on the litter surface. Conversely, the fact that only *M. catharinae* and *M. ruskii* nests were found in dead twigs fallen in the litter is interesting because other species (*Myrmelachista* sp. 4; *Myrmelachista* sp. 7; *M. nodigera* and *M. gallicola*) were recorded in the tips of dry twigs. Moreover, Nakano et al. (2012) reported *Myrmelachista* sp. 4, *M. catharinae*, *M. ruskii*, *M. nodigera*, *M. gallicola*, and *M. reticulata* in fallen twigs in the leaf litter of the sunny Atlantic Forest.

Specifically for *M. gallicola*, the 14 nests found in a single plant suggest a possible case of polydomy because all of the nests had workers, immature individuals and one fertilized queen (Debout et al. 2007, Lanan et al. 2011). However, no connection was found among them, corroborating the results of Carvalho & Vasconcelos (2002), Santos & Del Claro (2009) and Lanan et al. (2011). Polydomy is relatively common among ants, occurring in over 150 species (Debout et al. 2007).

The occurrence of polydomy may be an alternative for *Myrmelachista* colonies possessing growth that may be limited by the space available for nesting, but polydomy may also be related to polygyny (Walín et al. 2001), increased foraging area (Santos & Del Claro 2009, Schmolke 2009, Lanan et al. 2011), offspring thermoregulation (Pedersen & Boomsma 1999) and type of food source, such as extrafloral nectaries and insect exudates

in the case of *Myrmelachista* (Pfeiffer & Linsenmair 1998). *Myrmelachista* also uses other food sources external to the host plant (McNett et al. 2010). *M. ramulorum* Wheeler workers, for example, were observed carrying plant and animal parts (eggs, feces and larvae), and even fungi, into their nest (Torres 1984). The workers of *M. arthuri* kept in the laboratory collected insect pieces (cockroaches and termites) and carried them into the nest (Nakano MA personal obs.).

When disturbed, due to a more abrupt motion of the researcher, while foraging or at the nest entrance, individuals of *M. ruskii*, *M. catharinae*, *M. arthuri* and *Myrmelachista* sp. 7 responded aggressively, immediately elevating their gaster. Conversely, individuals of *M. nodigera*, *M. gallicola* and *Myrmelachista* sp. 4 quickly hid in nest holes. The aggressive behavior of gaster elevation has been previously observed for *M. zeledoni* Emery (Longino 2006) and *Myrmelachista* sp. (Espírito Santo 2008). Davidson et al. (1989) described a less aggressive behavior for another unidentified species.

The natural history of most ant species is still poorly understood, preventing broad generalizations (Del-Claro et al. 2009). This study partly fills the gap in the body of knowledge regarding the biology of the genus *Myrmelachista* by reporting observations concerning a number of its species, such as the ability of colonies to emergence winged reproductives especially during rainy season, that is an important stimulus for the synchronized release of winged ants (Torres et al. 2001; Santos & Del Claro 2009). This result complements the work of Nakano et al. (2012) that reported the emergence of males and females of *Myrmelachista* during rainy season and the confirmation of *M. ruskii* and *M. catharinae* as common species in Atlantic Forest remnants of southeastern Brazil. Furthermore, this study reports the preference of these species for sites with opener canopies.

**ACKNOWLEDGEMENTS:** The authors would like to thank the National Council for Scientific and Technological Development (Conselho Nacional de Desenvolvimento Científico e Tecnológico, CNPq) for the scholarships that were granted to MA Nakano (grant 132485/2008-7) and MSC Morini (grant 301151/2009-1), the São Paulo Research Foundation (FAPESP) for aid to RM Feitosa (grant 07/01310-2) and MSC Morini (grant 10/50294-

2) and Prof. Dr. CRF Brandão for making valuable suggestions.

# LITERATURE CITED

- ARMBRECHT I & MC GALLEGO (2007) Testing ant predation on the coffee berry borer in shaded and sun coffee plantations in Colombia. *Entomologia Experimentalis et Applicata* 124: 261-267.
- BENZING DH (1970) An investigation of two bromeliad myrmecophytes: *Tillandsia butzii* Mez, *T. caput-medusae* E. Morren, and their ants. *Bulletin of the Torrey Botanical Club (USA)* 97: 109-115.
- BOLTON B, G ALPERT, PS WARD & P NASKRECKI (2006) *Bolton's Catalogue of Ants of the World 1758-2005*. Harvard University Press, Cambridge. CD-ROM.
- BROWN WL (2000) Diversity of ants. In: Agosti D, JD Majer, Alonso LE & TR Schultz (eds) *Ants: standard methods for measuring and monitoring biodiversity*: 45-79. Washington: Smithsonian Institution, Washington D.C., USA.
- CAMPOS RI, HL VASCONCELOS, SP RIBEIRO, FS NEVES & JP SOARES (2006) Relationship between tree size and insect assemblages associated with *Anadenanthera macrocarpa*. *Ecography* 29: 442-450.
- CARVALHO KS & HL VASCONCELOS (2002) Comunidade de formigas que nidificam em pequenos galhos da serapilheira em floresta da Amazônia Central, Brasil. *Revista Brasileira de Entomologia* 46: 115-121.
- DAVIDSON DW & D MCKEY (1993) Ant-plant symbioses: stalking the chuyachaqui. *Trends in Ecology and Evolution* 8: 326-332.
- DAVIDSON DW, RR SNELLING & JT LONGINO (1989) Competition among ants for myrmecophytes and the significance of plant trichomes. *Biotropica* 21: 64-73.
- DEBOUT G, B SCHATZ, M ELIAS & D MCKEY (2007) Polydomy in ants: what we know, what we think we know, and what remains to be done. *Biological Journal of the Linnean Society* 90: 319-348.
- DELABIE JHC, B JAHYNY, IC NASCIMENTO, CSF MARIANO, S LACAU, S CAMPIOLO, S.M PHILPOTT & M LEPONCE (2007) Contribution of cocoa plantations to the conservation of native ants (Insect: Hymenoptera: Formicidae) with a special emphasis on the Atlantic forest fauna of southern Bahia, Brazil. *Biodiversity and Conservation (Netherlands)* 16: 2359-2384.
- DEL-CLARO K, HM TOREZAN-SILINGARDI, C BELCHIOR & E ALVES-SILVA (2009) *Ecologia Comportamental: uma ferramenta para a compreensão das relações animal-planta*. *Oecologia Brasiliensis (Brazil)* 13: 16-29.
- EDWARDS DP, ME FREDERICKSON, GH SHEPARD & DW YU (2009) A plant needs ants like a dog needs fleas: *Myrmelachista schumanni* ants gall many trees species to create housing. *American Naturalist* 174: 734-740.
- ESPÍRITO-SANTO NB (2008) *Assembléia de formigas do Parque Estadual do Itacolomi (MG) e relações intra e interespecíficas entre espécies dominantes*. Master's Dissertation, Institute of Biological Sciences, Federal University of Juiz de Fora.
- FERNÁNDEZ F & S SENDOYA (2004) List of Neotropical ants (Hymenoptera: Formicidae). *Biota Colombiana* 5: 3-93.
- FERNANDES TT, RR SILVA, DR SOUZA, N ARAÚJO & MSC MORINI (2012) Undecomposed twigs in the leaf litter as nest-building resources for ants (Hymenoptera: Formicidae) in areas of the Atlantic Forest in the southeastern region of Brazil. *Psyche (Egypt)* 2012: 1-8.
- FRAZER GW, CD CANHAM & KP LERTZMAN (1999) *Gap Light Analyzer (GLA): Imaging software to extract canopy structure and gap light transmission indices from true-colour fisheye photographs, users manual and program documentation*. Copyright © 1999: Simon Fraser University, Burnaby, British Columbia, and the Institute of Ecosystem Studies, Millbrook, New York.
- FREDERICKSON ME (2005) Ant species confer different partner benefits on two neotropical myrmecophytes. *Oecologia* 143: 387-395.
- FREDERICKSON ME (2006) The reproductive phenology of an Amazonian ant species reflects the seasonal availability of its nest sites. *Oecologia* 149: 418-427.
- FREDERICKSON ME, DM GORDON & M J GREENE (2005) "Devil's gardens" bedeviled by ants. *Nature* 437: 495-496.
- FREDERICKSON ME & DM GORDON (2009) The intertwined population biology of two Amazonian myrmecophytes and their symbiotic ants. *Ecology* 90: 1595-1607.
- GOITÍA W & K JAFFÉ (2009) Ant-plant associations in different forests in Venezuela. *Neotropical Entomology* 38: 7-31.
- HABER WA, GW FRANKIE, HG BAKER, I BAKER & S KOPTUR (1981) Ants like flower nectar. *Biotropica* 13: 211-214.
- HAMMEL BE (1986) The vascular flora of La Selva Biological Station, Costa Rica - Lauraceae. *Selbyana* 9: 218-233.
- HANSON PE & JT LONGINO (2006) *Hormigas (Formicidae)*. In: Hanson PE & ID Gauld (eds) *Hymenoptera de la región neotropical*: 617-644. Gainesville: *Memoirs of the American Entomological Institute, Florida, USA*.
- KEMPF WW (1972) *Catálogo Abreviado das Formigas da Região Neotropical*. *Studia Entomologica (Brazil)* 15: 3-344.
- KETTERL J, M VERHAAGH, JH BIHN, CRF BRANDÃO & W ENGELS (2003) Spectrum of ants associated with *Araucaria angustifolia* trees and their relations to hemipteran trophobionts. *Studies on Neotropical Fauna Environment (Germany)* 38: 199-206.
- KUSNEZOV N (1951) *Myrmelachista* en la Patagonia (Hymenoptera, Formicidae). *Acta Zoologica Liloana (Argentina)* 11: 353-365.
- LANAN MC, A DORHAUS & JL BRONSTEIN (2011) The function of polydomy: the ant *Crematogaster torosa* preferentially forms new nests near food sources and fortifies outstations. *Behavioral Ecology and Sociobiology (Germany)* 65: 959-968.
- LONGINO JT 2006 A taxonomic review of the genus *Myrmelachista* (Hymenoptera: Formicidae) in Costa Rica. *Zootaxa* 1141: 1-54.
- McNETT K, JT LONGINO, P BARRIGA, O VARGAS, K PHILLIPS, CL SAGERS (2010) Stable isotope investigation of a cryptic ant-plant association: *Myrmelachista flavocotea* (Hymenoptera: Formicidae) and *Ocotea* spp. (Lauraceae). *Insect Socialia (Belgium)* 57: 67-72.

- MINUZZI RB, GC SEDIYAMA, EM BARBOSA & JCF MELO JR (2007) Climatologia do comportamento do período chuvoso da região sudeste do Brasil. *Revista Brasileira de Meteorologia* (Brazil) 22: 338-344.
- MORAWETZ W, M HENZL & B WALLNOFER (1992) Tree killing by herbicide producing ants for the establishment of pure *Tococa occidentalis* populations in the Peruvian Amazon. *Biodiversity and Conservation* (Netherlands) 1: 19 - 33.
- MORINI MSC, N KAMAZUKA, R LEUNG, SS SUGUITURU & LF SILVA (2006) Ant fauna (Hymenoptera: Formicidae) in Magnoliophyta native to the Atlantic Forest. *Sociobiology* 47: 433-444.
- NAKANO MA, RM FEITOSA, CO MORAES, LDC ADRIANO, EP HENGLES, LE LONGUI & MSC MORINI (2012) Assembly of *Myrmelachista* Roger (Formicidae: Formicinae) in twigs fallen on the leaf litter of Brazilian Atlantic Forest. *Journal of Natural History* 46: 2103-2115
- PEDERSEN JS & JJ BOOMSMA (1999) Genetic analysis of colony structure in polydomous and polygynous ant populations. *Biological Journal of the Linnean Society* 66: 115-144.
- PFEIFFER M & KE LINSSENMAIR (1998) Polydomy and the organization of foraging in a colony of the Malaysian giant ant *Camponotus gigas* (Hymenoptera, Formicidae). *Oecologia* 117: 579-590.
- PHILPOTT SM, S UNO & J MALDONADO (2006) The importance of ants and high-shade management to coffee pollination and fruit weight in Chiapas, Mexico. *Biodiversity and Conservation* (Netherlands) 15: 487-501.
- PHILPOTT SM & PF FOSTER (2005) Nest-site limitation in coffee agroecosystems: artificial nests maintain diversity of arboreal ants. *Ecological Applications* 15: 1478-1485.
- QUIRÁN EM & JJ MARTÍNEZ (2006) Redescrición de la obrera de *Myrmelachista gallicola* (Hymenoptera: Formicidae) y primera cita para la provincial de La Pampa (Argentina). *Revista de la Sociedad Entomológica Argentina* (Argentina) 65: 89-92.
- RENNER SS & RE RICKLEFS (1998) Herbicidal activity of domatia-inhabiting ants in patches of *Tococa guianensis* and *Clidemia heterophylla*. *Biotropica* 30: 324-327.
- ROSUMÉK FB, MA ULYSSÉA, BC LOPES, J STEINER & A ZILLIKENS (2008) Formigas de solo e de bromélias em uma área de Mata Atlântica, Ilha de Santa Catarina, sul do Brasil: levantamento de espécies e novos registros. *Biotemas* 21: 81-89.
- SANTOS JC & K DEL-CLARO (2009) Ecology and behaviour of the weaver ant *Camponotus (Myrmobrachys) senex*. *Journal of Natural History* 43: 1423-1435.
- SCHMOLKE A (2009) Benefits of dispersed central-place foraging: an individual-based model of a polydomous ant colony. *American Naturalist* 173: 772-778.
- SILVESTRE R, CRF BRANDÃO & RR SILVA (2003) Grupos funcionales de hormigas: el caso de los grêmios del Cerrado. In: Fernández F (ed) *Introducción a las hormigas de la región Neotropical*: 113-148. Instituto de Investigación de Recursos Biológicos Alexander Von Humboldt, Colombia.
- STOUT J (1979) An association of an ant, a mealy bug, and an understory tree from a Costa Rican rain forest. *Biotropica* 11: 309-311.
- SUGUITURU SS, RR SILVA, DR SOUZA, CB MUNHAE, MSC MORINI (2011) Ant community richness and composition across a gradiente from *Eucalyptus* plantations to secondary Atlantic Forest. *Biota Neotropica* (Brazil) 11:1-8.
- SUGUITURU SS, DR SOUZA, CB MUNHAE, R PACHECO, MSC MORINI (2013) Diversidade e riqueza de formigas (Hymenoptera: Formicidae) em remanescentes de Mata Atlântica na Bacia Hidrográfica do Alto Tietê, SP. *Biota Neotropica* (Brazil) 13:1-8.
- TORRES JA (1994) Niches and coexistence of ant communities in Puerto Rico: Repeated patterns. *Biotropica* 26: 124-140.
- TORRES JA, RR SNELLING & M CANALS (2001) Seasonal and nocturnal periodicities in ant nuptial flights in the tropics (Hymenoptera: Formicidae). *Sociobiology* 37: 601-626.
- WALIN L, P SEPPA & L SUNDSTROM (2001) Reproductive allocation within a polygyne, polydomous colony of the ant *Myrmica rubra*. *Ecological Entomology* 26: 537-546.
- WHEELER WM (1934) Neotropical ants collected by Dr. Elisabeth Skwarra and others. *Bulletin of the Museum of Comparative Zoology* (USA) 77: 157-240.

*Editorial responsibility: Marco A. Méndez*

*Received received June 12, 2012; accepted May 17, 2013*

