



Brazilian Journal of Otorhinolaryngology

ISSN: 1808-8694

revista@aborlccf.org.br

Associação Brasileira de
Otorrinolaringologia e Cirurgia Cérvico-
Facial
Brasil

Du, Yuanfeng; Yu, Wenhua

Comparison of nerve combing and percutaneous radiofrequency thermocoagulation in the
treatment for idiopathic trigeminal neuralgia

Brazilian Journal of Otorhinolaryngology, vol. 83, núm. 2, marzo-abril, 2017, p. 240

Associação Brasileira de Otorrinolaringologia e Cirurgia Cérvico-Facial
São Paulo, Brasil

Available in: <http://www.redalyc.org/articulo.oa?id=392450410021>

- How to cite
- Complete issue
- More information about this article
- Journal's homepage in redalyc.org

redalyc.org

Scientific Information System

Network of Scientific Journals from Latin America, the Caribbean, Spain and Portugal

Non-profit academic project, developed under the open access initiative



Brazilian Journal of OTORHINOLARYNGOLOGY

www.bjorl.org



LETTER TO THE EDITOR

Comparison of nerve combing and percutaneous radiofrequency thermocoagulation in the treatment for idiopathic trigeminal neuralgia[☆]



Comparação entre neurólise interna e termocoagulação percutânea por radiofrequência no tratamento da neuralgia idiopática do trigêmeo

Dear Editor,

We read with great interest the article by Zhou et al.¹

Zhou and colleagues compared the efficacy between nerve combing (NC) and percutaneous radiofrequency thermocoagulation (RF) in the treatment for idiopathic trigeminal neuralgia. In the group NC, the rate of satisfactory relief was 82%, satisfactory relief was 76.4% in the group RF; there were no significant differences between the two groups. As known, microvascular decompression (MVD) is the best surgical treatment for trigeminal neuralgia in patients of any age,² only when the neurovascular compression (NVC) is absent, nerve combing or internal neurolysis may be considered.³

In this study, the authors performed nerve combing procedure in 55 cases, but they did not mention whether NVC was present or how many patients present NVC. If typical vascular compression was found at surgery, MVD was the only

procedure; NV might lead to facial numbness after operation. MVD combined with NV may be used in the patients with atypical vascular compression. So the procedure and conclusion might be inaccurate without excluding NVC.

Conflicts of interest

The authors declare no conflicts of interest.

References

1. Zhou X, Liu Y, Yue Z, Luan D, Zhang H, Han J. Comparison of nerve combing and percutaneous radiofrequency thermocoagulation in the treatment for idiopathic trigeminal neuralgia. *Braz J Otorhinolaryngo.* 2016;82:574–9.
2. Sekula RF Jr, Frederickson AM, Jannetta PJ, Quigley MR, Aziz KM, Arnone GD. Microvascular decompression for elderly patients with trigeminal neuralgia: a prospective study and systematic review with meta-analysis. *J Neurosurg.* 2011;114:172–9.
3. Ko AL, Ozpinar A, Lee A, Raslan AM, McCartney S, Burchiel KJ. Long-term efficacy and safety of internal neurolysis for trigeminal neuralgia without neurovascular compression. *J Neurosurg.* 2015;122:1048–57.

Yuanfeng Du, Wenhua Yu*

Nanjing Medical University Affiliated Hangzhou Hospital, The Hangzhou First People's Hospital, Department of Neurosurgery, Hangzhou, China

*Corresponding author.

E-mail: d542762345@126.com (W. Yu).

[☆] Please cite this article as: Du Y, Yu W. Comparison of nerve combing and percutaneous radiofrequency thermocoagulation in the treatment for idiopathic trigeminal neuralgia. *Braz J Otorhinolaryngol.* 2017;83:240.