



Ciência e Tecnologia de Alimentos

ISSN: 0101-2061

revista@sbcta.org.br

Sociedade Brasileira de Ciência e  
Tecnologia de Alimentos  
Brasil

Cruz-Ramírez, Susana-Gabriela; López-Saiz, Carmen-María; Rosas-Burgos, Ema-Carina;  
Cinco-Moroyoqui, Francisco-Javier; Velázquez, Carlos; Hernández, Javier; Burgos-  
Hernández, Armando

Antimutagenic, antiproliferative, and antioxidant effect of extracts obtained from octopus  
(*Paraoctopus limaculatus*)

Ciência e Tecnologia de Alimentos, vol. 35, núm. 4, outubro-diciembre, 2015, pp. 722-728

Sociedade Brasileira de Ciência e Tecnologia de Alimentos  
Campinas, Brasil

Available in: <http://www.redalyc.org/articulo.oa?id=395943288021>

- How to cite
- Complete issue
- More information about this article
- Journal's homepage in redalyc.org

redalyc.org

Scientific Information System

Network of Scientific Journals from Latin America, the Caribbean, Spain and Portugal

Non-profit academic project, developed under the open access initiative

## Antimutagenic, antiproliferative, and antioxidant effect of extracts obtained from octopus (*Paraoctopus limaculatus*)

Susana-Gabriela CRUZ-RAMÍREZ<sup>1</sup>, Carmen-María LÓPEZ-SAIZ<sup>2</sup>, Ema-Carina ROSAS-BURGOS<sup>1</sup>, Francisco-Javier CINCO-MOROYOQUI<sup>1</sup>, Carlos VELÁZQUEZ<sup>3</sup>, Javier HERNÁNDEZ<sup>4</sup>, Armando BURGOS-HERNÁNDEZ<sup>1\*</sup>

### Abstract

The search for chemopreventive/chemoprotective compounds in marine organism has been extensively reported; however, the presence of these compounds in octopus has been incipiently explored. In this research, the antimutagenic, antiproliferative, and antioxidant potential of three crude extracts (methanolic, acetonetic, and hexanic) from *Paraoctopus limaculatus* was investigated. Antimutagenic activity against aflatoxin B<sub>1</sub> (AFB<sub>1</sub>) was evaluated through the Ames test using *Salmonella typhimurium* tester strains TA98 and 100. Antiproliferative activity was assessed using the standard MTT (3-(4,5-dimethyl-2-thiazolyl)-2,5-diphenyl-2-H-tetrazolium bromide) assay on M12.C3.F6 murine cell line. Antioxidant activity was assessed using the DPPH (2,2-diphenyl-1-picrylhydrazyl) and ABTS (2,2'-Azino-bis(3-ethylbenzothiazoline-6-sulfonic acid) methods. Hexanic extract showed the highest antimutagenic and antiproliferative activities inhibiting 80 and 43% of mutagenicity induced by AFB<sub>1</sub> for TA98 and TA100, respectively, and showing a high antiproliferative activity at 200 and 100 µg/mL. However, when the antioxidant activity was evaluated at a concentration of 50 mg/mL, the methanolic fraction exerted inhibition of 98 and 96 % ABTS and DPPH radicals, respectively. RP-HPLC and <sup>1</sup>H-RMN analyses suggested the presence of double bonds with extended conjugation and oxygenated compounds such as alcohols, esters, ethers or ketones. These results suggested that hexanic and methanolic extract from octopus contained compounds with chemoprotective and antioxidant properties.

**Keywords:** octopus; biological activity; chemopreventive.

**Practical Application:** Octopus fractions can be used as potential chemopreventive and antioxidant compounds.

## 1 Introduction

Cancer is the leading cause of mortality worldwide and the number of cases is increasing every year (World Health Organization, 2010). Chemoprevention is a long-term strategy with the best cost-effectiveness ratio for cancer control, which consists in the use of certain chemicals, natural or synthetic, with the aim of preventing carcinogenesis or reversing the development of invasive malignancy (Martínez, 2006). In recent years, the marine environment has been recognized as a resource with high biodiversity but still unexplored. Nowadays is considered a potential source of novel bioactive compounds as new anticancer agents (Pramanik & Pandey, 2013).

A marine source of bioactive compounds are cephalopods which include nautilus, cuttlefishes, squids, and octopus that constitute an important source of many biological substances with antioxidant, antitumor, antileukemic, antibacterial, and antiviral activities (Rajaganapathi et al., 2000, Jayaraj et al., 2008, Jesy et al., 2014, Sudhakar & Nazeer, 2015). These researches suggest that these marine invertebrate may become an important source of new compounds with potential application in biomedicine.

Octopus is one of the species that has been reported as source of bioactive compounds with accentuated antiproliferative and antimutagenic activities, which are able to reverse the mutation of aflatoxin B<sub>1</sub> (AFB<sub>1</sub>) and also selectively act against the proliferation of tumor cells (Moreno-Félix et al., 2013). These antimutagenic compounds extracted with chloroform from octopus, were identified as butyl isobutyl phthalate derivatives (Cruz-Ramírez et al., 2015). Based on the above, the aim of this study was to assess the antimutagenic, antiproliferative, and antioxidant activities exerted by compounds in fractions from octopus.

## 2 Materials and methods

### 2.1 Bacterial cultures and cell lines

*Salmonella typhimurium* TA98 and TA100 (Molecular Toxicology Inc.) were routinely checked to confirm genetic features using the procedure described by Maron & Ames (1983).

Received 27 Aug., 2015

Accepted 25 Nov., 2015

<sup>1</sup>Departamento de Investigación y Posgrado en Alimentos, Universidad de Sonora, Hermosillo, Sonora, México

<sup>2</sup>Programa de Ingeniería Ambiental, Universidad Estatal de Sonora, Hermosillo, Sonora, México

<sup>3</sup>Departamento de Ciencias Químico-Biológicas, Universidad de Sonora, Hermosillo, Sonora, México

<sup>4</sup>Unidad de Servicios de Apoyo en Resolución Analítica, Universidad Veracruzana, México

\*Corresponding author: [aburgos@guayacan.uson.mx](mailto:aburgos@guayacan.uson.mx)

Cell line M12.C3.F6 (murine B-cell lymphoma) was kindly provided by Dr. Emil R. Unanue (Department of Pathology and Immunology, Washington University at St. Louis, MO, USA). M12.C3.F6 was cultured in Dulbecco's modified Eagle's medium (DMEM) (Gibco, Grand Island, NY, USA), supplemented with 5% fetal calf serum and grown at 37 °C at 5% CO<sub>2</sub>.

## 2.2 Testing species

Octopus (*Paraoctopus limaculatus*) was obtained from Bahía de Kino, Sonora, México (29° 22' 27" N, 112° 34' 08" O) and transported in ice to the University of Sonora, Seafood Laboratory. Organism were fresh packed and stored at -25 °C until further analysis. Tentacles were minced, separated in homogeneous portions and freeze-dried at -30 °C under vacuum (Unitop 600L, USA).

## 2.3 Extraction of bioactive compounds

A 40 g portion of lyophilized octopus tentacles was homogenized with 500 mL hexane and the mixture agitated for 60 min using a Burrell Wrist Action Shaker (Burrell Corporation Pittsburgh, PA). Then, the mixture was filtered through Whatman No. 42 filter paper under vacuum and the solvent was evaporated using a rotary evaporator (Yamato RE300 Scientific Co., Ltd, Japan) to dryness under a nitrogen stream and darkness (López-Saiz et al., 2014) in order to obtain the hexane soluble fraction (HSF). The residue obtained after filtration was subjected to the same procedure described above using acetone and later with methanol, in order to obtain the acetone- (ASF) and methanol-soluble fractions (MSF), respectively. Yields (dry basis) of the three fractions were quantified. Octopus fractions were used to assess biological activities.

## 2.4 Antimutagenicity Assay

The antimutagenic potential of the three extracts was evaluated using the mutagenicity assay described by Maron & Ames (1983) with *Salmonella typhimurium* TA98 and TA100 as tester strains in the presence of metabolic activation (S9 mix). Fractions were reconstituted and serially diluted with dimethyl sulfoxide (DMSO) and then spiked with enough pure AFB<sub>1</sub> to achieve a common final concentration of 500 ng of AFB<sub>1</sub>/100 µL. Different AFB<sub>1</sub> concentrations were used as positive controls for both strains. All assays were performed in triplicate.

## 2.5 Antiproliferation assay

The antiproliferative effect of octopus fractions was determined using the standard MTT (3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide) assay (Mosmann, 1983) using M12.C3.F6 (murine B-cell lymphoma) cell line. Fifty-µL aliquots of murine B-cell lymphoma culture were placed in wells of a flat 96-well plate (10,000 cells/ well) and then incubated for 12 h at 37 °C in an atmosphere of 5% CO<sub>2</sub> to allow cell attachment. Cell cultures were incubated with 50 µL of medium containing various concentrations of extract (200, 100, and 50 µg/mL) and incubated for 48 h. Extracts were first re-suspended in DMSO and diluted in supplemented Dulbecco's Modified Eagle's Medium

(DMEM) media. Control cell cultures were incubated with DMSO (final DMSO concentration 0.2% v/v). Prior to the last 4 h of the cell culture, 10 µL of a MTT stock solution (5 mg/mL) were added to each well and after this time, the formazan crystals formed were dissolved with acidic isopropanol (100 µL of 0.04 N HCl in isopropanol). Plates were read in an ELISA plate reader (Benchmark Microplate Reader; Bio-Rad, Hercules, CA, USA) using a test wavelength of 570 nm and a reference wavelength of 630 nm. Plates were read within 15 min after adding isopropanol. Results were expressed as percentages of cell proliferation.

## 2.6 Antioxidant assay

### Scavenging activity on 2,2-diphenyl-1-picrylhydrazyl (DPPH) radical assay

DPPH radical scavenging activity was determined according to the method described by Jie Li et al. (2012) with slight modifications. In this method, 50 µL-aliquots of octopus fractions at different concentrations (50, 25, 10, 5, and 2.5 mg/mL) were placed into wells of 96-well plates and then combined with 200 µL of a DPPH hexane-ethanol (1:1) solution (4 mg/100 mL of solution). Each well was mixed and incubated in the dark for 30 min at room temperature. The absorbance was measured at 540 nm in microplate reader (BIO-RAD xMark™, México). Antioxidant capacity was calculated using the following Equation 1:

$$DPPH \text{ scavenging activity (\%)} = \frac{(1 - \lambda_{540-s})}{\lambda_{540-c}} \times 100 \quad (1)$$

Where  $\lambda_{540-s}$  is the absorbance of the remaining DPPH in the presence of the scavenger and  $\lambda_{540-c}$  is the absorbance of the control with no radical scavenger.

### Scavenging activity on 2,2'-Azino-bis(3-ethylbenzothiazoline-6-sulfonic acid (ABTS) radical assay

The ABTS radical assay was determined according to Jie Li et al. (2012), modified as described below. The ABTS radical solution was prepared by dissolving 8 mg of the radical in 1 mL of H<sub>2</sub>O, reacted with an equal volume (1:1) of a potassium persulfate (K<sub>2</sub>S<sub>2</sub>O<sub>8</sub>) (1.3 mg/mL) solution, and then stored for 16 h in the dark at room temperature. Fifty-µL-aliquots of octopus fractions at different concentrations (50, 25, 10, 5, and 2.5 mg/mL), were placed into wells of 96-well plates and then combined with 200 µL of a previously activated ABTS radical solution. Each well was mixed and incubated in the dark for 30 min at room temperature. The absorbance was measure at 713 nm in microplate reader (BIO-RAD xMark™, México). The antioxidant capacity was calculated using the following Equation 2:

$$ABTS \text{ scavenging activity (\%)} = \frac{(1 - \lambda_{713-s})}{\lambda_{713-c}} \times 100 \quad (2)$$

Where  $\lambda_{713-s}$  is the absorbance of the remaining ABTS in the presence of scavenger and  $\lambda_{713-c}$  is the absorbance of the control with no radical scavenger.

## 2.7 Qualitative analysis by RP-HPLC

The objective of this analysis was to determine the compositional complexity octopus fractions and the retention times of the main compounds in them. The experiment was carried out in an Agilent Technologies 1260 Infinity High Performance Liquid Chromatoph equipped with a photodiode detector, a Zorbax Eclipse XDB-C18 semipreparative column (5 µm particle size, 250 x 9.4 mm i.d.), and a 20 µL-injection loop. Column temperature was set at 20 °C. Elution of components was carried out at a flow rate of 2 ml/min using a linear gradient from 50:50 to 0:100 during the first 9 min mobile phase consisting of a water/methanol mixture followed by acetone/hexane (100:0) during 20 min and concluding with 100% hexane over 35 min. All of the three octopus fractions were monitored at 350 nm and scanned (190 to 600 nm) to detect maximum absorption values.

## 2.8 <sup>1</sup>H-NMR Analysis

Hexanic and menthanolic octopus fractions were dissolved in deuterated chloroform (CDCl<sub>3</sub>) and analyzed by <sup>1</sup>H-NMR. Measurements were carried out in a Bruker Avance 400 MHz equipment using tetramethylsilane (TMS) as reference.

## 2.9 Statistical analysis

The statistical analysis was performed using the software JMP (Visual statistical discovery software, 5.01). Data were analyzed by ANOVA followed by a Tukey-Kramer test ( $P < 0.05$ ).

# 3 Results and discussion

## 3.1 Extraction yield

The extraction of octopus compounds was carried out using three different solvents: methanol, acetone, and hexane, obtaining the highest yield (16.91 % dry basis) with methanol. Athukorala et al. (2003) obtained similar results reporting that methanol was one of the most effective organic solvents for compounds extraction from a marine alga achieving an 18.4 % yield. Recently, Umayaparvathi et al. (2013), evaluated one methanolic extract from octopus and they obtained a 0.85 % yield in wet basis. These last results showed a lower extraction yield compared to those obtained in our experiment, the differences may be related to the extraction process. On the other hand, the HSF and ASF showed the lowest yield with 1.45 and 0.45 %, respectively. These results suggest that methanol is an effective solvent to extract a wide range of compounds from octopus.

## 3.2 Antimutagenic activity

The antimutagenic potential of the three fractions obtained from octopus is shown in Table 1, which corresponds to both *Salmonella typhimurium* tester strains. The antimutagenic activities of MSF and ASF were not statistically ( $p < 0.05$ ) different achieving 50 and 45 % inhibition of 500 ng of pure AFB<sub>1</sub>, respectively, when exposing tester strain TA98 to the highest octopus fraction concentration tested. Similar results were observed for TA100 tester strain, but in this case the octopus fraction used decreased the reversion rate achieved by AFB<sub>1</sub> by 13 and 2 % for MSF and ASF.

However, the inhibition of the mutagenicity of 500 ng AFB<sub>1</sub> exerted by the HSF at 5 mg/mL, was significantly ( $p < 0.05$ ) higher than that observed for the MSF or ASF. These results were also observed in both *Salmonella* tester strains, suggesting the presence of compounds that might protect the genetic material from suffering both types of mutation, base pair or frame-shift.

Stankevics et al. (2008), evaluated the antimutagenic activity from two extract: acetone- and ethanol-soluble fractions, from the sponge *Arenosclera brasiliensis*, finding the acetone-soluble fraction was the most antimutagenic using *Salmonella* TA100 tester strain. More recently, extracts from the marine organism *Sphyrna lewini* were evaluated for antimutagenic potential by Samadi et al. (2011), revealing that methanolic extracts from cartilage exerted the highest antimutagenic effect against sodium azide (as a control mutagen).

Nevertheless, the antimutagenic activity of hexane-soluble fractions has been evaluated in medical plants such as *Astragalus* species (Gulluce et al., 2010). They found five hexane-soluble fractions with antimutagenic activity in *Salmonella typhimurium* TA1537 strain that may be due to their inhibition capabilities by blocking 9-AA binding to DNA *in vitro*. In this context, the results obtained in the present study are promising in finding biologically active compounds that might be candidates as possible cancer preventive agents.

## 3.3 Antiproliferative activity

The cell-growth inhibitory effect of octopus fractions on murine B-cell lymphoma cell line (M12.C3.F6) is shown in Figure 1. The HSF exhibited an antiproliferative activity significantly ( $p < 0.05$ ) higher than either MSF or ASF, tested at concentrations of 200 and 100 µg/mL. On the other hand, whereas the ASF only inhibited approximately 20 % cell proliferation, the MSF did not show inhibition of cellular proliferation compared to the control. Althunibat et al. (2009) evaluated the antiproliferative effects of aqueous and organic extracts from three species of sea cucumbers; the organic extract from the species *Holothuria scabra* inhibited the growth of both cancer cell lines, C33A (human cervix) and A549 (human lung), at cytotoxic concentrations (IC<sub>50</sub>) of

**Table 1.** Revertants<sup>†</sup> induced by AFB<sub>1</sub> spiked obtained of extracts from *Paraoctopus limaculatus* evaluated on *Salmonella typhimurium* TA98 and TA100 tester strain with S9.

Octopus fraction	5 mg/mL	0.5 mg/mL	0.05 mg/mL
<b>TA98</b>			
Hexane-soluble	133 ± 4.1 <sup>b</sup>	373 ± 11.3 <sup>a</sup>	523 ± 10 <sup>ab</sup>
Acetone-soluble	340 ± 3.2 <sup>a</sup>	429 ± 7.7 <sup>a</sup>	476 ± 5.8 <sup>a</sup>
Methanol-soluble	312 ± 1.4 <sup>a</sup>	385 ± 7.4 <sup>a</sup>	358 ± 5.1 <sup>a</sup>
<b>TA100</b>			
Hexane-soluble	602 ± 18.2 <sup>b</sup>	694 ± 6.3 <sup>b</sup>	919 ± 13.7 <sup>a</sup>
Acetone-soluble	921 ± 8.4 <sup>a</sup>	1012 ± 8.3 <sup>a</sup>	1048 ± 5.6 <sup>a</sup>
Methanol-soluble	1030 ± 2.9 <sup>a</sup>	1052 ± 1.5 <sup>a</sup>	1063 ± 4.4 <sup>a</sup>


<sup>†</sup>Positive control AFB<sub>1</sub> (500 ng/plate) induced 602 ± 62 and 1051 ± 69 revertants/plate for TA98 and TA100 tester strains, respectively. Spontaneous revertants were 22 ± 4 and 96 ± 6 for TA98 and TA100 tester strains, respectively. <sup>a</sup>Values are means of three replicates ± standard error mean. Different letters in a column represent significant differences ( $p < 0.05$ ).




15.5 and 3.5 mg/mL. Furthermore, recent research evaluated the antiproliferative effect of lipids fractions from octopus by-products on human breast cancer cell lines MCF-7 and MDA-MB-231. Two lipidic sub-fractions exhibited significant anti-proliferative effect on MC-7; authors found that compounds like sterols or terpenes present in octopus by-products may be responsible for these biological activity (Fitahia et al., 2015).

### 3.4 Antioxidant activity

Figure 2a and 2b show the inhibition percentage for both radicals exerted by the three octopus fractions. MSF showed the highest percentage of radical inhibition at concentration of 50 mg/mL evaluated on ABTS radical-scavenging capacity.



**Figure 1.** Antiproliferative effect of fractions obtained from octopus on transformed cell line M12.C3.F6 (murine B-cell lymphoma). All values are mean from three determinations. Values with the same letters are not significantly ( $p > 0.05$ ) different.



**Figure 2.** Antioxidant activity of acetone-, hexane-, and methanol-soluble fractions obtained from octopus (*Paraoctopus limaculatus*) against ABTS (a) and DPPH (b) radicals. Values are means  $\pm$  SD of triplicate experiments.


These results are consistent with the DPPH radical-scavenging capacity; because MSF exhibited values of 96.84, 83.28, 80.5, 77.45, and 75.98 % at concentrations of 50, 25, 10, 5, and 2.5 mg/mL, respectively. These results are supported by Senthilkumar & Sudha (2012); they found a methanolic extract from seaweed *Chaetomorpha linum* that contains compounds with high scavenging activity attributed to phenolic OH groups, which are major contributors to antioxidant activity and anticancer effects. Moreover, peptide-like compounds have been isolated from marine organisms, which act as ROS radical scavengers, preventing DNA damage, and inhibiting lipidic peroxidation (Sudhakar & Nazeer et al., 2015).

Many studies correlate the antimutagenic effect with antioxidant activity because antioxidant compounds may act in the early stages of carcinogenesis, either interacting with mutagens or by stabilizing free radicals that cause damage to genetic material (Aqil et al., 2008; Zahin et al., 2010a, b, 2013; Bhattacharya, 2011; Sultana et al., 2014). However, antimutagens present in the HSF were not able to act as antioxidant. The differences in scavenging antioxidant and antimutagenic activity may be due to polarity characteristic of compounds extracted with different solvents and to the mechanism of action by which they act.

### 3.5 Qualitative analysis by RP-HPLC

Chromatographical analysis and the spectrophotometric scanning of three fractions obtained from octopus are shown in Figure 3. Results of MSF analysis suggest the presence of polar compounds evidenced by the presence of signals at times  $< 10$  min. The highest absorption signals were observed at 350 nm but most were absorbed in the near UV spectrum.

The ASF emitted signals at retention times similar to those observed (10 min) for the MSF, but also this fraction emitted signals at retention times  $> 10$  min, suggesting the presence of




**Figure 3.** RP-HPLC chromatograms (a-c) and spectrometric scanning (d-f) of methanol-, acetone-, and hexane-soluble fractions, respectively, all obtained from octopus (*Paraoctopus limaculatus*).

compounds with lower polarity than those found in the MSF. Also, most of the absorption was detected in the near UV spectrum, which is also consistent with the presence of non-polar compounds. Spectrophotometric scanning of the HSF showed absorption signals also in the near UV spectrum (after 18 min), also consistent with the presence of non-polar compounds.

### 3.6 $^1\text{H-NMR}$ Analysis

$^1\text{H-NMR}$  analysis of HSF (Figure 4a) shows chemical shifts of the protons between  $\delta_{\text{H}}$  0.7 – 2.4 that correspond to hydrogen directly bound to carbon atoms, which is attributed to methine, methyl, and methylene protons.

The spectrum shows the presence of protons bound to carbons attached to an oxygen atom  $\delta_{\text{H}}$  2.7 – 5.5 which may be attributed to functional groups such as alcohols, acetals, esters, ethers, and ketones; however, in this field (between 4.6 and 5.9) signals corresponding to vinyl protons in alkenes may be found, therefore, compounds with conjugated double bond systems may also be present. These conjugated systems have great absorption intensities in the near UV spectrum, which was previously observed in the RP-HPLC scanning analyses. This spectrum shows no characteristic signals of aromatic protons, suggesting that antimutagenic compounds may have different structures, which is consistent with results reported by Cruz-Ramírez et al.



**Figure 4.**  $^1\text{H}$ -NMR spectrum of (a) antimitagenic and antiproliferative hexanic-soluble fraction, and (b) antioxidant methanol-soluble fraction, both obtained from octopus (*Paraoctopus limaculatus*). TMS was used as a reference.

(2015).  $^1\text{H}$ -NMR spectrum (Figure 4b) of MSF shows irradiation signals of methyl, methylene, and methine ( $\delta_{\text{H}}$  0.6 – 2.6), which suggest the presence of protons associated to aliphatic chains. The signals displayed between  $\delta_{\text{H}}$  3.2 – 4.0 correspond to protons adjacent to carbon attached to oxygen atoms. Previously, we discussed that the antioxidant activity of the MSF may due to the presence of phenolic compounds; however, these compounds have a complex aromatic ring system not observed in the  $^1\text{H}$ -NMR spectrum ( $\delta_{\text{H}}$  6.0 – 8.5), suggesting the presence of different compounds capable of inhibiting ABTS and DPPH radicals.

## 4 Conclusions

Results from this research suggest that HSF from octopus contains antimitagenic and antiproliferative compounds, while potent antioxidant compound are mainly contained in the MSF. Based on RP-HPLC and  $^1\text{H}$ -NMR data, both MSF and HSF contain biologically active non-polar compounds, possibly featuring conjugated double-bond systems; however, further research is required for their full characterization.

## Acknowledgements

Authors wish to thank Consejo Nacional de Ciencia y Tecnología (CONACYT) Mexico for financing the project 241133 and for the graduate scholarship granted to Susana-Gabriela Cruz-Ramírez.

## References

- Althunibat, O. Y., Hashim, R. B., Taher, M., Daud, J. M., Ikeda, J. M., & Zali, B. I. (2009). In vitro antioxidant and antiproliferative activities of three malaysian sea cucumber species. *European Journal of Scientific Research*, 37(3), 376-387.
- Aqil, F., Zahin, M., & Ahmad, I. (2008). Antimutagenic activity of methanolic extracts of four ayurvedic medicinal plants. *Indian Journal of Experimental Biology*, 46(9), 668-672. PMID:18949897.
- Athukorala, Y., Lee, K.-W., Song, C., Ahn, C.-B., Shin, T.-S., Cha, Y.-J., Shahidi, F., & Jeon, Y.-J. (2003). Potential, antioxidant activity of marine red alga *Grateloupia Filicina* extracts. *Journal of Food Lipids*, 10(3), 251-265. <http://dx.doi.org/10.1111/j.1745-4522.2003.tb00019.x>.
- Bhattacharya, S. (2011). Natural antimutagens: a review. *Research Journal of Medicinal Plant*, 5(2), 116-126. <http://dx.doi.org/10.3923/rjmp.2011.116.126>.
- Cruz-Ramírez, S. G., López-Saiz, C. M., Plascencia-Jatomea, M., Machi-Lara, L., Rocha-Alonzo, F., Márquez-Ríos, E., & Burgos-Hernández, A. (2015). Isolation and identification of an antimitagenic phthalate derivative compound from octopus (*Paractopus limaculatus*). *Tropical Journal of Pharmaceutical Research*, 14(17), 1257-1267. <http://dx.doi.org/10.4314/tjpr.v14i7.19>.
- Fitahia, E. M., Raheriniaina, C. E., Bazin, M. A., Huvelin, J.-M., Logé, C., Ranaivoson, E., & Nazih, H. (2015). Anti-proliferative and pro-apoptotic effect of dichloromethane extract of *Octopus vulgaris* by-products on human breast cancer Cell Lines. *Waste Biomass Valor*, 6(2), 237-242. <http://dx.doi.org/10.1007/s12649-014-9344-1>.
- Gulluce, M., Agar, G., Baris, O., Karadayi, M., Orhan, F., & Sahin, F. (2010). Mutagenic and Antimutagenic effects of hexane extract of some *Astragalus* species grown in the eastern anatolia region of Turkey. *Phytotherapy Research*, 24(7), 1014-1018. PMID:19957249.
- Jayaraj, S. S., Thiagarajan, R., Arumugam, M., & Mullainadhan, P. (2008). Isolation, purification and characterization of [beta]-1,3-glucan binding protein from the plasma of marine mussel *Perna viridis*. *Fish & Shellfish Immunology*, 24(6), 715-725. <http://dx.doi.org/10.1016/j.fsi.2007.11.012>. PMID:18420422.
- Jesy, J. H., Thilaga, R. D., & Mary, J. (2014). Anticancer activity of marine cephalopod (*S. Pharaonis*) from Gulf of mannar. *International Journal of Current Research*, 6(11), 10009-10012.
- Jie Li, W., Long Cheng, X., Liu, J., Chao Lin, R., Li Wang, G., Shan Du, S., & Long Liu, Z. (2012). Phenolic compounds and antioxidant activities of *Liriope muscari*. *Molecules (Basel, Switzerland)*, 17(2), 1797-1808. PMID:22328078.
- López-Saiz, C. M., Velázquez, C., Hernández, J., Cinco-Moroyoqui, F. C., Plascencia-Jatomea, M., Robles-Sánchez, M., Machi-Lara, L., & Burgos-Hernandez, A. (2014). Isolation and structural elucidation of antiproliferative compounds of lipidic fractions from white shrimp muscle (*Litopenaeus vannamei*). *International Journal of Molecular Sciences*, 15(12), 23555-23570. <http://dx.doi.org/10.3390/ijms151223555>. PMID:25526568.
- Maron, D. M., & Ames, B. N. (1983). Revised methods for the *Salmonella* mutagenicity test. *Mutation Research*, 113(3-4), 173-215. PMID:6341825.
- Martínez, J. E. (2006). Quimioprevención del cáncer de mamá. *Oncología*, 29(1), 16-21.

- Moreno-Félix, C., Wilson-Sánchez, G., Cruz-Ramírez, S. G., Velázquez-Contreras, C., Plascencia-Jatomea, M., Acosta, A., Machi-Lara, L., Aldana-Madrid, M. L., Ezquerra-Brauer, J. M., Rocha-Alonzo, F., & Burgos-Hernández, A. (2013). Bioactive lipidic extracts from octopus (*Paraoctopus limaculatus*): antimutagenicity and antiproliferative studies. *Evidence-Based Complementary and Alternative Medicine*, 2013, 1-12. <http://dx.doi.org/10.1155/2013/273582>. PMID:23401709.
- Mosmann, T. (1983). Rapid colorimetric assay for cellular growth and survival: application to proliferation and cytotoxicity assays. *Journal of Immunological Methods*, 65(1-2), 55-63. [http://dx.doi.org/10.1016/0022-1759\(83\)90303-4](http://dx.doi.org/10.1016/0022-1759(83)90303-4). PMID:6606682.
- Pramanik, K., & Pandey, A. (2013). Natural compounds: prospective of chemoprevention. *Endocrinology & Metabolic Syndrome*, 2(2), 2-3. <http://dx.doi.org/10.4172/2161-1017.1000e115>.
- Rajaganapathi, J., Thyagarajan, S. P., & Edward, J. K. (2000). Study on cephalopod's ink from anti-retroviral activity. *Indian Journal of Experimental Biology*, 38(5), 519-520. PMID:11272422.
- Samadi, S., Emtiazjoo, M., Khanipour, A. A., Moghadasi, Z., & Nazarhaghghi, F. (2011). Antimutagenic activity of Chloroformic and Methanolic extracts of muscle, liver and cartilage of *Sphyrna lewini* with the Ames test. *Iranian Journal of Fisheries Sciences*, 10(1), 135-142.
- Senthilkumar, P. & Sudha, S. (2012). Antioxidant and antibacterial properties of methanolic extract of green seaweed *Chaetomorpha linum* from Gulf of Mannar: Southeast coast of India. *Jundishapur Journal of Microbiology*, 5(2), 411-415. <http://dx.doi.org/10.5812/jjm.3400>.
- Stankevics, L., Aiub, C. A., Mazzei, J. L., Lobo-Hajdu, G., & Felzenszwalb, I. (2008). Cytotoxic, mutagenic and antimutagenic screening of *Arenosclera brasiliensis* acetone and ethanol extracts. *Genetics and Molecular Research*, 7(2), 542-548. <http://dx.doi.org/10.4238/vol7-2gmr460>. PMID:18752179.
- Sudhakar, S., & Nazeer, R. (2015). Preparation of potent antioxidant peptide from edible part of shortclub cuttlefish against radical mediated lipid and DNA damage. *LWT - Food Science and Technology*, 64, 593-601.
- Sultana, B., Anwar, F., Mushtaq, M., Aslam, M., & Ijaz, S. (2014). In vitro antimutagenic, antioxidant activities and total phenolics of clove (*Syzygium aromaticum* L.) seed extracts. *Pakistan Journal of Pharmaceutical Sciences*, 27(4), 893-899. PMID:25015457.
- Umayaparvathi, S., Arumugam, M., Meenakshi, S., & Balasubramanian, T. (2013). Studies on Antifungal, Cytotoxic and HPTLC profiles of Mollusks and Echinoderm extracts from Southeast Coast of India. *Journal of Medical & Allied Health Sciences*, 2(1), 24-32.
- World Health Organization – WHO. (2010). *Prevención del cáncer*. Retrieved from <http://www.who.int/cancer/prevention/es/>
- Zahin, M., Ahmad, I., & Aqil, F. (2010a). Antioxidant and antimutagenic activity of *Carum copticum* fruit extracts. *Toxicology In Vitro*, 24(4), 1243-1249. <http://dx.doi.org/10.1016/j.tiv.2010.02.004>. PMID:20149861.
- Zahin, M., Aqil, F., & Ahmad, I. (2010b). Broad spectrum antimutagenic activity of antioxidant active fraction of *Punica granatum* L. peel extracts. *Mutation Research/Genetic Toxicology and Environmental Mutagenesis*, 703(2), 99-107. <http://dx.doi.org/10.1016/j.mrgentox.2010.08.001>. PMID:20708098.
- Zahin, M., Aqil, F., Husain, F. M., & Ahmad, I. (2013). Antioxidant capacity and antimutagenic potential of *Murraya koenigii*. *BioMed Research International*, 2013, 1-10. <http://dx.doi.org/10.1155/2013/263509>. PMID:23853769.