



Revista Brasileira de Parasitologia
Veterinária

ISSN: 0103-846X

zacariascbpv@fcav.unesp.br

Colégio Brasileiro de Parasitologia
Veterinária
Brasil

Santiago Alberto Carlos, Renata; Carvalho, Fábio Santos; Arias Wenceslau, Amauri;
Pereira Almosny, Nadia Regina; Rêgo Albuquerque, George

Risk factors and clinical disorders of canine ehrlichiosis in the South of Bahia, Brazil
Revista Brasileira de Parasitologia Veterinária, vol. 20, núm. 3, julio-septiembre, 2011, pp.
210-214

Colégio Brasileiro de Parasitologia Veterinária
Jaboticabal, Brasil

Available in: <http://www.redalyc.org/articulo.oa?id=397841481005>

- How to cite
- Complete issue
- More information about this article
- Journal's homepage in redalyc.org

redalyc.org

Scientific Information System

Network of Scientific Journals from Latin America, the Caribbean, Spain and Portugal

Non-profit academic project, developed under the open access initiative

Risk factors and clinical disorders of canine ehrlichiosis in the South of Bahia, Brazil

Fatores de risco e alterações clínicas da ehrlichiose canina no Sul da Bahia, Brasil

Renata Santiago Alberto Carlos¹; Fábio Santos Carvalho²; Amauri Arias Wenceslau¹;
Nadia Regina Pereira Almosny³; George Rêgo Albuquerque^{1*}

¹Departamento de Ciências Agrárias e Ambientais, Universidade Estadual de Santa Cruz – UESC

²Programa de Pós-Graduação em Ciência Animal, Universidade Estadual de Santa Cruz – UESC

³Departamento de Patologia Clínica Veterinária, Universidade Federal Fluminense – UFF

Received December 13, 2010

Accepted August 17, 2011

Resumo

Objetivou-se com este trabalho estudar as alterações clínicas, fatores de risco da ehrlichiose canina nos municípios de Ilhéus e Itabuna, Bahia, e comparar diferentes métodos de diagnóstico. Amostras de sangue foram coletadas de 200 cães e cada animal foi examinado clinicamente. Foi preenchido um questionário para avaliar os fatores de risco. As amostras de sangue foram analisadas pelo teste Dot-ELISA e foram realizadas hematimetria, contagem de plaquetas e procura de mórulas em esfregaço de sangue. Nested-PCR foi realizada em 50 amostras positivas e 50 negativas na sorologia. Três amostras PCRs positivas foram sequenciadas. Foi encontrado 36,0% de positividade na sorologia e 5,5% nos esfregaços sanguíneos. Os animais apresentavam anemia e trombocitopenia. Ter carrapatos e residir em áreas suburbanas foram considerados fatores de risco ($p < 0,05$). A Nested-PCR identificou 11 cães positivos, sendo 9 com sorologia positiva e 2 negativos. O sequenciamento de DNA foi compatível com a presença de *Ehrlichia canis*.

Palavras-chave: *Ehrlichia canis*, epidemiologia, ELISA, Nested PCR.

Abstract

The aim of this work was to study the clinical disorders and risk factors of canine ehrlichiosis in Ilhéus and Itabuna, Bahia, and compare different diagnostic methods. Blood samples were collected from 200 dogs. Each dog was clinically examined. A questionnaire was used to evaluate the risk factors. The blood samples were analyzed using the Dot-ELISA test; hematometry, platelet counts and searches for morulae on blood smears were performed. Nested PCR was carried out on 50 serologically positive samples and 50 negative samples. Three positive PCRs were sequenced. Thirty-six percent were serologically positivity and 5.5% from blood smears. The animals were anemic and thrombocytopenic. Presence of ticks and living in areas on the urban periphery were considered to be risk factors ($p < 0.05$). Nested PCR identified 11 positive dogs of which nine were serologically positive and two were negative. The DNA sequencing was consistent with the presence of *Ehrlichia canis*.

Keywords: *Ehrlichia canis*, epidemiology, ELISA, Nested PCR.

Introduction

Ehrlichia canis is the agent for canine monocytic ehrlichiosis (CME), which is transmitted by ticks, especially *Rhipicephalus sanguineus*. The first clinical manifestations occurred after an incubation period that varies from 8 to 20 days, with three distinct infection phases: acute, subclinical and chronic (HARRUS et al., 1997). Clinical signs such as fever, anorexia, vomiting, weight loss, hepatomegaly, splenomegaly, lymphadenopathy, epistaxis and hemorrhage are observed (HARRUS et al., 1997; MOREIRA et al.,

2003). Thrombocytopenia and anemia are the most common blood abnormalities in dogs that have been naturally or experimentally infected (M'GHIRBI et al., 2009).

Epidemiological studies in Brazil have revealed that the seroprevalence of CME ranges from 1.7 to 65.6% among dogs in urban or rural environments (LABARTHE et al., 2003; COSTA JUNIOR et al., 2007). Early diagnosis of ehrlichiosis is important because this is an emerging zoonotic disease (PEREZ et al., 2006). The diagnostic techniques available include blood smears to view inclusions of corpuscles or morulae (BORIN et al., 2009), serum tests, especially Dot-ELISA and the indirect immunofluorescence assay (IFA) (TRAPP et al., 2006), and molecular techniques such as nested PCR (SOUZA et al., 2010).

*Corresponding author: George Rêgo Albuquerque
Departamento de Ciências Agrárias e Ambientais, Universidade Estadual de Santa Cruz – UESC, CEP 45662-900, Ilhéus - BA, Brazil;
e-mail: gralbu@uesc.br

The aim of the present study was to describe the clinical disorders and risk factors of canine ehrlichiosis in Ilhéus and Itabuna, Bahia, and compare the different diagnostic methods.

Materials and Methods

1. Location

This study was conducted in Ilhéus (Latitude 14° 47' S and Longitude 39° 02' W) and Itabuna (latitude 14° 47' S and longitude 39° 16' W), in the Southern region of the State of Bahia, from August 2004 to April 2005. During the study period, the average temperature was 23.4 °C, with relative humidity of 84.8% and total rainfall of 1,320 mm.

2. Population studied, samples and procedures

Blood samples were collected from 200 dogs (100 dogs per city) at the Veterinary Teaching Hospital of the State University of Santa Cruz (UESC) (Ilhéus n = 35, Itabuna n = 20), at participating local veterinary clinics (Ilhéus n = 20, Itabuna n = 45), and selected households in both cities (Ilhéus n = 45, Itabuna n = 35). The animals were selected for the researchers' convenience, without specific inclusion criteria, exclusion criteria or randomization. At the local veterinary clinics, dogs were enrolled in this study when they were presented due to illness or the need for a surgical procedure or annual checkup. The participating households were invited to join in by staff at the clinics and were selected based on the owners' willingness to participate in this study. Only one dog per household was enrolled.

Any clinical abnormalities identified from each dog's history or during the physical examination were recorded. The epidemiological questionnaire was filled out by the owner, and the variables studied were gender (male, female), age (1-4 years, >4 years), contact with other dogs (yes, no), presence of ticks (yes, no), breed (purebred, mixed-breed), wellness care (yes, no), petechiae (presence, absence), apathy (presence, absence), fever (presence, absence), dehydration (presence, absence), lymphadenopathy (presence, absence), weight loss (presence, absence) and household location (urban, urban periphery). The dogs living on the urban periphery were in households in peripheral neighborhoods, i.e. outside the urban centers of the cities. Thrombocytopenia (presence, absence) and anemia (presence, absence) were included in the analysis based on the hematological tests.

Blood samples were collected into tubes with and without EDTA. Blood samples were examined for *E. canis* antibodies using a commercial kit for dot-ELISA (Snap 3DX®, IDEXX Laboratories), in accordance with the manufacturer's instructions. Blood smears were prepared, fixed in methanol for 5 minutes and stained with 10% Giemsa. Complete blood counts were performed. Dogs with counts of less than 200,000 platelets.mL⁻¹ were considered to be thrombocytopenic (HARRUS et al., 1997). Animals with packed cell volume (PCV) below 37% were considered anemic.

Fifty seropositive and 50 seronegative dogs were tested by means of nested PCR for molecular detection of *E. canis*. Only 100 PCRs were possible because of losses of several samples at the beginning

of the study. DNA was extracted from 100 mL of EDTA blood using a commercial kit (ChargeSwitch™ gDNA 50-100 mL Blood Kit, Invitrogen™, USA), in accordance with the manufacturer's instructions, thus resulting in a final DNA concentration of at least 3 mg of total extracted volume. Molecular-grade water was extracted to confirm that there was no cross-contamination between samples during DNA extraction. The genetic presence of any species of the *Ehrlichia* genus was investigated using the ECC forward primer (5'-AGAACGAACGCTGGCGGCAAGC-3') and ECB reverse primer (5'-CGTATTACCGCGGCTGCTGGCA-3'). Samples that were positive for *Ehrlichia* sp. were then tested with the ECAN5 forward primer (5'-CAATTATTTATAGCCTCTGGCTATAGGA-3') and HE3 reverse primer (5'-TATAGGTACCGTCA TTATCTTCCCTAT-3') to obtain a specific amplification of *E. canis*. The PCR mixture consisted of 5 mL of purified DNA, 0.4 mM of ECB/ECC, 200 mM of each dNTP, 5.0 mM of MgCl₂, 2.5 U of Taq DNA polymerase (Invitrogen™) and 1.6x PCR buffer (Invitrogen™), to make up a total volume of 25 mL of reaction mixture. The amplification program used to identify the genetic sequence of the genus *Ehrlichia* spp. consisted of a first step of denaturation for 3 minutes at 94 °C, followed by 35 cycles of denaturation at 94 °C for 1 minute, annealing at 68 °C for 2 minutes and extension at 72 °C for 2 minutes. For the nested PCR, 3 mL of amplicons from the first PCR and 0.2 mM of ECAN5/HE3 were used to identify the specific genetic sequence for *E. canis*, in an amplification program consisting of denaturation for 3 minutes at 94 °C, followed by 35 denaturation cycles at 94 °C for 1 minute, annealing at 58 °C for 2 minutes and extension at 72 °C for 1.5 minute. The PCR products were analyzed using 1.5% agarose gel electrophoresis, stained with ethidium bromide and photographed. The positive control used was a blood sample from a dog that was positive for *E. canis*, and which had previously been confirmed by sequencing. For the negative control, ultra-pure water was used in both PCRs.

The sequencing of the samples was carried out using the automatic sequencer ABI-PRISM 3100 Genetic Analyzer (Applied Biosystems). The DNA template (45 ng) was marked with 3.2 pmol of the primer ECAN5 and 2 mL of the reagent BigDye Terminator v3.1 Cycle Sequencing RR-100 (Applied Biosystems), in a final volume of 10 mL. The BLAST software (<http://www.ncbi.nlm.nih.gov/blast>) was used for comparison and analysis of the sequences obtained.

3. Statistical analysis

Normal data distribution was confirmed by means of the Kolmogorov-Smirnov test in the SPSS software, and statistical differences in platelet counts and PCV between *E. canis* seropositive and seronegative dogs were analyzed by means of the *t*-test, to 5% (RODRIGUEZ-VIVAS et al., 2005).

The association between presence of antibodies against *E. canis* and the study variables was evaluated by means of the chi-square (χ^2) test and Fisher's exact test. The Epi-Info software, version 3.5.1, was used to evaluate the dispersion frequency. The variables that presented *P* < 0.2 in the univariate analysis were selected for multivariate analysis by means of logistic regression, also using the Epi-Info software, version 3.5.1.

Results

Out of all the samples tested, 72 (36%) were positive for *E. canis* according to ELISA Snap 3DX[®]: 43 in Ilhéus and 29 in Itabuna (CARLOS et al., 2007). In relation to the place where the samples were collected, the rates of positive findings were 52.7% (29) at the Veterinary Hospital, 26.15% (17) for the dogs at private veterinary clinics and 32.5% (26) among samples collected from households. Evaluating according to municipality, the rates of positive findings in Ilhéus were 65.7% (23) for the samples collected at UESC, 35% (7) in private veterinary clinics and 28.9% (13) from households. In Itabuna, the rates of positive findings were 30% (6) at UESC, 22.2% (10) in veterinary clinics and 37.1% (13) from households. A statistical difference was observed in relation to the samples collected in Ilhéus, at UESC ($p = 0.02$). Only 11 dogs (5.5%) presented typical morulae in monocytes (Table 1).

The clinical examination detected fever ($n = 24$, 12.0%), apathy ($n = 13$, 6.5%), petechiae ($n = 3$, 1.5%), dehydration ($n = 13$, 6.5%), lymphadenopathy ($n = 49$, 24.5%), and weight loss ($n = 10$, 5.0%). Among the 72 seropositive animals, 10 presented fever

(13.8%), 9 apathy (12.5%), 3 petechiae (4.2%), 7 dehydration (9.7%) and 28 lymphadenopathy (38.9%). All the animals with petechiae were seropositive.

The average number of platelets was 249,000 (range: 23,000 – 628,000) platelets.mL⁻¹ of blood. The seropositive dogs presented an average count of 180,000 platelets.mL⁻¹ of blood (range: 23,000 – 590,000). The seronegative dogs presented an average of 288,000 platelets.mL⁻¹ of blood (range: 76,000 – 628,000) ($p < 0.0001$). Regarding hematocrit values, the total average was 40.5% (range: 8.0 – 60.0). The seropositive animals presented an average of 35.2% (range: 8.0 – 56.0) whereas the seronegative animals presented an average of 43.5% (range: 22.0 – 60.0) ($p < 0.0001$).

The univariate analysis used to identify risk factors for *E. canis* seropositivity are presented in Table 2. Logistic regression revealed a positive relationship between the presence of ticks and dogs living on the urban periphery (Table 3). Regarding clinical abnormalities, based on univariate analysis, seropositive animals more frequently were anemic ($p \leq 0.01$) and thrombocytopenic ($p \leq 0.01$) and presented lymphadenopathy ($p \leq 0.01$), in comparison with seronegative dogs. There were no differences in temperature between the groups (Table 4). Logistic regression revealed positive relationships with anemia, thrombocytopenia and lymphadenopathy (Table 5).

Among the 100 blood samples tested for *E. canis* DNA by means of nested PCR, 11 were considered positive. Out of these, nine were from seropositive dogs and two from seronegative dogs. All the sequenced fragments had approximately 364 base pairs (bp). These sequences had 97-99% identity with the corresponding sequences of *E. canis* strains that have been deposited at GenBank (access codes: GU386289.1; GU386285.1).

Table 1. Presence of ectoparasites, morulae and antibodies against *Ehrlichia canis* in dogs in the Southern region of Bahia.

Location	Dogs	Dogs with ticks	Presence of morulae	Presence of antibodies
Ilhéus	100	54	6	43
Itabuna	100	40	5	29
Total	200	94	11	72

Table 2. Univariate analysis on risk factors associated with seroprevalence of *Ehrlichia canis* in dogs in Ilhéus and Itabuna, Bahia.

Variable	N	Positive dogs	Prevalence (%)	OR	95% CI	P-value
Gender						
Female	93	33	35.48	-	-	-
Male	107	39	36.45	1	0.5-1.9	0.99
Age (years)						
1-4	109	36	33.03	-	-	-
>4	91	36	39.56	0.75	0.4-1.4	0.41
Breed						
Mixed breed	139	44	31.65	-	-	-
Pure breed	61	28	45.90	1.83	0.9-3.6	0.07
Ticks						
Presence	95	43	45.26	2.17	1.1-4.1	0.01
Absence	105	29	27.62	-	-	-
Household						
Urban periphery	45	27	62.5	3.67	1.7-7.7	0.0002
Urban	155	45	27.5	-	-	-
Wellness care						
Yes	138	43	31.2	0.52	0.27-1.0	0.049
No	62	29	46.8	-	-	-
Contact with other dogs						
Yes	148	54	36.5	0.92	0.4-1.7	0.94
No	52	18	34.5	-	-	-

95% CI: 95% confidence interval; OR: odds ratio (relative risk); P-value: probability.

Discussion

The results obtained from this study showed that dogs from the cities of Ilhéus and Itabuna, in the State of Bahia, Brazil, are exposed to *E. canis*. The serum prevalence in this study was similar to that found by Souza et al. (2010) using IFA, which found a prevalence of 35.6% in Salvador, Bahia. Using the Dot-ELISA test on 2,553 dogs from several Brazilian States, Labarthe et al. (2003) found that 505 (19.8%) were positive, with highest prevalence in the States of the Northeast (43%) and lowest in the Southern region (1.7%). Among the 2,553 dogs, 117 of the samples were collected in Salvador, and 42 (35.9%) were positive, which was also similar to the present study. In the same region of Bahia, with 153 different dogs, Carvalho et al. (2008) found using PCR that 12 (7.8%) were positive. In the present study, there was a statistical difference in relation to the dog samples from UESC in Ilhéus, because the majority of these animals came from the urban periphery, which was considered to be a risk factor for infection.

The serum prevalence of *E. canis* varies according to climatic conditions and certain epidemiological factors, such as the

presence and distribution of the vector, the animal's behavior, the age and type of the population under study, and the habitat (RODRIGUEZ-VIVAS et al., 2005). The dogs that lived on the urban periphery presented a higher seropositivity rate for *E. canis* (ORS= 3.7) than shown by those in the urban area. The same association was observed by Trapp et al. (2006). These results suggest that dogs that live on the urban periphery are more exposed to vector ticks, which may be related to the fact that urban dogs are more frequently sprayed with acaricides than are dogs on the urban periphery or rural dogs.

The weather is an important factor that influences the population dynamics of ticks. The average temperature in the region of Ilhéus and Itabuna provides excellent conditions for *R. sanguineus* ticks, the natural vector for *E. canis*, to develop. Ticks were found on 59.7% of the positive animals, whereas on 59.4% of the negative animals, this vector was not found. Presence of the tick as a risk factor for ehrlichiosis was also detected by Trapp et al. (2006) and Costa Junior et al. (2007).

A higher rate of seronegative animals ($p = 0.049$) was also observed among the dogs that were not provided with wellness care, thereby showing that visits to veterinary clinics were a protection factor. This is explained by the care that owners provide for their animals and the consequent lower infestation rate by the vector.

The main clinical signs observed during clinical examinations were similar to those described by Moreira et al. (2003) and Borin et al. (2009). *Ehrlichia canis* may cause variable signs, which make the diagnosis more challenging. The hematological abnormalities identified in the present study (thrombocytopenia and anemia) were similar to those previously described by Rodriguez-Vivas et al. (2005), Borin et al. (2009) and M'Ghirbi et al. (2009). The effects of the mononuclear phagocytic system, cell lysis mediated by the complement system and suppression of erythropoiesis at the bone marrow may be the mechanisms responsible for the anemia

Table 3. Final result from logistic regression on risk factors associated with *Ehrlichia canis* exposure in dogs in Ilhéus and Itabuna, Bahia.

Variable	OR	95% CI	P-value
Household			
Urban periphery	1.00	-	-
Urban	0.30	0.15 - 0.62	0.001
Ticks			
No	1.00	-	-
Yes	1.81	0.98 - 3.35	0.05

95% CI: 95% confidence interval; OR: odds ratio (relative risk); P-value: probability.

Table 4. Univariate analysis on clinical and hematological abnormalities in dogs exposed to *Ehrlichia canis* in Ilhéus and Itabuna, Bahia.

Variable	N	+	Prevalence (%)	OR	95% CI	P-value
Fever						
Yes	24	10	41.7	-	-	-
No	176	62	35.2	0.76	0.30-1.98	0.69
Petechiae						
Yes	3	3	100.0	-	-	-
No	197	69	35.0	-	-	0.045*
Apathy						
Yes	13	9	69.2	4.43	1.18-17.87	0.022
No	187	63	33.7	-	-	-
Lymph nodes						
Normal	151	44	29.1	-	-	-
Increased	49	28	57.1	3.24	1.58-6.67	0.0007
Thrombocytopenia						
Yes	76	48	63.2	7.14	3.58-14.38	<0.0001
No	124	24	19.4	-	-	-
Anemia						
Yes	78	42	53.8	3.58	1.87-6.88	<0.0001
No	122	30	24.6	-	-	-
Dehydration						
Yes	13	7	53.8	-	-	-
No	187	65	34.8	0.45	0.14-1.41	0.27

95% CI: 95% confidence interval; OR: odds ratio (relative risk); P-value: probability. * Fisher's exact test.

Table 5. Final result from logistic regression on clinical and hematological abnormalities in dogs exposed to *Ehrlichia canis* in Ilhéus and Itabuna, Bahia.

Variable	OR	95% CI	P-value
Anemia			
Yes	2.25	1.13-4.51	0.02
No	1.00	-	-
Thrombocytopenia			
Yes	6.60	3.32-13.12	<0.0001
No	1.00	-	-
Lymph nodes			
Normal	0.34	0.15-0.75	0.007
Increased	1.00	-	-

95% CI: 95% confidence interval; OR: odds ratio (relative risk); P-value: probability.

relating to the disease (MOREIRA et al., 2003). *Ehrlichia canis* induces thrombocytopenia through destruction and consumption of platelets, increased hepatic or splenic platelets sequestration, decreased platelet production due to bone marrow hypoplasia (WOODY; HOSKINS, 1991) and production of antiplatelet antibodies (GAUNT et al., 2010).

Diagnosing of ehrlichiosis through blood smear analysis is difficult because intracytoplasmic morulae are only occasionally seen during the acute phase of the disease (RODRIGUEZ-VIVAS et al., 2005). In the present study, we observed that 5.5% of the blood smears presented the typical morulae of *E. canis*, and this was similar to the observations of Rodriguez-Vivas et al. (2005), who found that 44.1% of the dogs were seropositive but that only six dogs (5%) presented morulae detected in blood smears.

Serological tests (dot-ELISA and IFA) are the method most commonly used for veterinary diagnosis. However, the fact that an animal is seropositive does not mean that it is sick. Since the presence of antibodies reveals exposure to the agent, PCR may help in reaching a diagnostic conclusion. Souza et al. (2010) found that 35.6% (168/472) of the dogs examined were seropositive, but that only 58 (34.5%) of these animals were PCR-positive.

In the present study, nested PCR was used on 100 samples (50 seropositive and 50 seronegative for *E. canis*), and nine dogs were found to be PCR-positive among the seropositive dogs. It was found that two dogs were seronegative for *E. canis*, but PCR-positive. Similar results were obtained by Diniz et al. (2007). Hyperacute cases may be PCR-positive with no antibody response, in which the molecular test detects the presence of the agent before the dog's immunological response occurs. Nested PCR is the technique most used for diagnosing *E. canis*, but false-positive results may occur if primers that generate unspecific fragments are used. In the present study, such results did not occur because the primers used were ECAN5 and HE-3. In sequencing the *E. canis* DNA fragments using the ECAN5 primer, the homology ranged from 97 to 99%. This low homology may have occurred because of the few samples sequenced (three), as well as incomplete DNA purification prior to sequencing.

In the present study, subject enrollment done according to convenience, because of certain constraints, such as structural and financial limitations. Despite the lack of random sampling in this study, the results provide important information about disease prevalence and the risk factors and clinical abnormalities associated with *E. canis* infection in the Southern region of Bahia.

References

- BORIN, S.; CRIVELENTI, L. Z.; FERREIRA, F. A. Aspectos epidemiológicos, clínicos e hematológicos de 251 cães portadores de mórula de *Ehrlichia* spp. naturalmente infectados. **Arquivo Brasileiro de Medicina Veterinária e Zootecnia**, v. 61, n. 3, p. 566-571, 2009.
- CARLOS, R. S. A. et al. Frequência de anticorpos anti-*Ehrlichia canis*, *Borrelia burgdorferi* e antígenos de *Dirofilaria immitis* em cães na microrregião Ilhéus-Itabuna, Bahia, Brasil. **Revista Brasileira de Parasitologia Veterinária**, v. 16, n. 3, p. 117-120, 2007.
- CARVALHO, F. S. et al. Epidemiological and molecular study of *Ehrlichia canis* in dogs in Bahia, Brazil. **Genetics and Molecular Research**, v. 7, n. 3, p. 657-662, 2008. PMID:18752193. <http://dx.doi.org/10.4238/vol7-3gmr468>
- COSTA JUNIOR, L. M. et al. Sero-prevalence and risk indicators for canine ehrlichiosis in three rural areas of Brazil. **Veterinary Journal**, v. 174, n. 3, p. 673-676, 2007. PMID:17204439. <http://dx.doi.org/10.1016/j.tvjl.2006.11.002>
- DINIZ, P. P. V. et al. Surveillance for zoonotic vector-borne infections using sick dogs from southeastern Brazil. **Vector-Borne and Zoonotic Diseases**, v. 7, n. 4, p. 689-697, 2007.
- GAUNT, S. D. et al. Experimental infection and co-infection of dogs with *Anaplasma platys* and *Ehrlichia canis*: hematologic, serologic and molecular findings. **Parasites and Vectors**, v. 3, n. 1, p. 33, 2010. <http://dx.doi.org/10.1186/1756-3305-3-33>
- HARRUS, S. et al. Canine monocytic ehrlichiosis: an update. **Parasitology**, v. 19, p. 431-444, 1997.
- LABARTHE, N. et al. Serologic prevalence of *Dirofilaria immitis*, *Ehrlichia canis*, and *Borrelia burgdorferi* infections in Brazil. **Veterinary Therapeutics**, v. 4, p. 67-75, 2003.
- M'GHIRBI, Y. et al. Clinical, serological and molecular evidence of ehrlichiosis and anaplasmosis in dogs in Tunisia. **Parasitology Research**, v. 104, n. 4, p. 767-774, 2009.
- MOREIRA, S. M. et al. Retrospective study (1998-2001) on canine ehrlichiosis in Belo Horizonte, MG, Brazil. **Arquivo Brasileiro de Medicina Veterinária e Zootecnia**, v. 55, n. 2, p. 141-147, 2003.
- PEREZ, M. et al. Human infection with *Ehrlichia canis* accompanied by clinical signs in Venezuela. **Annals of New York Academic Science**, v. 1078, p. 110-117, 2006. PMID:17114689. <http://dx.doi.org/10.1196/annals.1374.016>
- RODRIGUEZ-VIVAS, R. I.; ALBORNOZ, R. E. F.; BOLIO, G. M. E. *Ehrlichia canis* in dogs in Yucatan, México: seroprevalence, prevalence of infection and associated factors. **Veterinary Parasitology**, v. 127, n. 1, p. 75-79, 2005. PMID:15619376. <http://dx.doi.org/10.1016/j.vetpar.2004.08.022>
- SOUZA, B. M. P. S. et al. Prevalence of ehrlichial infection among dogs and ticks in Northeastern Brazil. **Revista Brasileira de Parasitologia Veterinária**, v. 19, n. 2, p. 89-93, 2010.
- TRAPP, S. M. et al. Seroepidemiology of canine babesiosis and ehrlichiosis in a hospital population. **Veterinary Parasitology**, v. 140, n. 3-4, p. 223-230, 2006. PMID:16647817. <http://dx.doi.org/10.1016/j.vetpar.2006.03.030>
- WOODY, B. J.; HOSKINS, J. D. Ehrlichial diseases of dogs. **Veterinary Clinics of North America: Small Animal Practice**, v. 21, n. 1, p. 75-98, 1991. PMID:2014630.