



Revista Brasileira de Parasitologia
Veterinária

ISSN: 0103-846X

zacariascbpv@fcav.unesp.br

Colégio Brasileiro de Parasitologia
Veterinária
Brasil

da Fonseca Fernandes, Annielle Regina; Rodrigues Menezes Pimenta, Carla Lauise;
Fernandes Vidal, Ivana; Capriogli Oliveira, Gabriela; Saran Sartori, Raissa; Barros Araújo,
Raizza; Almeida Melo, Márcia; Langoni, Hélio; Santos Azevedo, Sérgio
Risk factors associated with seropositivity for *Leishmania* spp. and *Trypanosoma cruzi* in
dogs in the state of Paraíba, Brazil
Revista Brasileira de Parasitologia Veterinária, vol. 25, núm. 1, enero-marzo, 2016, pp. 90
-98
Colégio Brasileiro de Parasitologia Veterinária
Jaboticabal, Brasil

Available in: <http://www.redalyc.org/articulo.oa?id=397844775010>

- How to cite
- Complete issue
- More information about this article
- Journal's homepage in redalyc.org

redalyc.org

Scientific Information System

Network of Scientific Journals from Latin America, the Caribbean, Spain and Portugal

Non-profit academic project, developed under the open access initiative

Risk factors associated with seropositivity for *Leishmania* spp. and *Trypanosoma cruzi* in dogs in the state of Paraíba, Brazil

Fatores de risco associados às soropositividades para *Leishmania* spp. e *Trypanosoma cruzi* em cães no Estado da Paraíba, Brasil

Annielle Regina da Fonseca Fernandes¹; Carla Lauise Rodrigues Menezes Pimenta¹; Ivana Fernandes Vidal¹; Gabriela Capriogli Oliveira²; Raissa Saran Sartori²; Raizza Barros Araújo³; Márcia Almeida Melo³; Hélio Langoni²; Sérgio Santos Azevedo^{1*}

¹ Laboratório de Doenças Transmissíveis, Programa de Pós-graduação em Medicina Veterinária, Universidade Federal de Campina Grande – UFCG, Patos, PB, Brasil

² Laboratório de Zoonoses, Faculdade de Medicina Veterinária e Zootecnia, Universidade Estadual Paulista “Júlio de Mesquita Filho” – UNESP, Botucatu, SP, Brasil

³ Laboratório de Biologia Molecular do Semi-árido, Programa de Pós-graduação em Medicina Veterinária, Universidade Federal de Campina Grande – UFCG, Patos, PB, Brasil

Received January 19, 2016

Accepted February 16, 2016

Abstract

The aim of this survey was to determine the seropositivity and risk factors for *Leishmania* spp. and *Trypanosoma cruzi* in dogs in the State of Paraíba, Northeastern Brazil. A total of 1,043 dogs were tested, and the serological diagnoses of Chagas disease (CD) and canine visceral leishmaniasis (CVL) was performed by the indirect fluorescent antibody test (IFAT). Animals that tested seropositive for both diseases (by IFAT) were further subjected to ELISA. Of the 1,043 dogs 81 (7.8%; 95% CI = 6.1-9.4%) tested seropositive for *Leishmania* spp., while 83 were seropositive for *T. cruzi* (7.9%; 95% CI = 6.3-9.6%). Simultaneous serological reactions were detected in 49 animals (4.6%; 95% CI = 3.6-6.2%). Semi-domiciled housing (OR = 2.044), free housing (OR = 4.151), and soil (OR = 3.425) and soil/cement (OR = 3.065) environmental conditions were identified as risk factors for CVL seropositivity. The risk factors identified for CD seropositivity were semi-domiciled (OR = 2.353) or free housing (OR = 3.454), and contact with bovine (OR = 2.015). This study revealed the presence of dogs in the Paraíba State seropositive for CVL and CD, suggesting the need for revisiting and intensification of disease control measures through constant monitoring of the canine population.

Keywords: Dogs, *Leishmania infantum*, *Trypanosoma cruzi*, risk factors, Northeastern region of Brazil.

Resumo

O objetivo do presente trabalho foi determinar a soropositividade para *Leishmania* spp. e *Trypanosoma cruzi* em cães do Estado da Paraíba, Nordeste do Brasil, bem como identificar fatores de risco. Foram utilizados 1.043 cães e, para o diagnóstico sorológico de doença de Chagas (DC) e leishmaniose visceral canina (LVC), foi utilizada a reação de imunofluorescência indireta (RIFI). Animais positivos para ambas as doenças (pela RIFI) foram submetidos ao ELISA. Dos 1.043 cães investigados, 81 foram soropositivos para *Leishmania* spp., resultando em prevalência de 7,8% (IC 95% = 6,1-9,4%) e, para *T. cruzi*, 83 (7,9%; IC 95% = 6,3-9,6%) animais foram soropositivos. Quarenta e nove animais (4,6%; IC 95% = 3,6-6,2%) apresentaram sororeatividade mista. Criação semidomiciliar (OR = 2,044), criação solta (OR = 4,151), ambiente de terra (OR = 3,425) e ambiente de terra/cimento (OR = 3,065) foram apontados como fatores de risco para LVC, e criação semidomiciliar (OR = 2,353), criação solta (OR = 3,454) e contato com bovinos (OR = 2,015) para DC. Conclui-se que LVC e DC estão presentes em cães do Estado da Paraíba, o que sugere revisão e intensificação das medidas de controle através do constante monitoramento da população canina.

Palavras-chave: Cães, *Leishmania infantum*, *Trypanosoma cruzi*, fatores de risco, Nordeste do Brasil.

*Corresponding author: Sérgio Santos Azevedo. Laboratório de Doenças Transmissíveis, Unidade Acadêmica de Medicina Veterinária, Universidade Federal de Campina Grande – UFCG, Av. Universitária, s/n, Santa Cecília, CEP 58700-970, Patos, PB, Brasil. e-mail: sergio@vps.fmvz.usp.br

Introduction

The zoonoses visceral leishmaniasis (VL) and Chagas disease (CD) have been known to significantly affect human health in Brazil, prompting the need for repeated medical assistance. These diseases are caused by the protozoans *Leishmania infantum* and *Trypanosoma cruzi*, respectively; both are known to be carried by hematophagous insect (*Lutzomyia* spp. and *Triatoma* spp.) vectors. Wild and domesticated canids have been identified as the reservoirs of these parasites (SIMÕES-MATTOS et al., 2005; LUCIANO et al., 2009).

Canine visceral leishmaniasis (CVL) is an important zoonosis, which is associated with rapid geographical expansion. This disease has been observed in 47 countries, and is caused by species *Leishmania (L.) infantum* (KUHLS et al., 2011). The disease is widely distributed throughout Brazil, and presents specific geographical, climatic, and social characteristics. CVL, which was mostly diagnosed in rural areas in the past, has recently migrated to medium and large urban areas, which has led to changes in its epidemiological profile. In Brazil, CVL occurs in the central-western, southeastern, northern, and northeastern regions, however, the majority of cases have been reported in northeastern region (BRASIL, 2006).

Dogs are the major domestic reservoirs of VL, and play a major role in maintaining the disease cycle (MELO, 2004). Their relevance is attributed to the greater prevalence of VL in the canine than in the human population, since infections in humans are often preceded by infections in dogs. Furthermore, dogs carry a greater number of parasites on their skin than humans, which favors the infection of the vectors (CASTRO, 1996; SANTA ROSA & OLIVEIRA, 1997; BANETH, 2006).

CD, also known as American trypanosomiasis, is a major public health concern in several countries around the world (BORCHHARDT et al., 2010). According to the World Health Organization (WHO), over 6 million people from 21 countries are estimated to be infected with CD, with an annual incidence of 100,000 to 200,000 cases (WHO, 2015). With regard to Brazil, it is estimated that the number of infected individuals is around of three million, and this zoonotic disease is present in the list of neglected tropical diseases (DIAS, 2011). Despite the main reservoirs of CD being wild species, cats and dogs are known to get infected by the causative protozoan; this plays an important role in the ecology and epidemiology of this disease (GÜRTLER et al., 2007). The natural infection of dogs by *T. cruzi* occurs in a manner similar to human infections, occurring either through active transmission by the vector, or through contamination of the skin and/or conjunctiva by infected feces. However, Barr (2006) stated that the transmission frequently occurs through the ingestion of infected vectors or infected tissues from rodents or other wild animals found in around shelters/residences.

The phylogenetic proximity between the parasites, and the fact that both diseases are endemic to some regions of South America, necessitate the analysis of the two infections in parallel. These zoonoses must be monitored and controlled through surveys that combine serological and epidemiological approaches for each geographical location, as these strategies can lead to the allocation

of specific funds that will allow policy-makers to organize and direct new policies towards strengthening public health as a whole.

The aforementioned reasons, the effect of CVL and CD on public health, the role of dogs as parasite reservoirs, and the scarce evidence-based data available in the State of Paraíba, have motivated this study, which aims to determine the seropositive and associated risk factors for *Leishmania* spp. and *T. cruzi* in the canine population of this region.

Materials and Methods

Ethics Committee

This work was approved by the Ethics Committee on Animal Use (CEUA/CESED), Faculty of Medical Sciences of Campina Grande-FCM, under the code 0041/280314.

Samples

Dogs older than three months were included in this study; the test subjects were identified by visiting the residences of their owners, and from those admitted to veterinary clinics and analysis laboratories. Animals from the João Pessoa, Campina Grande, Patos, Sousa, and Cajazeiras counties (Figure 1), five regional urban centers in the State of Paraíba situated along one of the major highways (BR-230; also known as the Trans-Amazonian highway), were included in this study. The sample size was determined according to the total population of dogs (141,863 animals) in these counties (6,843 dogs in Sousa, 10,553 in Patos, 78,073 in João Pessoa, 6,103 in Cajazeiras, and 40,291 in Campina Grande). These numbers were estimated based on human population data for the year 2013, provided by the Brazilian Institute of Geography and Statistics (IBGE, 2013). The dog/human distribution in urban areas was calculated at a ratio of 1:10 (WHO, 1990; REICHMANN et al., 1999). The sample size was determined for each county based on an estimated seropositivity of 50% (value adopted for sample maximization), a confidence level of 95%, and an error of 10% (THRUSFIELD, 2007); this provided the required sample size of at least 96 animals per county. Ultimately, 1,043 animals were included in this study (125 dogs in Sousa, 206 in Patos, 338 in João Pessoa, 125 in Cajazeiras, and 249 in Campina Grande).

Probabilistic criteria were not established for animal selection, i.e., inclusion of the animals depended on previous contact with the owners and their agreement to taking part in the study. Blood samples were collected from the external jugular or cephalic veins using 5 mL disposable syringes, in the period from January 2013 to June 2014; the serum samples were stored at -20 °C until serological tests.

Serology

Serum antibodies for both diseases were searched by indirect fluorescent antibody test (IFAT), using the protocol by Camargo (1966); the samples were diluted as follows: 1:40, 1:80, 1:160, 1:320, and 1:640. The antigen used to coat the slides for CVL diagnosis




Figure 1. Geographical distribution of counties used. Detail shows the State of Paraíba within Brazil.

was prepared from *L. major*-like promastigotes, whereas the antigen for CD was prepared from *T. cruzi* (strain Y) epimastigotes; both cultures were maintained in LIT (Liver Infusion Triptose) and NNN (Neal, Novy, Nicolle) culture media. Positive and negative control sera, for both parasites, were provided by the Núcleo de Pesquisa em Zoonoses (NUPEZO), Universidade Estadual Paulista “Júlio de Mesquita Filho” (UNESP), Botucatu – SP. Based on the results obtained for the controls, the final antibody titer was determined to correspond to the highest dilution of the sera; under these conditions, the membranes of at least 50% of the promastigotes (CVL) and epimastigotes (CD) emitted readable fluorescence, with a cutoff of 40 or higher.

Animals that tested seropositive for both parasites (by IFAT) were further subjected to enzyme-linked immunosorbent assay (ELISA) using the ELISA S7® kit (Biogene Indústria & Comercio Ltda ME, Recife-PE, Brazil) for the serological diagnosis of CVL. ELISA was performed as per the manufacturer protocols in order to minimize cross-reactivity and accurately detect possible co-infections.

Serum samples that showed positive results at the 1:40 or higher dilutions by IFAT, and which were also determined to be reactive by ELISA, were established to be positive for *Leishmania* spp. Dogs were determined to be positive for *T. cruzi* when results from the 1:40 or higher serum dilutions were detected by IFAT. Simultaneous serological reactions were diagnosed when the animal was seropositive for both parasites.

Epidemiological questionnaires

During the collection of blood samples from the dogs, their owners were also provided with an epidemiological questionnaire. This questionnaire requested information regarding a series of variables, in order to investigate certain behaviors and conditions that could act as risk factors for CVL and CD. The variables analyzed and respective categories were as follow:

- Owner's information: county of origin (Sousa, Patos, João Pessoa, Cajazeiras, Campina Grande), level of education (illiterate, 1st degree, 2nd degree, 3th degree), and traveling with the dog (yes, no);
- Animal's information: gender (male, female), age (≤ 12 months, 13-48 months, 49-72 months, > 72 months), breed (undefined, defined), conditions of housing (domiciled environment [dogs without access to streets], semi-domiciled [dogs with restricted access to streets], free [dogs with unrestricted access to streets]), and dog food (commercial, homemade, homemade meals, a combination of commercial and homemade, a combination of commercial and homemade meals);
- Environment's information: contact with other dogs (yes, no), contact with bovine (yes, no), contact with horses (yes, no), contact with cats (yes, no), contact with

goat/sheep (yes, no), contact with pigs (yes, no), contact with wild animals (yes, no), environmental conditions (soil, cement, soil and cement), environmental hygiene (yes, no), presence of rodents (yes, no), and access to water dams (yes, no).

Risk factor analysis

The risk factors were analyzed from the data obtained by the epidemiological questionnaires, using univariable approaches. Two groups of animals, seropositive and seronegative dogs, were formed for univariable analysis; these were compared with the tested variables. Variables with $p \leq 0.2$, determined by the chi-square or Fisher's exact tests (ZAR, 1999), were selected for multivariable analysis, using multiple logistic regression (HOSMER & LEMESHOW, 2000). The significance level considered to discarding a determined variable was 5%. The collinearity among independent variables was assessed using correlation analysis, and when two variables were highly collinear (correlation coefficient > 0.90), only one variable was likely to enter into analysis. In such situations, selection of which collinear variable to put into the model was guided by biological plausibility (DOHOO et al., 1997). The tests were performed using the SPSS software package, version 13.0 for Windows (SPSS, Inc., Chicago, IL, USA).

Results

Of the 1,043 dogs included in the study, 81 tested seropositive for *Leishmania* spp. (7.8%; 95%CI = 6.1-9.4%). The following antibody titers were detected: 38 (46.9%), 15 (18.5%), 11 (13.5%), 8 (9.8%), and 9 (11.1%) dogs had titers of 40, 80, 160, 320, and 640, respectively. Eighty-three (7.9%; 95% CI = 6.3-9.6%) dogs tested seropositive for *T. cruzi*; forty-five (54.2%) animals displayed antibodies at a serum titer of 40, while 12 (14.4%), 9 (10.8%), 10 (12%), and 7 (8.4%) displayed antibodies at titers of 80, 160, 320 and 640, respectively. Simultaneous serological reactions were detected in 49 (4.6%; 95% CI = 3.6-6.2%) dogs. Table 1 shows the distribution of seropositive animals according to the county of origin.

Univariable analysis (Table 2) of the risk factors for CVL seropositivity ($p \leq 0.2$) focused on the following variables: county of origin ($p < 0.001$), level of education of the owner ($p = 0.055$),

breed ($p < 0.001$), conditions of housing ($p < 0.001$), dog food ($p < 0.001$), contact with other dogs ($p < 0.001$), contact with bovine ($p < 0.001$), contact with cats ($p < 0.001$), contact with goat/sheep ($p < 0.001$), contact with wild animals ($p < 0.001$), environmental conditions ($p < 0.001$), environmental hygiene ($p = 0.001$), traveling with the dog ($p = 0.047$), presence of rodents ($p = 0.138$), and access to water dams ($p < 0.001$). Semi-domiciled housing (OR = 2.044), free housing (OR = 4.151), and soil (OR = 3.425) and soil/cement (OR = 3.065) environmental conditions were identified as risk factors for CVL seropositivity in the logistic regression analysis (Table 3).

Univariable analysis (Table 2) of the risk factors for CD seropositivity ($p \leq 0.2$) focused on the following variables: county of origin ($p < 0.001$), level of education of the owner ($p = 0.043$), age ($p = 0.194$), breed ($p < 0.004$), housing conditions ($p < 0.001$), dog food ($p = 0.056$), contact with other dogs ($p = 0.002$), contact with bovine ($p < 0.001$), contact with cats ($p = 0.001$), contact with goat/sheep ($p < 0.001$), contact with wild animals ($p < 0.006$), environmental conditions ($p = 0.009$), environmental hygiene ($p = 0.025$), and access to water dams ($p < 0.001$). Logistic regression identified the following risk factors (Table 3): semi-domiciled (OR = 2.353) or free housing (OR = 3.454), and contact with bovine (OR = 2.015).

Discussion

Several surveys have been conducted in Brazil with the aim of establishing the prevalence of CVL; these studies have produced variable results depending on the characteristics of the study population and the methods used. Alves et al. (1998), Martins (2008), and Barboza et al. (2009), who conducted studies in Fortaleza, CE, Maceió, AL, and Salvador, BA, observed disease frequencies of 1.59%, 1.9%, and 0.7%, respectively; these disease frequencies were lower than the ones reported in this study. In contrast, similar disease frequency was reported by Azevedo et al. (2008) in a survey conducted in Poxoréo, MT (7.8%). The highest disease frequencies were identified by Matos et al. (2006) in animals admitted to the UFERSA veterinary hospital, Mossoró, RN (28%); in addition, Amora et al. (2006), observed very high disease frequencies of 45% and 34% in rural and urban areas of Mossoró, RN, respectively, while Abreu-Silva et al. (2008), Almeida et al. (2012), and Morais et al. (2013) observed high disease frequencies in São Luiz, MA (51.6%), Cuiabá, MT

Table 1. Seropositivity for visceral leishmaniasis, Chagas disease and both diseases in dogs in the period from January 2013 to June 2014, in the State of Paraíba, Brazil.

County	Number of dogs		Leishmaniasis			Chagas disease			Both diseases		
	Total	Sampled	Positive	%	95% CI	Positive	%	95% CI	Positive	%	95% CI
Sousa	6,843	125	9	7.2	2.7-11.7	6	4.8	1.1-8.5	5	4.0	1.7-9.0
Patos	10,553	206	38	18.4	13.1-23.7	39	18.9	13.6-24.3	23	11.1	7.6-16.2
João Pessoa	78,073	338	20	5.9	3.4-8.4	16	4.7	2.5-7.0	10	2.9	1.6-5.4
Cajazeiras	6,103	125	5	4.0	1.7-9.2	7	5.6	1.6-9.6	4	3.2	1.3-7.9
Campina Grande	40,291	249	9	3.6	1.3-5.9	15	6.0	3.1-9.0	7	2.8	1.4-5.7
Total	141,863	1,043	81	7.8	6.1-9.4	83	7.9	6.3-9.6	49	4.6	3.6-6.2

95% CI: 95% confidence interval.

Table 2. Univariable analysis for risk factors associated with the seropositivity for visceral leishmaniasis and Chagas disease in dogs in the period from January 2013 to June 2014, in the State of Paraíba, Brazil.

Variable	Category	Total number of dogs	Leishmaniasis		Chagas disease	
			Positive dogs (%)	p	Positive dogs (%)	p
County of origin	Sousa	125	9 (7.2)		6 (4.8)	
	Patos	206	38 (18.4)		39 (18.9)	
	João Pessoa	338	20 (5.9)		16 (4.7)	
	Cajazeiras	125	5 (4.0)		7 (5.6)	
	Campina	249	9 (3.6)	<0.001	15 (6.0)	<0.001
Level of education of the owner	Illiterate	20	3 (15.0)		2 (10.0)	
	1 st degree	272	22 (8.1)		21 (7.7)	
	2 nd degree	474	44 (9.3)		48 (10.1)	
	3 rd degree	277	12 (4.3)	0.055*	12 (4.3)	0.043*
Gender	Male	554	48 (8.7)		48 (8.7)	
	Female	489	33 (6.7)	0.299	35 (7.2)	0.434
Age	≤ 12 months	345	22 (6.4)		21 (6.1)	
	13-48 months	538	46 (8.6)		46 (8.6)	
	49-72 months	97	10 (10.3)		12 (12.4)	
	> 72 months	63	3 (4.8)	0.382	4 (6.3)	0.194*
Breed	Undefined	567	60 (10.6)		58 (10.2)	
	Defined	476	21 (4.4)	<0.001	25 (5.3)	0.004*
Condition of housing	Domiciled	724	30 (4.1)		35 (4.8)	
	Semi-domiciled	195	22 (11.3)		23 (11.8)	
	Free	124	29 (23.4)	<0.001	25 (20.2)	<0.001
Dog food	Commercial	411	17 (4.1)		23 (5.6)	
	Homemade	104	20 (19.2)		14 (13.5)	
	Homemade meals	283	22 (7.8)		21 (7.4)	
	Commercial + homemade	49	5 (10.2)		5 (10.2)	
	Commercial + homemade meals	196	17 (8.7)	<0.001	20 (10.2)	0.056*
Contact with other dogs	No	451	19 (4.2)		22 (4.9)	
	Yes	592	62 (10.5)	<0.001	61 (10.3)	0.002*
Contact with bovine	No	960	59 (6.1)		63 (6.6)	
	Yes	83	22 (26.5)	<0.001	20 (24.1)	<0.001
Contact with horses	No	1031	79 (7.7)		81 (7.9)	
	Yes	12	2 (16.7)	0.238	2 (16.7)	0.246
Contact with cats	No	805	44 (5.5)		51 (6.3)	
	Yes	238	37 (15.5)	<0.001	32 (13.4)	0.001*
Contact with goat/sheep	No	972	61 (6.3)		67 (6.9)	
	Yes	71	20 (28.2)	<0.001	16 (22.5)	<0.001
Contact with pigs	No	1034	81 (7.8)		82 (7.9)	
	Yes	9	0 (0.0)	1.000	1 (11.1)	0.527

*Variables selected for the multiple analysis ($p \leq 0.2$); p: probability of casual occurrence.

Table 2. Continued...

Variable	Category	Total number of dogs	Leishmaniasis		Chagas disease	
			Positive dogs (%)	P	Positive dogs (%)	P
Contact with wild animals	No	911	60 (6.6)		64 (7.0)	
	Yes	132	21 (15.9)	<0.001	19 (14.4)	0.006*
Environmental conditions	Soil	198	24 (12.1)		17 (8.6)	
	Soil/cement	398	46 (11.6)		43 (10.8)	
	Cement	447	11 (2.5)	<0.001	23 (5.1)	0.009*
Environmental hygiene	No	137	21 (15.3)		18 (13.1)	
	Yes	906	60 (6.6)	0.001*	65 (7.2)	0.025*
Traveling with the dog	No	947	79 (8.3)		78 (8.2)	
	Yes	96	2 (2.1)	0.047*	5 (5.2)	0.397
Presence of rodents	No	771	66 (8.6)		66 (8.6)	
	Yes	272	15 (5.5)	0.138*	17 (6.3)	0.280
Access to water dams	No	867	53 (6.1)		56 (6.5)	
	Yes	176	28 (15.9)	<0.001	27 (15.3)	<0.001

*Variables selected for the multiple analysis ($p \leq 0.2$); p: probability of casual occurrence.

Table 3. Risk factors associated with the seropositivity for visceral leishmaniasis and Chagas disease in dogs in the period from January 2013 to June 2014, in the State of Paraíba, Brazil.

Risk factor	Odds ratio (OR)	95% IC	P
Leishmaniasis			
Semi-domiciled housing	2.044	1.107-3.777	0.022
Free housing	4.151	2.046-8.423	<0.001
Environmental condition (soil)	3.425	1.514-7.747	0.003
Environmental condition (soil/cement)	3.065	1.493-6.290	0.002
Chagas disease			
Semi-domiciled housing	2.353	1.331-4.159	0.003
Free housing	3.454	1.740-6.854	<0.001
Contact with bovine	2.015	1.005-4.039	0.048

95% CI: 95% confidence interval; probability of casual occurrence.

(22.1%), and Araguaína, TO (51.35%), respectively. However, it should be emphasized that a majority of these studies were conducted in single locations, whereas the current study focused on five different urban hubs in the State of Paraíba, which could explain the differences in seropositivity. Rondon et al. (2008) reported that the distribution of CVL frequency suggested a seasonal variation, which was caused by the high and low peaks of the vector population, which could justify the variability of data for CVL prevalence throughout Brazil.

Paraíba is an endemic region for CVL, and according to the SINAN Health Information System (Sistema de Informação de Agravos de Notificação), 162 cases of human visceral leishmaniasis have been recorded between 2007 and 2013 (BRASIL, 2014) in Paraíba. Therefore, the measures developed for the control of CVL and VL must be revisited by the responsible policy-makers to facilitate constant monitoring of the canine population for

the presence of anti-*L. chagasi* antibodies, in order to prevent transmission to humans.

It should be highlighted that the criterion adopted for positive serological results in this study (IFAT associated with ELISA, instead of IFAT alone) may have influenced the results. Because of the controversy and the fact that many owners are reluctant to euthanize their animals, since the adoption of that measure has no sufficient effectiveness in reducing the prevalence of cases of the disease, there is the need to carry out more than one test for the confirmatory diagnosis of this disease as well as a standardization of diagnostic techniques to be used both in surveys and in individual cases. This question becomes even more delicate when many of the animals are asymptomatic.

The housing conditions of the dogs, particularly semi-domiciled and free environments, have previously been identified as risk factors for *L. chagasi* infection by Oliveira & Araújo (2003),

Amora et al. (2006), and Naveda et al. (2006), in Feira de Santana, BA, Mossoró, RN, and Pedro Leopoldo, MG, respectively; these studies have strongly suggested the greater exposure of free animals to the vector. These results are also in agreement with the results obtained by Uchôa et al. (2001), who reported that the lack of organized human occupation (proximity to hillsides and/or forest areas) caused an environmental imbalance that favored the occurrence of disease cycle outside the forests, and closer to the urban areas. The variables related to the environmental conditions (soil, soil/cement), also identified as risk factors for CVL, suggested that a strong presence of organic materials contributes to the proliferation of synanthropic species, in addition to creating a favorable habitat for the spread of the vector (as eggs are usually laid on organic materials).

For DC, 7.9% of the animals were seropositive for *T. cruzi*. This value differs from that (22.7%) described by Souza et al. (2009), which analyzed the seroprevalence of *T. cruzi* infection in dogs from Mato Grosso do Sul using IFAT and ELISA. Mendes et al. (2013) found a prevalence of 4.08% using IFAT, ELISA and indirect hemagglutination (IHA), in Patos, PB. The higher frequencies could probably be attributed to the predominantly rural survey areas, which are the usual ecotopes of disease vectors. Silva & Fernandes (2013) utilized IFAT and ELISA and identified a prevalence for CD of 31% in domiciled dogs in São Domingos do Capim, PA. Silva et al. (2014) used IFAT to detect a CD prevalence of 22.2% in a rural area of Bragança, PA. These values, which are higher than the ones reported in this study, could be attributed to the prior instances of human CD infections in both locations. Despite the low positivity reported in this study, the general population and the authorities must be alerted to the presence of infectious agents, in order to plan and establish epidemiological strategies for the effective control of CD.

The housing conditions of the dogs, particularly semi-domiciled and free environments, have also been identified as risk factors for *T. cruzi*. This is a reason for concern; despite the major vector for this disease (*Triatoma infestans*) being no longer responsible for CD infections in Brazil (ARGOLO et al., 2008), other species could contribute to its spread in the country. *Triatoma* sp. insects are currently migrating to urban areas, and have contributed to the strengthening of the disease cycle outside the forest (and closer to urban areas), thereby infecting dogs (especially the ones in free environments) and humans. Contact with bovine is another risk factor for CD. Many wild and domestic mammals are known reservoirs of CD; although dogs are the major domestic reservoir for human infection, other animals can also contribute to CD ecology (DIAS & COURA, 1997). Furthermore, it is usual the occurrence of bovine roaming free in many urban areas of the State of Paraíba. Dias et al. (2000) have reported the decades-old existence of peri-urban disease foci, and have suggested that the constant migration from rural to urban areas, and the poverty and semi-rural characteristics of the periphery neighborhoods determine disease distribution.

According to the criteria established in this study, 4.6% (49/1043) of the dogs tested positive (by IFAT assay) for *Leishmania* spp. and *T. cruzi*, possibly indicating the occurrence of mixed (simultaneous) infections (UMEZAWA & SILVEIRA, 1999). These results corroborate that ones obtained by Luciano et al.

(2009), who observed intense cross-reactions following higher differences in dog serum titers (tested by IFAT) for *Leishmania* spp. and *T. cruzi* antigens. It must be highlighted that the remaining *T. cruzi* titers varied between 40 and 640. In contrast, Souza et al. (2009) mentioned the difficulties in discriminating between infections by *T. cruzi* and *Leishmania* spp. in asymptomatic dogs using conventional diagnostic techniques. Several South American locations are endemic to CVL and CD, with a high possibility of mixed infections. Moraes et al. (2013) have also observed the simultaneous infection of canines by *L. chagasi* and *T. cruzi* in Araguaína, TO; of the 111 samples tested 57 were observed to be positive (IFAT and ELISA) for CVL (51.35%). The same sera were also analyzed by TESA-blot, which suggested the possibility of CD infection in five animals (4.5%); among these, 3 tested ELISA-positive and IFAT-negative for leishmaniasis.

In conclusion, seropositive animals for *Leishmania* spp. and *T. cruzi* were detected in the canine population in the State of Paraíba, suggesting the need for revisiting and intensification of disease control measures through constant monitoring by the competent authorities. Based on the identified risk factors for CVL and CD, we propose the observation of certain measures when allowing dogs on the streets; in addition, the environmental and housing conditions that the dogs are subjected must be improved. Contact with bovine was identified as risk factor for CD, which emphasizes the need for future studies on the role of this species in the transmission cycle of the disease.

References

- Abreu-Silva AL, Lima TB, Macedo AA, Moraes-Júnior FJ, Dias EL, Batista ZS, et al. Soroprevalência, aspectos clínicos e bioquímicos da infecção por *Leishmania* em cães naturalmente infectados e fauna de flebotomíneos em uma área endêmica na ilha de São Luís, Maranhão, Brasil. *Rev Bras Parasitol Vet* 2008;17(Suppl 1): 197-203. PMID:20059848.
- Almeida ABPF, Sousa VRF, Cruz FACS, Dahroug MAA, Figueiredo FB, Madeira MF. Canine visceral leishmaniasis: seroprevalence and risk factors in Cuiabá, Mato Grosso, Brazil. *Rev Bras Parasitol Vet* 2012; 21(4): 359-365. <http://dx.doi.org/10.1590/S1984-29612012005000005>. PMID:23184322.
- Alves AL, Bevilacqua CML, Moraes NB, Franco SO. Levantamento epidemiológico da leishmaniose visceral em cães vadios da cidade de Fortaleza, Ceará. *Ciênc Anim* 1998; 8(2): 63-68.
- Amóra SSA, Santos MJP, Alves ND, Costa SCG, Calabrese KS, Monteiro AJ, et al. Fatores relacionados com a positividade de cães para leishmaniose visceral em área endêmica do Estado do Rio Grande do Norte, Brasil. *Ciênc Rural* 2006; 36(6): 1854-1859. <http://dx.doi.org/10.1590/S0103-84782006000600029>.
- Argolo AM, Felix M, Pacheco R, Costa J. *Doença de Chagas e seus principais vetores no Brasil*. Rio de Janeiro: Imperial Novo Milênio/Fundação Oswaldo Cruz; 2008.
- Azevedo MA, Dias AKK, Paula HB, Perri SHV, Nunes CM. Avaliação da leishmaniose visceral canina em Poxoréo, Estado do Mato Grosso, Brasil. *Rev Bras Parasitol Vet* 2008; 17(3): 123-127. PMID:19245756.
- Baneth G. Leishmaniasis. In: Greene CE. *Infectious diseases of the dog and cat*. 3th ed. Saint Louis: Saunders Elsevier; 2006. p. 685-698.

- Barboza DCPM, Leal DC, Souza BMPS, Carneiro AJB, Gomes Neto CMB, Alcântara AC, et al. Inquérito epidemiológico da leishmaniose visceral canina em três distritos sanitários do Município de Salvador, Bahia, Brasil. *Rev Bras Saúde Prod Anim* 2009; 10(2): 434-447.
- Barr SC. American trypanosomiasis. In: Greene CE. *Infectious diseases of the dog and cat*. 3th ed. Philadelphia: WB Saunders; 2006. p. 676-680.
- Borchhardt DM, Mascarello A, Chiaradia LD, Nunes RJ, Oliva G, Yunes RA, et al. Biochemical evaluation of a series of synthetic chalcone and hydrazide derivatives as novel inhibitors of cruzain from *Trypanosoma cruzi*. *J Braz Chem Soc* 2010; 21(1): 142-150. <http://dx.doi.org/10.1590/S0103-50532010000100021>.
- Brasil. Ministério da Saúde, Secretaria de Vigilância em Saúde, Departamento de Vigilância Epidemiológica. *Manual de vigilância e controle da Leishmaniose Visceral* [online]. Brasília; 2006. Série A: Normas e manuais Técnicos [cited 2006 Oct 30]. Available from: http://bvmsms.saude.gov.br/bvs/publicacoes/manual_vigilancia_controle_leishmaniose_visceral.pdf
- Brasil. Ministério da Saúde, Secretaria de Vigilância em Saúde, Departamento de Vigilância Epidemiológica. *Casos confirmados de Leishmaniose Visceral, Brasil, grandes regiões e unidades federadas 1990 a 2013* [online]. Brasília; 2014 [cited 2014 Mar 27]. Available from: <http://portalsaude.saude.gov.br/index.php/o-ministerio/principal/leia-mais-o-ministerio/726-secretaria-svs/vigilancia-de-a-a-z/leishmaniose-visceral-lv/11334-situacao-epidemiologica-dados>
- Camargo ME. Fluorescent antibody test for the sorodiagnosis of American trypanosomiasis: technical modification employing preserved culture forms of *Trypanosoma cruzi* in a slide test. *Rev Inst Med Trop* 1966; 8(5): 227-235. PMID:4967348.
- Castro AG. *Controle, diagnóstico e tratamento da leishmaniose visceral (calazar): normas técnicas*. Brasília: Fundação Nacional de Saúde; 1996.
- Dias JCP, Coura JR. Epidemiologia. In: Dias JCP, Coura JR. *Clínica e terapêutica da Doença de Chagas: uma abordagem prática para o clínico geral*. Rio de Janeiro: Fiocruz; 1997. p. 33-65.
- Dias JCP, Machado EMM, Fernandes AL, Vinhaes MC. Esboço geral e perspectivas da Doença de Chagas no Nordeste do Brasil. *Cad Saude Publica* 2000; 16(Suppl 2): 13-34. <http://dx.doi.org/10.1590/S0102-311X2000000800003>. PMID:11119317.
- Dias JCP. Os primórdios do controle da doença de Chagas (em homenagem a Emmanuel Dias, pioneiro do controle, no centenário de seu nascimento). *Rev Soc Bras Med Trop* 2011; 44(Suppl 2): 12-18. <http://dx.doi.org/10.1590/S0037-86822011000800003>. PMID:21584352.
- Dohoo IR, Ducrot C, Fourichon C, Donald A, Hurnik D. An overview of techniques for dealing with large numbers of independent variables in epidemiologic studies. *Prev Vet Med* 1997; 29(3): 221-239. [http://dx.doi.org/10.1016/S0167-5877\(96\)01074-4](http://dx.doi.org/10.1016/S0167-5877(96)01074-4). PMID:9234406.
- Gürtler RE, Cecere MC, Lauricella MA, Cardinal MV, Kitron U, Cohen JE. Domestic dogs and cats as sources of *Trypanosoma cruzi* infection in rural northwestern Argentina. *Parasitology* 2007; 134(1): 69-82. <http://dx.doi.org/10.1017/S0031182006001259>. PMID:17032467.
- Hosmer DW, Lemeshow S. *Applied logistic regression*. New York: John Wiley & Sons; 2000.. <http://dx.doi.org/10.1002/0471722146>.
- Instituto Brasileiro de Geografia e Estatística – IBGE. *Estimativas da população residente nos municípios brasileiros com data de referência em 1º de julho de 2013* [online]. Brasília: IBGE; 2013 [cited 2015 Oct 10]. Available from: ftp://ftp.ibge.gov.br/Estimativas_de_Populacao/Estimativas_2013/nota_metodologica_2013.pdf
- Kuhls K, Alam MZ, Cupolillo E, Ferreira GEM, Mauricio IL, Oddone R, et al. Comparative microsatellite typing of new world *Leishmania infantum* reveals low heterogeneity among populations and its recent old world origin. *PLoS Negl Trop Dis* 2011; 5(6): e1155. <http://dx.doi.org/10.1371/journal.pntd.0001155>. PMID:21666787.
- Luciano RM, Lucheis SB, Troncarelli MZ, Luciano DM, Langoni H. Avaliação da reatividade cruzada entre antígenos de *Leishmania* spp e *Trypanosoma cruzi* na resposta sorológica de cães pela técnica de imunofluorescência indireta (RIFI). *Braz J Vet Res Anim Sci* 2009; 46(3): 181-187.
- Martins IV. *Aspectos epidemiológicos e de hemostasia na leishmaniose visceral canina*. [Dissertation]. Recife: Universidade Federal Rural de Pernambuco; 2008.
- Matos MM, Filgueira KD, Amora SSA, Suassuna ACD, Ahid SMM, Alves ND. Ocorrência da Leishmaniose Visceral em cães em Mossoró, Rio Grande do Norte. *Ciênc Anim* 2006; 16(1): 51-54.
- Melo MN. Leishmaniose Visceral no Brasil: desafios e perspectivas. *Rev Bras Parasitol Vet* 2004; 13(S1): 41-45.
- Mendes RS, Santana VL, Jansen AM, Xavier SCC, Vidal IF, Rotondano TEF, et al. Aspectos epidemiológicos da Doença de Chagas canina no semiárido paraibano. *Pesq Vet Bras* 2013; 33(12): 1459-1465. <http://dx.doi.org/10.1590/S0100-736X2013001200011>.
- Moraes AN, Sousa MG, Meireles LR, Kesper N Jr, Umezawa ES. Canine visceral leishmaniasis and Chagas disease among dogs in Araguaína, Tocantins. *Rev Bras Parasitol Vet* 2013; 22(2): 225-229. <http://dx.doi.org/10.1590/S1984-29612013005000024>. PMID:23802237.
- Naveda LAB, Moreira EC, Machado JG, Moraes JRC, Marcelino AP. Aspectos epidemiológicos da leishmaniose visceral canina no município de Pedro Leopoldo, Minas Gerais, 2003. *Arq Bras Med Vet Zootec* 2006; 58(6): 988-993. <http://dx.doi.org/10.1590/S0102-09352006000600003>.
- Oliveira SS, Araújo TM. Avaliação das ações de controle da leishmaniose visceral (calazar) em uma área endêmica do estado da Bahia, Brasil (1995-2000). *Cad Saude Publica* 2003; 19(6): 1681-1690. <http://dx.doi.org/10.1590/S0102-311X2003000600012>. PMID:14999334.
- Reichmann MLAB, Pinto HBF, Nunes VP. *Vacinação contra raiva de cães e gatos* [online]. São Paulo; 1999 [cited 2015 Oct 10]. Available from: http://www.saude.sp.gov.br/recursos/instituto-pasteur/pdf/manuais/manual_03.pdf
- Rondon FCM, Bevilacqua CML, Franke CR, Barros RS, Oliveira FR, Alcântara AC, et al. Cross-sectional serological study of canine *Leishmania* infection in Fortaleza, Ceará state, Brazil. *Vet Parasitol* 2008; 155(1-2): 24-31. <http://dx.doi.org/10.1016/j.vetpar.2008.04.014>. PMID:18565676.
- Santa Rosa ICA, Oliveira ICS. Leishmaniose Visceral: zoonose reemergente. *Clin Vet* 1997; 2(11): 24-28.
- Silva MRM, Nascimento RCS, Negrão AMG, Valente VC, Valente SAS. *Perfis sorológicos e hematológicos de cães em área de ocorrência de doença de chagas aguda no município de Bragança, estado do Pará* [online]. Santa Catarina: Anclivepa; 2014 [cited 2014 May 04]. Available from: www.anclivepa2014.com.br/353/135.pdf
- Silva PB, Fernandes JI. *Aspectos clínicos epidemiológicos da infecção por Trypanosoma cruzi em cães naturalmente infectados no município de São Domingos Do Capim – Pará* [online]. Pará: UFPA; 2013 [cited 2013 Sept 17]. Available from: http://www.pibic.ufpa.br/ANAISSEMINIC/XXIVSEMINIC/arquivos/resumos/Ciencias_Agrarias/ciencias_agrarias_014.pdf
- Simões-Mattos L, Mattos MRF, Teixeira MJ, Oliveira-Lima JW, Bevilacqua CML, Prata-Júnior RC, et al. The susceptibility of domestic cats (*Felis*

catus) to experimental infection with *Leishmania braziliensis*. *Vet Parasitol* 2005; 127(3-4): 199-208. <http://dx.doi.org/10.1016/j.vetpar.2004.10.008>. PMID:15710520.

Souza AI, Oliveira TMFS, Machado RZ, Camacho AA. Soroprevalência da infecção por *Trypanosoma cruzi* em cães de uma área rural do Estado de Mato Grosso do Sul. *Pesqui Vet Bras* 2009; 29(2): 150-152. <http://dx.doi.org/10.1590/S0100-736X2009000200011>.

Thrusfield M. *Veterinary epidemiology*. 3rd ed. Oxford: Blackwell Science; 2007.

Uchôa CMA, Serra CMB, Duarte R, Magalhães CM, Silva RM, Theophilo F, et al. Aspectos sorológicos e epidemiológicos da leishmaniose tegumentar americana canina em Maricá, Rio de Janeiro, Brasil. *Rev Soc*

Bras Med Trop 2001; 34(6): 563-568. <http://dx.doi.org/10.1590/S0037-86822001000600011>. PMID:11813064.

Umezawa ES, Silveira JF. Serological diagnosis of Chagas disease with purified and defined *Trypanosoma cruzi* antigens. *Mem Inst Oswaldo Cruz* 1999;94(S1 Suppl Suppl 1): 285-288. <http://dx.doi.org/10.1590/S0074-02761999000700051>. PMID:10677737.

World Health Organization – WHO. *Guidelines for dog population management*. Geneva: WHO; 1990.

World Health Organization – WHO. *Chagas disease (American trypanosomiasis)* [online], Geneva: WHO; 2015 [cited 2015 Mar 01]. Available from: <http://www.who.int/mediacentre/factsheets/fs340/en/>

Zar JH. *Biostatistical analysis*. 4th ed. Upper Saddle River: Prentice Hall; 1999.