



Revista Brasileira de Cirurgia
Cardiovascular/Brazilian Journal of
Cardiovascular Surgery

ISSN: 0102-7638

revista@sbccv.org.br

Sociedade Brasileira de Cirurgia
Cardiovascular

Aygün, Fatih; Özülkü, Mehmet; Günday, Murat
Effects of Cardiopulmonary Bypass on Mediastinal Drainage and the Use of Blood
Products in the Intensive Care Unit in 60- to 80-Year-Old Patients Who Have Undergone
Coronary Artery Bypass Grafting
Revista Brasileira de Cirurgia Cardiovascular/Brazilian Journal of Cardiovascular Surgery,
vol. 30, núm. 6, 2015, pp. 597-604
Sociedade Brasileira de Cirurgia Cardiovascular
São José do Rio Preto, Brasil

Available in: <http://www.redalyc.org/articulo.oa?id=398944486002>

- How to cite
- Complete issue
- More information about this article
- Journal's homepage in redalyc.org

redalyc.org

Scientific Information System

Network of Scientific Journals from Latin America, the Caribbean, Spain and Portugal

Non-profit academic project, developed under the open access initiative

Effects of Cardiopulmonary Bypass on Mediastinal Drainage and the Use of Blood Products in the Intensive Care Unit in 60- to 80-Year-Old Patients Who Have Undergone Coronary Artery Bypass Grafting

Fatih Aygün¹, MD; Mehmet Özülkü¹, MD; Murat Günday¹, MD



DOI: 10.5935/1678-9741.20150086

Abstract

Objective: The present study consisted of patients who underwent on-pump coronary artery bypass grafting (CABG) and off-pump CABG and investigated effect of using cardiopulmonary bypass (CPB) on the amount of postoperative drainage and blood products, red blood cell (RBC), free frozen plasma (FFP) given in the intensive care unit in 60-80-year-old patients who underwent CABG.

Methods: The present study comprises a total of 174 patients who have undergone coronary artery bypass graft (off-pump or on-pump CABG) surgery in our clinic in between 2012-2015 year.

Results: It was observed that the amount of drainage in the first 24 postoperative hours was lower in the on-pump CABG group (Group 1) when compared to off-pump group (Group 2) (Group 1 vs. Group 2; 703.5±253.8 ml vs. 719.6±209.4 ml; $P=0.716$). However, the amount of drainage in the second 24 hours was

statistically significantly lower in the off-pump CABG group (Group 1 vs. Group 2; 259.8±170.6 ml vs. 190.1±129.1 ml; $P=0.016$). With regard to the amount of overall drainage, no statistically significant difference was observed between the two groups. Group 1 needed RBC transfusion higher than Group 2 (Group 1 vs. Group 2; 2.2±1.3 bag vs. 1.2±0.9 bag; $P<0.001$).

Conclusion: We can say that CPB influences the amount of second 24-hour drainage which indexed body surface area. In addition, CPB decreases hct, hb, thrombocyte count in ICU arrived, after 24 hours in postoperative period. Reduced thrombocyte counting effect can be appeared after 48 hours in the postoperative period of CPB.

Keywords: Coronary Artery Bypass. Cardiopulmonary Bypass. Coronary Artery Bypass, Off-Pump. Health of the Elderly. Drainage. Blood Transfusion.

Abbreviations, acronyms & symbols

ACT	= Activated clotting time
BMI	= Body mass index
CABG	= Coronary artery bypass grafting
COPD	= Chronic obstructive pulmonary disease
CPB	= Cardiopulmonary bypass
CVS	= Cardiovascular surgery
FFP	= Fresh frozen plasma
Hb	= Hemoglobin
Hct	= Hematocrit
ICU	= Intensive care unit
LIMA	= Left internal mammary artery
MCPB	= Mini cardiopulmonary bypass
PAD	= Peripheral arterial disease
RBC	= Red blood cell
SPSS	= Statistical Package for the Social Sciences

INTRODUCTION

Coronary artery bypass graft (CABG) surgery is one of the most common surgical procedures worldwide; however, it can also lead to numerous complications. The conventional CABG procedure is performed with the use of a cardiopulmonary bypass (CPB) device or heart-lung machine and is known as "on-pump" CABG, whereas coronary bypass surgery performed without the use of CPB is known as "off-pump" CABG. CABG surgery has a greater level of risk when performed in elderly patients^[1]. Hence, cardiac surgeons have to consider the disadvantages of using CPB in older patients, particularly with regard to increased drainage and other coincident effects. Ferraris & Ferraris^[2] identified a relationship between the greater use of blood and its products after CABG and postoperative infection, because excessive postoperative drainage can lead to infection and other complications resulting in an increased need for blood transfusions.

¹Başkent University, Konya Research and Medical Center, Turkey.

This study was carried out at Department of Cardiovascular Surgery, Konya Medical and Research Center, Başkent University, Konya, Turkey.

No financial support.

Correspondence Address:

Fatih Aygün

Department of Cardiovascular Surgery, Konya Medical and Research Center, Başkent University, Konya

Hocacıhan Mah. Saray Cad No:1, 42000 Selçuklu-Konya, Turkey

E-mail: fatihaygun@ttmail.com

Article received on May 4th, 2015
Article accepted on November 25th, 2015

On-pump CABG is considered to be the gold standard, but this method has some physiological consequences, including thrombocytopenia, activation of the complement system, immune suppression and an inflammatory response that causes organ dysfunction. In addition, manipulation of the ascending aorta during cannulation (cannulation and cross-clamping) poses the ensuing risk of embolization and stroke^[3]. Thus, drainage tubes are inserted into the mediastinum and thoracic cavity at the end of the CABG procedure to monitor leakage and bleeding throughout these regions. A large volume of drained fluid implies the need for increased use of blood products (erythrocyte suspension and fresh frozen plasma [FFP]), and can lead to unfavorable consequences.

The risk factors for transfusion after CABG include advancing age, preoperative aspirin therapy, priority of the operation, lower preoperative volume of red blood cells (RBCs), duration of the CPB, recent fibrinolytic therapy, reoperative CABG, and differences in heparin management^[4-9]. Off-pump CABG may reduce the need for the transfusion of blood products, lead to shorter stays in the intensive care unit (ICU), and reduce the postoperative hospitalization time^[10-12]. However, in one previous study, there was no difference in the level of bleeding, mediastinitis, and rate of mortality between off-pump and on-pump CABG patients^[13].

We investigated the effects of CPB on the level of postoperative drainage and the number of blood products administered in the ICU to patients between the ages of 60 and 80 years who underwent both on-pump and off-pump CABG.

METHODS

Clinical characteristics of the patients

A total of 174 individuals who underwent CABG surgery at our clinic between 2012 and 2015 were included in this study. Data were collected retrospectively, and approval for the study was obtained by the ethics committee of our institution. The patients were reviewed consecutively and included in the study according to predetermined criteria. Care was taken to ensure that the make-up of each group was similar.

All of the patients had their medical history taken and underwent a detailed physical examination. In the preoperative period, standard preoperative laboratory tests including a pulmonary function test (Spirobank spirometer; MIR, Waukesha, WI, USA), transthoracic echocardiography (Sequoia C256; Acuson, Mountain View, CA, USA), and bilateral carotid artery Doppler ultrasonography (Xario prime ultrasound; Toshiba, Japan) were performed in the cardiovascular surgery (CVS) clinic. Calcifications in the thoracic and ascending aorta and aortic arch were evaluated by chest radiography prior to surgery. During surgery, the starting points of the ascending aorta and aortic arch were thoroughly examined through manipulation. The procedure was changed if plaque was detected, and these patients were not included in the study.

During the preoperative period, clopidogrel (if any was being administered) was discontinued 5 days prior to surgery, and acetylsalicylic acid (if any) was discontinued 1 day prior to surgery in patients scheduled to undergo on-pump (cross-clamped) CABG.

The same was done for those scheduled for off-pump (beating-heart) CABG. The blood glucose levels of patients with type 2 diabetes were regulated with insulin both before

and after surgery, and were maintained below 200 mg/dL. Preoperative blood samples were collected in standard ethylenediaminetetraacetic acid vacuum tubes, and analyzed using a Cell-Dyn 3700 instrument (Abbott Laboratories, Abbott Park, IL, USA). Prothrombin time, activated partial thromboplastin time, and international normalized ratio levels were measured. The preoperative data of all study participants were within the normal ranges.

Dyslipidemia was defined as a fasting total cholesterol level in serum ≥ 240 mg/dL, triglyceride level ≥ 200 mg/dL, low-density lipoprotein cholesterol level ≥ 160 mg/dL, and/or high-density lipoprotein cholesterol level < 40 mg/dL; anyone receiving active medical therapy for this condition was also considered to have it^[14]. Cholesterol levels were measured using an enzymatic method.

Height and body weight were measured (SECA, Vogel & Halke, Hamburg, Germany) before surgery, and body mass index (BMI) was calculated. A staged approach was adopted in patients with a carotid artery stenosis of 70-100%, and was reserved for the post-CABG period. Preoperative data, including age, gender, incidence of chronic obstructive pulmonary disease (COPD), and peripheral arterial disease rate are provided in Table 1.

Surgical procedure

First, isolated CABG surgery was performed in all of the study participants by the same surgical team. Fentanyl, midazolam, and pancuronium bromide were administered for the induction of anesthesia. Then, standard median sternotomy was performed. The left internal mammary artery (LIMA) and other vascular conduits (including the saphenous vein and radial artery) were prepared. Heparin sodium (Nevparin® flacon 25000 IU/5 mL) was administered at a dose of 300 IU/kg. CPB and a cross clamp, as well as standard aortic and two-stage venous cannulas, were used. A Jostra-Cobe heart-lung machine (model 043213 105; VLC 865, Sweden) was used. All of the patients received crystalloid cardioplegy during surgery and hot-shot cardioplegy at the end of surgery. While the LIMA was used in all of the cases, the right internal mammary artery was not used. The great saphenous vein and radial artery were used as grafts. A scrupulous aseptic regime was implemented in all of the surgeries. The unnecessary use of electrocautery, as well as luxury perfusion in CPB, was avoided. Sodium heparin was administered at a dose of 150 IU/kg in patients who underwent the beating-heart technique. Distal anastomoses were performed using the Octopus tissue stabilizer and Starfish heart positioner (Medtronic, Dublin, Ireland). Hematocrit (Hct) and hemoglobin (Hb) values were monitored every 20 min, beginning from the induction of anesthesia until the end of surgery. Intraoperative blood transfusion was performed when the Hct value decreased to 20%. These cases underwent full vascularization during the CABG procedure. Mediastinal and chest drainage tubes were situated subxiphoidally.

Proximal anastomoses to the aorta were performed with the assistance of a side clamp in both the on-pump and off-pump techniques. At the end of surgery, patients who underwent on-pump CABG received protamine hydrochloride (Protamin® ampoule 1000 IU/1 mL) at appropriate doses for full-dose neutralization, and the activated clotting time (ACT) was maintained between 100 and 120. Patients who underwent off-pump CABG received appropriate doses of protamine

Table 1. Preoperative data according to groups.

	Group 1 (n=91) (On-pump CABG)	Group 2 (n=83) (Off-pump CABG)	P values
Age (±SD) (year)	68±5.2	67±4.6	0.437 ^T
Gender (male)	53 (58.2%)	58 (69.9%)	0.118 ^P
Smoking	34 (37.4%)	37 (44.6%)	0.357 ^P
COPD	16 (17.6%)	21 (25.3%)	0.266 ^P
Hypertension	82 (90.1%)	65 (78.3%)	0.037 ^P
PAD	5 (5.5%)	1 (1.2%)	0.214 ^F
Preoperative stroke story	10 (11%)	5 (6%)	0.288 ^P
Diabetes	*59 (64.8%)	*49 (59%)	*0.439 ^P
no-diabetes			
oral a/d	22 (24.2%)	26 (31.3%)	
parenteral a/d	10 (11%)	8 (9.6%)	
Right carotid artery			
stenosis<50%	55 (60.4%)	29 (34.9%)	*0.641 ^P
50%< stenosis≤70%	*5 (5.5%)	*5 (6%)	
70%≤ stenosis<100%	*2 (2.2%)	—	
stenosis=100%	—	—	
Left carotid artery			
stenosis<50%	48 (52.7%)	30 (36.1%)	*0.003 ^F
50%< stenosis≤70%	*11 (12.1%)	*1 (1.2%)	
70%≤ stenosis<100%	—	—	
stenosis=100%	*1 (1.1%)	—	
BMI	29.9±5.4	29.9±5.1	0.983 ^T
Ejection Fraction	52.3±7.5	54.6±8.4	0.070 ^T

^T=P value was presented as a result Student-t test

^P=P value was presented as a result Pearson chi-square test

^F=P value was presented as a result Fischer's Exact tests

*=P value was calculated according to carotid artery stenosis <50%

CABG=coronary artery bypass grafting; BMI=body mass index; SD=standard deviation. PAD=peripheral artery disease;

COPD =chronic obstructive pulmonary disease

Table 2. Operative data according to groups.

	Group 1 (n=91) (On-pump CABG)	Group 2 (n=83) (Off-pump CABG)	P values
Saphenous graft	91 (100%)	73 (88%)	-
Radial artery graft	56 (61.5%)	16 (19.3%)	<0.001 ^P
LIMA graft	89 (97.8%)	83 (100%)	-
Numbers of grafting	3.7±0.7	2.5±0.9	<0.001 ^T
RBC (bag) (while in the operatin groom)	0.5±0.6	0.4±0.6	<0.001 ^T

^T=P value was presented as a result Student-t test

^P=P value was presented as a result Pearson chi-square test

CABG=coronary artery bypass grafting; LIMA=left internal mammary artery; RBC=red blood cell

hydrochloride, as before, to maintain the ACT between 140 and 150. Data related to the surgery, including choice of graft and graft number, are listed in Table 2. Table 3 provides the postoperative Hb and Hct values, thrombocyte count, and drainage on arrival in the CVS ICU, and at the first and second 24 hours periods.

Postoperative care

After the surgery was completed, the patients were admitted to the CVS ICU, and their Hct and Hb values were monitored

at 4 hours intervals. The criterion for blood transfusion was an Hct value of 28%. During the postoperative period, 300 mg/day acetylsalicylic acid (Coraspin 300®) was commenced in all of the patients along with enteral nutrition to reduce the risk of post-CABG stroke. Cefazolin sodium (Cefamezin®-IM/IV), which is used as the standard antibiotic in our clinic, was initially administered at a dose of 1 g, 30 minutes before surgery, and then every 8 hours after surgery, and was continued for a total of 72 hours. Blood glucose regulation in diabetic patients was strictly maintained after surgery using 100 IU/

Table 3. Postoperative data according to groups.

	Group 1 (n=91) (On-pump CABG)	Group 2 (n=83) (Off-pump CABG)	P values
Preoperative Hct	39.8±3.7	38.5±5.1	0.067 ^T
Preoperative Hb	12.3±1.6	12.5±1.6	0.322 ^T
Preoperative trombocyst count	259.4±80.6	260.9±77.1	0.899 ^T
Hct (arrived ICU)	30.1±1.9	32±4.4	0.001 ^T
Hb (arrived ICU)	10±0.8	10.9±2	0.001 ^T
Trombocyst (arived ICU)	164.2±58.2	241.4±91.9	<0.001 ^T
Hct (24 th hour in the ICU)	30.2±1.5	31.4±3.0	0.003 ^T
Hb (24 th hour in the ICU)	10±0.5	10.4±1.0	0.005 ^T
Trombocyst (24 th hour in the ICU)	187.9±74.7	228.8±78.2	0.001 ^T
Hct (48 th hour in the ICU)	31.2±1.9	31.5±2.9	0.402 ^T
Hb (48 th hour in the ICU)	10.4±0.6	10.5±1.0	0.495 ^T
Trombocyst (48 th hour in the ICU)	184.1±63.7	207.5±75.2	0.029 ^T
First 24 th hour drainage (ml)	704.3±253	713.8±209.8	0.788 ^T
Second 24 th hour drainage (ml)	275.2±176.9	188.5±132.3	<0.001 ^T
Total drainage count (ml)	979.6±374.7	902.4±299.1	0.133 ^T
First 24 th hour drainage (ml)/m ²	382.6±139.5	384.4±116.8	0.925 ^T
Second 24 th hour drainage (ml)/m ²	150.2±97.7	100.9±71.5	<0.001 ^T
Total drainage count (ml)/m ²	533.4±208.2	485.9±165.7	0.096 ^T
FFP (bag) (used in the ICU)	2.7±1.7	2.9±1.3	0.366 ^T
RBC (bag) (used in the ICU)	2.2±1.3	1.2±0.9	<0.001 ^T

^T=P value was presented as a result Student-t test

Hct=haematocryt; Hb=hemoglobin; CABG=coronary artery bypass grafting; ICU=intensive care unit; FFP=fresh frozen plasma; RBC=red blood cell

mL insulin glargine (Lantus® flacon), and 100 IU/mL human soluble insulin (Humulin-R® flacon), and insulin infusions were administered when required. The blood glucose concentration was maintained below 200 mg/dL in all of the diabetic patients.

The patients' drainage tubes were checked over a 48 hours period after the patient was admitted to the CVS ICU. After a stay in the ICU, patients were transferred to the CVS clinic with their drains and arterial catheters removed, and on postoperative day 4, the central venous line was removed. Patients were discharged from the hospital on postoperative days 6–11.

Statistical analyses

Statistical analyses were performed with Statistical Package for the Social Sciences (SPSS) software (SPSS Inc., Chicago, IL, USA). The statistical significance of nonparametric data between the groups was analyzed by a Chi-square test and Fisher's exact test, because the observed values were below those expected. While the parametric data were represented as minimum, maximum, and mean ± standard deviation, the statistical significance of the parametric data between the groups was analyzed by an independent Student's t-test. Results were considered statistically significant at $P<0.05$.

Study groups

Consecutive patients who underwent the CABG procedure between 2012 and 2015 were grouped randomly according to two different surgical techniques. The first group (Group 1) consisted of patients who underwent CABG by CPB and the cross-clamp (on-pump with cross-clamp) technique. The second

group (Group 2) consisted of patients who underwent CABG by the beating-heart (off-pump) technique. Proximal anastomosis was performed using side clamps. The duration of use of the cross-clamp did not exceed 90 minutes, and the length of CPB did not exceed 120 minutes in Group 1.

To create a homogeneous group, the exclusion criteria were as follows: hypoalbuminemia (a serum albumin level < 3.5 gr/dL); nutritional disorders; dialysis or a serum creatinine level of 2 gr/dL; aortic pathology detected during surgery and during which the operational procedure was changed; needing postoperative intra-aortic balloon pump support; and having undergone surgery under emergency conditions, reoperative CABG, CABG without a touching ascending aorta (no-touch), LIMA-left anterior descending (for single vascular disease patients) CABG, a second CABG, valvular and CABG during the same session, or surgical exploration for any other postoperative reason, including chronic renal insufficiency and dialysis.

RESULTS

In all, 91 (52.2%) patients underwent CABG by CPB, and 83 (47.8%) underwent CABG by the beating-heart technique. The age range of subjects was 61–79 years (mean ± standard deviation, 68±5 years); there were 111 (63.8%) males and 63 (36.2%) females. There were 147 (84.5%) patients with hypertension, 71 (40.8%) smokers, and 37 (21.3%) patients with COPD (Table 1). The number of grafts was 3.1±1 (Table 2). The preoperative Hct value was 39.2±4.4%, the Hb was 12.4±1.6 gr/dL, and the thrombocyte count was 260.1±78.7. At the time of admission to the CVS ICU after surgery, the

Hct value was $31 \pm 3.5\%$, the Hb was 10.4 ± 1.5 gr/dL, and the thrombocyte count was 201.1 ± 85.2 . The drainage was 708.9 ± 232.8 mL during the first postoperative 24 hours, whereas it was 233.9 ± 162.7 mL during the second 24 hours. The number of RBC collection bags administered was 1.7 ± 1.2 and the number of bags of FFP was 2.8 ± 1.5 (Table 3).

Although it was not statistically significant, the amount of drainage during the first postoperative 24 hours was lower in Group 1 than in Group 2 (704.3 ± 253 mL vs. 713.8 ± 209.8

mL, respectively; $P=0.788$). Indexing this value according to body surface area led to similar results (382.6 ± 139.5 mL vs. 384.4 ± 116.8 mL, respectively; $P=0.925$) (Figures 1 and 2). However, the amount of drainage during the second postoperative 24 hours was significantly lower in Group 1 than in Group 2 (275.2 ± 176.9 mL vs. 188.5 ± 132.3 mL, respectively; $P<0.001$). Again, indexing this value led to similar results (150.2 ± 97.7 mL vs. 100.9 ± 71.5 mL, respectively; $P<0.001$) (Figures 3 and 4).




Fig. 1 - Group 1 first 24 hours drainage.




Fig. 2 - Group 2 first 24 hours drainage.




Fig. 3 - Group 1 second 24 hours drainage.




Fig. 4 - Group 2 second 24 hours drainage.




Fig. 5 - Group 1 total drainage.




Fig. 6 - Group 2 total drainage.

With regard to the amount of overall drainage, no statistically significant differences were observed between the two groups (Figures 5 and 6). The mean number of bypass grafts used and the use of the radial artery were significantly higher in Group 1 than in Group 2. In addition, the Hct and Hb values and thrombocyte count were significantly lower in Group 1 at the time of admission to the ICU; however, the thrombocyte count subsequently recovered to the value of the off-pump group after 48 hours. In addition, there were no differences in mean Hct and Hb values between the groups at the end of 48 hours (Table 3).

DISCUSSION

The conventional CABG procedure is characterized by precisely performed coronary anastomoses using CPB. However, ensuring a bleeding-free surgical area and performing precise coronary anastomosis using CPB may lead to complications such as blood trauma, inflammatory responses, a negative non-pulsatile flow, and potential debris embolization^[3]. Off-pump CABG is considered to be a means of avoiding such unfavorable outcomes of CPB. Therefore, we included 91 (52%) on-pump CABG patients in our study.

Recently, CABG has been performed in a growing number of patients with a higher risk profile. The benefits of off-pump CABG with regard to aortic complications are obvious. Recent studies have demonstrated an improvement in high-risk patients who have undergone off-pump CABG^[15-18]. In our study, we included 83 (48%) off-pump CABG patients.

Off-pump and the gold standard on-pump CABG have been compared in both large retrospective observational studies and randomized controlled studies. The results of these studies have revealed comparable outcomes. However, there is a lack

of small, prospective, randomized and controlled studies in the literature. Some reports have indicated that such studies are unable to demonstrate the early and long-term outcomes related to incomplete revascularization, a decreased rate of long-term graft patency, enhanced repetitive revascularization, and survival. This has discouraged research on off-pump CABG and led to the abeyance of this technique. Other reports have claimed that studies that doubt the applicability and benefits of off-pump CABG have ignored research indicating comparable long-term outcomes and a greatly reduced period of hospitalization with the on-pump technique^[19-23].

Van Dijk et al.^[24], Angelini et al.^[25], and Cooley^[26] identified a greater amount of drainage following on-pump CABG as opposed to off-pump CABG. Abdel Aal et al.^[27] identified hemodilution as the reason for the difference between the two techniques. In our study, we did not find a statistically significant difference in either the first postoperative 24 hours drainage volume or the total drainage between the two groups. The second postoperative 24 hours drainage was significantly higher in Group 1 ($P<0.001$). When we indexed the drainage according to the body surface area, we achieved similar results.

Beghi et al.^[28] and Hall et al.^[29] emphasized that, besides hemodilution, blood and air contact, hypothermia, and use of the CPB device also contribute to this difference. In our study, the Hct and Hb values, and thrombocyte count upon arrival in the CVS ICU and also after the first postoperative 24 hours, were significantly lower in Group 1 ($P=0.01$; $P=0.01$; $P<0.01$, respectively, on ICU arrival, and $P=0.003$; $P=0.005$; $P=0.001$, respectively, after 24 hours). After 48 hours, only the thrombocyte count was significantly lower ($P=0.029$).

Tsai et al.^[30] reported an almost similar amount of drainage in their on-pump and off-pump CABG groups, with a slightly

lower volume observed in the off-pump CABG group. The authors found that a shortened CPB time and elimination of hypothermia reduced the amount of postoperative drainage during the on-pump CABG procedure. In our study, the length of time that the cross-clamp was used did not exceed 90 minutes, and the duration of CPB did not exceed 120 minutes, for patients in Group 1.

Panday et al.^[31] determined that the amounts of blood and blood products used in cases of off-pump CABG were slightly lower than in cases of CABG performed by the on-pump technique. In addition, the amounts used preoperatively were almost equal between a group that underwent mini cardiopulmonary bypass (MCPB) and an on-pump CPB group. We did not find a statistical significance in terms of the postoperative use of FFP between the two groups ($P=0.036$); however, RBC usage in the ICU following CABG surgery was significantly higher in Group 1 ($P<0.001$).

Ranucci & Isgrò^[32] reported that MCPB implementations are characterized by decreased postoperative bleeding, but not by a low transfusion rate. Folliguet et al.^[33] compared the postoperative transfusion rates of patients who underwent MCPB to conventional CPB, and found no statistically significant differences between the two groups. The researchers also noted the importance of the roles of anesthesiologists and pump technicians during surgery.

Study Limitations

In the present study, all study participants were Caucasians and do not represent the other ethnic groups. Patients that would impair the similarity between the groups, such as patients with renal insufficiency and dialysis patients and redo CABG cases etc., have not been included in the study.

CONCLUSION

The use of CPB increases the level of drainage during the second 24 hours postoperative period (Group 1 vs. Group 2; 275.2 ± 176.9 mL vs. 188.5 ± 132.3 mL; $P<0.001$), and also increases the second 24 hours drainage volume if indexed by the body surface area (Group 1 vs. Group 2; 150.2 ± 97.7 mL/m² vs. 100.9 ± 71.5 mL/m²; $P<0.001$) in 60- to 80-year-old patients who undergo the CABG procedure. We found that CPB decreased Hct and Hb values and thrombocyte counts on arrival at the ICU and after 24 postoperative hours. In addition, CPB increased RBCs usage during the intraoperative (Group 1 vs. Group 2; 0.5 ± 0.6 bag vs. 0.4 ± 0.6 bag; $P<0.001$) and postoperative periods (Group 1 vs. Group 2; 2.2 ± 1.3 bag vs. 1.2 ± 0.9 bag; $P<0.001$). These data should be verified by large studies in the future.

Authors' roles & responsibilities

FA	Execution of surgeries and/or experiments; analysis and/or interpretation of data; statistical analysis; writing of the manuscript or critical review of its contents; final approval of the manuscript
MÖ	Design and drawing of the study; final approval of the manuscript
MG	Final approval of the manuscript

REFERENCES

1. Iglezias JCR, Dallan LA, Oliveira SF, Ramires JA, Oliveira SA, Verginelli G, et al. Aorto-coronary bypass in elderly. *Rev Bras Cir Cardiovasc*. 1990;5(3):183-6.
2. Ferraris VA, Ferraris SP. Limiting excessive postoperative blood transfusion after cardiac procedures: a review. *Tex Heart Inst J*. 1995;22(3):216-30.
3. Raja SG, Husain M, Popescu FL, Chudasama D, Daley S, Amrani M. Does off-pump coronary artery bypass grafting negatively impact long-term survival and freedom from reintervention? *Biomed Res Int*. 2013;2013:602871.
4. Ferraris VA, Gildengorin V. Predictors of excessive blood use after coronary artery bypass grafting: a multivariate analysis. *J Thorac Cardiovasc Surg*. 1989;98(4):492-7.
5. Scott WJ, Kessler R, Wernly JA. Blood conservation in cardiac surgery. *Ann Thorac Surg*. 1990;50(5):843-51.
6. Cosgrove DM, Loop FD, Lytle BW, Gill CC, Golding LR, Taylor PC, et al. Determinants of blood utilization during myocardial revascularization. *Ann Thorac Surg*. 1985;40(4):380-4.
7. Hardy JF, Perrault J, Tremblay N, Robitaille D, Blain R, Carrier M. The stratification of cardiac surgical procedures according to use of blood products: a retrospective analysis of 1480 cases. *Can J Anaesth*. 1991;38(4 pt 1):511-7.
8. Bracey AW, Radovancevi R. The hematologic effects of cardiopulmonary bypass and the use of hemotherapy in coronary artery bypass grafting. *Arch Pathol Lab Med*. 1994;118(4):411-6.
9. Goodnough LT, Johnston MF, Toy PT. The variability of transfusion practice in coronary artery bypass surgery. *Transfusion Medicine Academic Award Group. JAMA*. 1991;265(1):86-90.
10. Puskas JD, Thourani VH, Marshall JJ, Dempsey SJ, Steiner MA, Sammons BH, et al. Clinical outcomes, angiographic patency, and resource utilization in 200 consecutive off-pump coronary bypass patients. *Ann Thorac Surg*. 2001;71(5):1477-83.
11. Calafiore AM, Di Mauro M, Contini M, Di Giammarco G, Pano M, Vitolla G, et al. Myocardial revascularization with and without cardiopulmonary bypass in multivessel disease: impact of the strategy on early outcome. *Ann Thorac Surg*. 2001;72(2):456-62.
12. Sabik JF, Gillinov AM, Blackstone EH, Vacha C, Houghtaling PL, Navia J, et al. Does off-pump coronary surgery reduce morbidity and mortality? *J Thorac Cardiovasc Surg*. 2002;124(4):698-707.
13. Hernandez F, Cohn WE, Baribeau YR, Tryzelaar JF, Charlesworth DC, Clough RA, et al., Northern New England Cardiovascular Disease Study Group. In-hospital outcomes of off-pump versus on-pump coronary artery bypass procedures: a multicenter experience. *Ann Thorac Surg*. 2001;72(5):1528-33.
14. Reiner Z, Catapano AL, De Backer G, Graham I, Taskinen MR, Wiklund O, et al.; ESC Committee for Practice Guidelines (CPG) 2008-2010 and 2010-2012 Committees. ESC/EAS Guidelines for the management of dyslipidaemias: the Task Force for the management of dyslipidaemias of the European Society of Cardiology (ESC) and the European Atherosclerosis Society (EAS). *Eur Heart J*. 2011;32(14):1769-818.
15. Lemma MG, Coscioni E, Tritto FP, Centofanti P, Fondacone C, Salica A, et al. On-pump versus off-pump coronary artery bypass surgery in high-risk patients: operative results of a prospective randomized trial (on-off study). *J Thorac Cardiovasc Surg*. 2012;143(6):625-31.
16. Barandon L, Richebé P, Munos E, Calderon J, Lafitte M, Lafitte S, et al. Off-pump coronary artery bypass surgery in very high-risk patients: adjustment and preliminary results. *Interact Cardiovasc Thorac Surg*. 2008;7(5):789-93.
17. Marui A, Okabayashi H, Komiya T, Tanaka S, Furukawa Y, Kita T, et al.; CREDO-Kyoto Investigators. Benefits of off-pump coronary artery bypass grafting in high-risk patients. *Circulation*. 2012;126(11 suppl. 1):S151-7.
18. Mack MJ, Pfister A, Bachand D, Emery R, Magee MJ, Connolly M, et al. Comparison of coronary bypass surgery with and without cardiopulmonary bypass in patients with multivessel disease. *J Thorac Cardiovasc Surg*. 2004;127(1):167-73.

19. Zangrillo A, Crescenzi G, Landoni G, Leoni A, Marino G, Calabrò MG, et al. Off-pump coronary artery bypass grafting reduces postoperative neurologic complications. *J Cardiothorac Vasc Anesth*. 2005;19(2):193-6.
20. Farrokhhyar F, Wang X, Kent R, Lamy A. Early mortality from off-pump and on-pump coronary bypass surgery in Canada: a comparison of the STS and the EuroSCORE risk prediction algorithms. *Can J Cardiol*. 2007;23(11):879-83.
21. Racz MJ, Hannan EL, Isom OW, Subramanian VA, Jones RH, Gold JP, et al. A comparison of short- and longterm outcomes after off-pump and on-pump coronary artery bypass graft surgery with sternotomy. *J Am Coll Cardiol*. 2004;43(4):557-64.
22. Gobran SR, Goldman S, Ferdinand F, Wertan MA, Trace C, Grunkemeier GL, et al. Outcomes after usage of a quality initiative program for off-pump coronary artery bypass surgery: a comparison with on-pump surgery. *Ann Thorac Surg*. 2004;78(6):2015-21.
23. Polomsky M, Puskas JD. Off-pump coronary artery bypass grafting: the current state. *Circ J*. 2012;76(4):784-90.
24. van Dijk D, Nierich AP, Jansen EW, Nathoe HM, Suyker WJ, Diephuis JC, et al.; Octopus Study Group. Early outcome after offpump versus on-pump coronary bypass surgery: results from a randomized study. *Circulation*. 2001;104(15):1761-6.
25. Angelini GD, Taylor FC, Reeves BC, Ascione R. Early and midterm outcome after off-pump and on-pump surgery in Beating Heart Against Cardioplegic Arrest Studies (BHACAS 1 and 2): a pooled analysis of two randomised controlled trials. *Lancet*. 2002;359(9313):1194-9.
26. Cooley DA. Con: beating-heart surgery for coronary revascularization: is it the most important development since the introduction of the heart-lung machine? *Ann Thorac Surg*. 2000;70(5):1779-81.
27. Abdel Aal M, Elnahal N, Bakir BM, Fouda M. Mini-cardiopulmonary bypass impact on blood conservation strategy in coronary artery bypass grafting. *Interact Cardiovasc Thorac Surg*. 2011;12(4):600-4.
28. Beghi C, Nicolini F, Agostinelli A, Borrello B, Budillon AM, Bacciottini F, et al. Mini-cardiopulmonary bypass system: results of a prospective randomized study. *Ann Thorac Surg*. 2006;81(4):1396-400.
29. Hall MW, Hopkins RO, Long JW, Mohammad SF, Solen KA. Hypothermia-induced platelet aggregation and cognitive decline in coronary artery bypass surgery: a pilot study. *Perfusion*. 2005;20(3):157-67.
30. Tsai YT, Lin FY, Lai CH, Lin YC, Lin CY, Tsai CS. On-pump beating-heart coronary artery bypass provides efficacious short- and long-term outcomes in hemodialysis patients. *Nephrol Dial Transplant*. 2012;27(5):2059-65.
31. Panday GF, Fischer S, Bauer A, Metz D, Schubel J, El Shouki N, et al. Minimal extracorporeal circulation and off-pump compared to conventional cardiopulmonary bypass in coronary surgery. *Interact Cardiovasc Thorac Surg*. 2009;9(5):832-6.
32. Ranucci M, Isgrò G. Minimally invasive cardiopulmonary bypass: does it really change the outcome? *Crit Care*. 2007;11(2):R45.
33. Folliguet TA, Villa E, Vandeneiden F, Labored F. Coronary artery bypass graft with minimal extracorporeal circulation. *Heart Surg Fórum*. 2003;6(5):297-301.