



Revista Brasileira de Cirurgia  
Cardiovascular/Brazilian Journal of  
Cardiovascular Surgery

ISSN: 0102-7638

revista@sbccv.org.br

Sociedade Brasileira de Cirurgia  
Cardiovascular

Zhao, Xiaoqi; Gu, Tianxiang  
Dysfunctional Hyperpolarization-Activated Cyclic Nucleotide-gated Ion Channels in  
Cardiac Diseases  
Revista Brasileira de Cirurgia Cardiovascular/Brazilian Journal of Cardiovascular Surgery,  
vol. 31, núm. 2, 2016, pp. 203-206  
Sociedade Brasileira de Cirurgia Cardiovascular  
São José do Rio Preto, Brasil

Available in: <http://www.redalyc.org/articulo.oa?id=398946693022>

- How to cite
- Complete issue
- More information about this article
- Journal's homepage in [redalyc.org](http://redalyc.org)

[redalyc.org](http://redalyc.org)

Scientific Information System

Network of Scientific Journals from Latin America, the Caribbean, Spain and Portugal

Non-profit academic project, developed under the open access initiative

# Dysfunctional Hyperpolarization-Activated Cyclic Nucleotide-gated Ion Channels in Cardiac Diseases

DOI: 10.5935/1678-9741.20160030

## Abstract

Hyperpolarization-activated cyclic nucleotide-gated (HCN) channels are reverse voltage-dependent, and their activation depends on the hyperpolarization of the membrane and may be directly or indirectly regulated by the cyclic adenosine monophosphate (cAMP) or other signal-transduction cascades. The distribution, quantity and activation states of HCN channels differ in tissues throughout the body. Evidence exhibits that HCN

channels play critical roles in the generation and conduction of the electrical impulse and the physiopathological process of some cardiac diseases. They may constitute promising drug targets in the treatment of these cardiac diseases. Pharmacological treatment targeting HCN channels is of benefit to these cardiac conditions.

**Keywords:** Ion Channel Gating. Cardiac Electrophysiology. Electrophysiological Processes.

## INTRODUCTION

The hyperpolarization-activated cyclic nucleotide-gated (HCN) channels comprise a family of cation channels activated by hyperpolarized membrane potentials and stimulated by intracellular cyclic nucleotides. The members of the family are four human isoforms (HCN1-4). Upon hyperpolarization, all four isoforms generate a inward current (I<sub>f</sub>) in the heart and in the nervous system<sup>[1-3]</sup>. The hyperpolarization-activated funny current (I<sub>f</sub>) is a major determinant of cardiac diastolic depolarization and plays a key role in controlling heart rate<sup>[4]</sup>. The expression of HCN channel is age-related, the expression has difference in distinct growth stages, HCN genes are inactivated in ventricular myocytes cells following maturation whereas they expressed in these cells in the fetal and/or neonatal heart<sup>[5-8]</sup>. Three isoforms (HCN1, HCN2, HCN4) are expressed in cardiac tissues, with HCN2 and HCN4 the dominant subtypes. In the healthy adult heart, HCN channels are predominantly expressed in the conduction system, especially in the sinoatrial node, HCN4 has been determined as the principal HCN isoform in sinoatrial node cells. However, some studies demonstrate a significant expression of HCN1 and also a small amount of HCN2 beneath HCN4 in murine sinoatrial cells at the protein level, indicating that the three subunits together contribute to I<sub>f</sub> in these cells<sup>[8,9]</sup>. HCN2 has long been known to be expressed in cardiac ventricular myocytes<sup>[10]</sup>, and recently, the expression of HCN4 and low amounts of HCN1 and in addition a functional role of HCN3 in ventricular myocytes has been demonstrated<sup>[11,12]</sup>, but the expression of HCN channels in healthy adult ventricular myocardium is generally weaker than in the conduction system, so I<sub>f</sub> currents are rarely detectable in normal ventricular myocytes<sup>[13,14]</sup>. The kinetics of HCN1 is the fastest, followed by HCN2, and the slowest is HCN4<sup>[15]</sup>. cAMP can

make the potential of HCN channel open probability curve more positive. HCN2 and HCN4 channels increased 10-20 mV, and HCN1 only 5 mV. HCN channel is also affected by the endocrine system. Studies have demonstrated that HCN2 and HCN4 gene transcription is controlled by thyroid hormone, so it is speculated that HCN2 and HCN4 gene down-regulation level is an important mechanism for bradycardia induced by hypothyroidism. Positive inotropic effect of thyroid hormone may be associated with up-regulation of the channel molecules<sup>[16,17]</sup>.

## HCN AND CARDIOMYOPATHY

A large number of studies have found that in heart failure, atrial fibrillation, myocardial hypertrophy and myocardial infarction, atrial and ventricular HCN2 and HCN4 channels gene express unusually causing atrial or ventricular myocyte I<sub>f</sub> current rise or fall, and all of these may be associated with heart disease fatal arrhythmia<sup>[18]</sup>. Under normal conditions, HCN channels are poorly expressed outside the cardiac pacemaking and conduction system<sup>[11,18-20]</sup>, but it changes during cardiac disease. During hypertension and heart failure, I<sub>f</sub> activity is observed in the ventricular myocardium due to re-expression of HCN genes<sup>[18,21-23]</sup>, several studies proved that I<sub>f</sub> current density and occurrence is significantly greater in hypertrophic cardiomyocytes and end-stage failing hearts and this is directly related to the arrhythmias<sup>[18,21,24-27]</sup>. Basing on the presence of I<sub>f</sub> current in ventricular myocytes isolated from severely hypertrophied rat hearts, the current arrhythmogenic role in cardiac hypertrophy and failure has been inferred<sup>[25]</sup>, and its density is larger in human ventricular myocytes isolated from the hearts of patients with ischemic than in those with dilated cardiomyopathy<sup>[21]</sup>. Stillitano et al.<sup>[18]</sup> demonstrated for the first

time comparing the mRNA and protein expression of HCN subunits in the human atrial and ventricle under normal and heart failure conditions in human heart failure, an upregulation of ventricular HCN2 and HCN4 underlies the increase in functional If current, Michael et al.<sup>[28]</sup> studied the model with ventricular HCN2/4 knockout and identified the enhanced If as an important contributor to the typical electrophysiological alterations observed in the hypertrophic heart, including prolonged ventricular action potentials and lengthened QT intervals. So the overexpression of If may contribute to increased ventricular (and atrial) propensity to arrhythmias which leading to sudden death caused by ventricular tachycardia or fibrillation in heart failure and hypertrophic heart. Cardiac dysfunction is a complex and important problem for people with epilepsy. Recently, a study has showed that cardiac electrophysiology was significantly altered in adult rat models of genetic or acquired epilepsy, with slower heart rate, shorter QRS duration, longer QTc interval, greater standard deviation of RR intervals, and bradycardia, molecular analysis demonstrated significant reductions in cardiac HCN2 mRNA and protein expression in models, providing a molecular correlate of these electrophysiologic abnormalities<sup>[29]</sup>.

## HCN AND ARRHYTHMIAS

HCN channels are essential for cardiac pacemaker and electric conduction. The cardiac hyperpolarization activated cation current (If) is known to be present in region with primary or second pacemaker activity, which is found in non-pacemaking regions of the heart. In pacemaker regions, If is believed to contributed to spontaneous diastolic depolarization. HCN1-knockout animals display a dysfunction of the sinoatrial node resulting in sinus dysrhythmia, sinus pauses and a reduced cardiac output, demonstrating that HCN1 contributes to a stable cardiac rhythm generation<sup>[30]</sup>. Atrial fibrillation is the most common sustained tachyarrhythmia and is one of the major causes of cerebrovascular accident<sup>[31,32]</sup>. Clinical electrophysiology studies in patients have demonstrated that rapid focal activity originating from PV can trigger and maintain atrial fibrillation<sup>[33-35]</sup>. A study found that the pacing-induced spontaneous cation potential and larger If current were observed in the PVs cardiac myocytes from canine using RAP (rapid atrial pacing), and a slow diastolic depolarization in the PVs cells. Meanwhile If current densities were significantly higher. As mentioned above, delayed ventricular repolarization and prolonged QT intervals exist in hypertrophic heart, and they can promote early after-depolarizations and electrical instability, so ventricular HCN channels may play a significant role in the diseased heart by increasing the risk for severe ventricular arrhythmias<sup>[36,37]</sup>. For the patients with advanced atrioventricular block or sick sinus syndrome, pacemakers are implanted. Biological pacemakers have several advantages (improved cardiac output, no need for periodic resizing in a child during maturation, among others), so the efforts to use gene therapy to create a biological pacemaker have been ongoing for many years. And most of the efforts focused on HCN gene family of channels. Plotnikov et al.<sup>[38]</sup> has proved that HCN212 chimeric channel is noteworthy, therefore it resulted in tachycardia when overexpressed in the canine left bundle branch to achieve the desired physiological range.

## HCN CHANNEL BLOCKERS

It is well known that slow heart beat is good for the heart. Therefore, HCN channels become the main target of the drug study controlling rapid arrhythmia. Recently, there are many studies about ivabradine, the first selective blocker of If current. A work done by Kuwabara et al.<sup>[39]</sup> analyzed the effect of ivabradine on survival and rhythmicity in a transgenic mouse model (dnNRSF) of dilated cardiomyopathy, and the founding is although an increased cardiac expression of If accompanied by ventricular tachyarrhythmias and sudden arrhythmic death, the survival was improved significantly when ivabradine was administrated at low doses without affecting heart rate. Ivabradine can markedly reduce If currents in the RAP (rapid atrial pacing) cells in a concentration-dependence manner<sup>[40]</sup>. It has been proved that Ivabradine can reduce heart rate without exerting negative inotropic or vasodilatory effects<sup>[41]</sup>. A recent clinical trial, the Systolic Heart failure treatment with the If inhibitor ivabradine Trial (SHIFT), showed the outcomes of patients with resting heart rates  $\geq 70$  bpm were significantly improved with the heart rate decreased<sup>[42]</sup>. A subgroup analysis of BEAUTIFUL trial (morbidity-mortality evaluation of the If inhibitor ivabradine in patients with coronary artery disease and left ventricular dysfunction study) suggested a specific benefit for patients with heart rates over 70 bpm<sup>[43]</sup>, and echocardiography revealed that the reduction in the left ventricular systolic volume index observed in the ivabradine treatment was related to the degree of heart rate reduction, although the large study showed the composite endpoint was not significantly different between the ivabradine and the placebo groups. All of the results suggest that ivabradine improves cardiac diseases outcome depends on its reduction of heart rates. In the treatment of chronic heart failure,  $\beta$ -receptor blockers are used widely, but some patients cannot tolerant  $\beta$ -receptor blockers (because  $\beta$ -blockers may induce asthma, negative inotropic effect and so on), thus, ivabradine maybe the alternative. Another study examined the effects of ivabradine on survival and arrhythmicity in transgenic mice (dnNRSF-Tg). In this study, ivabradine was used 7 mg/kg per day orally, and the results showed it significantly reduced ventricular tachyarrhythmias and improved survival among dnNRSF-Tg mice while having no significant effect on heart rate or cardiac structure or function. Ivabradine inhibited the abnormal automaticity induced by  $\beta$ -adrenergic stimulation and prevented the increase in automaticity in dnNRSF-Tg mice<sup>[39]</sup>. It is well known that the incidence of sinoatrial node dysfunction increases with age and is accompanied by atrial fibrillation (AF) and other tachyarrhythmia<sup>[44]</sup>. AF is one of the most common arrhythmias. It is estimated that 84% of the patients with AF are over 65 years old, and the incidence rate of AF doubles every 10 years, the incidence rate of AF is 10% in people over 85 years old. It can lead to congestive heart failure, thromboembolism, and cerebrovascular accident. So the mortality rate of AF is very high<sup>[45,46]</sup>. Recent animal study showed that dogs with age-related AF induced by rapid atrial pacing administered ivabradine for intervention, and ivabradine could increase the effective refractory periods in the left atrium and left superior pulmonary vein, reduce the duration of AF after induction, and significantly decrease the inducing rate of AF (the incidence rate of AF

decreased from 50% to 25%). Therefore, ivabradine has potential effects in preventing age-related AF<sup>[47]</sup>. Amiodarone is one of the most frequently used anti-arrhythmic drugs. Amiodarone was shown to inhibit HCN channel currents in oocytes and HEK-cells. To determine the effects of amiodarone under pathological conditions, a study monitored If under the amiodarone treatment in ventricular myocytes from spontaneously hypertensive rats with left ventricular hypertrophy using the whole-cell patch-clamp technique. The findings of the study showed that the incubation of myocytes with amiodarone significantly suppressed If density and downregulated HCN2 and HCN4 expression at both the mRNA and protein level. This is the first study indicating that amiodarone inhibits If under hypertrophied condition<sup>[48]</sup>.

## PERSPECTIVES

With the development of the study on HCN channels, the importance of HCN channels on heart diseases attracted more attention. The biological pacemaker based on HCN channels has attracted much attention, and a lot of *in vitro* and *in vivo* gene transduction experiments proved the feasibility of constructing cardiac biological pacemaker, but its clinical application and safety issues is yet to be confirmed further. In order to avoid side effects of HCN channel blockers, the subtype-selective HCN channel blocker is promising.

**Xiaoqi Zhao, MD; Tianxiang Gu, MD** - Department of Cardiac Surgery ICU, The First Affiliated Hospital of China Medical University, Shenyang, P.R. China

## REFERENCES

- Ludwig A, Zong X, Jeglitsch M, Hofmann F, Biel M. A family of hyperpolarization-activated mammalian cation channels. *Nature*. 1998;393(6685):587-91.
- Robinson RB, Siegelbaum SA. Hyperpolarization-activated cation currents: from molecules to physiological function. *Annu Rev Physiol*. 2003;65:453-80.
- Wahl-Schott C, Biel M. HCN channels: structure, cellular regulation and physiological function. *Cell Mol Life Sci*. 2009;66(3):470-94.
- DiFrancesco D. The role of the funny current in pacemaker activity. *Circ Res*. 2010;106(3):434-46.
- Cho HJ, Furness JB, Jennings EA. Postnatal maturation of the hyperpolarisation-activated cation current, I(h), in trigeminal sensory neurons. *J Neurophysiol*. 2011;106(4):2045-56.
- Huang X, Yang P, Du Y, Zhang J, Ma A. Age-related down-regulation of HCN channels in rat sinoatrial node. *Basic Res Cardiol*. 2007;102(5):429-35.
- Surges R, Brewster AL, Bender RA, Beck H, Feuerstein TJ, Baram TZ. Regulated expression of HCN channels and cAMP levels shape the properties of the h current in developing rat hippocampus. *Eur J Neurosci*. 2006;24(1):94-104.
- Herrmann S, Layh B, Ludwig A. Novel insights into the distribution of cardiac HCN channels: an expression study in the mouse heart. *J Mol Cell Cardiol*. 2011;51(6):997-1006.
- Fenske S, Krause SC, Hassan SI, Becirovic E, Auer F, Bernard R, et al. Sick sinus syndrome in HCN1-deficient mice. *Circulation*. 2013;128(24):2585-94.
- Ludwig A, Zong X, Jeglitsch M, Hofmann F, Biel M. A family of hyperpolarization-activated mammalian cation channels. *Nature*. 1998;393(6685):587-91.
- Herrmann S, Layh B, Ludwig A. Novel insights into the distribution of cardiac HCN channels: an expression study in the mouse heart. *J Mol Cell Cardiol*. 2011;51(6):997-1006.
- Fenske S, Mader R, Scharr A, Paparizos C, Cao-Ehlker X, Michalakakis S, et al. HCN3 contributes to the ventricular action potential waveform in the murine heart. *Circ Res*. 2011;109(9):1015-23.
- Yasui K, Liu W, Opthof T, Kada K, Lee JK, Kamiya K, et al. I(f) current and spontaneous activity in mouse embryonic ventricular myocytes. *Circ Res*. 2001;88(5):536-42.
- Cerbai E, Pino R, Sartiani L, Mugelli A. Influence of postnatal-development on I(f) occurrence and properties in neonatal rat ventricular myocytes. *Cardiovasc Res*. 1999;42(2):416-23.
- Pian P, Bucchi A, Decostanzo A, Robinson RB, Siegelbaum SA. Modulation of cyclic nucleotide-regulated HCN channels by PIP(2) and receptors coupled to phospholipase C. *Pflugers Arch*. 2007;455(1):125-45.
- Gloss B, Trost S, Bluhm W, Swanson E, Clark R, Winkfein R, et al. Cardiac ion channel expression and contractile function in mice with deletion of thyroid hormone receptor alpha or beta. *Endocrinology*. 2001;142(2):544-50.
- Pachucki J, Burmeister LA, Larsen PR. Thyroid hormone regulates hyperpolarization-activated cyclic nucleotide-gated channel (HCN2) mRNA in the rat heart. *Circ Res*. 1999;85(6):498-503.
- Stillitano F, Lonardo G, Zicha S, Varro A, Cerbai E, Mugelli A, et al. Molecular basis of funny current (If) in normal and failing human heart. *J Mol Cell Cardiol*. 2008;45(2):289-99.
- Shi W, Wymore R, Yu H, Wu J, Wymore RT, Pan Z, et al. Distribution and prevalence of hyperpolarization-activated cation channel (HCN) mRNA expression in cardiac tissues. *Circ Res*. 1999;85(1):e1-6.
- Marionneau C, Couette B, Liu J, Li H, Mangoni ME, Nargeot J, et al. Specific pattern of ionic channel gene expression associated with pacemaker activity in the mouse heart. *J Physiol*. 2005;562(Pt 1):223-34.
- Cerbai E, Sartiani L, DePaoli P, Pino R, Maccherini M, Bizzarri F, et al. The properties of the pacemaker current I(f) in human ventricular myocytes are modulated by cardiac disease. *J Mol Cell Cardiol*. 2001;33(3):441-8.
- Sridhar A, Dech SJ, Lacombe VA, Elton TS, McCune SA, Altschuld RA, et al. Abnormal diastolic currents in ventricular myocytes from spontaneous hypertensive heart failure rats. *Am J Physiol Heart Circ Physiol*. 2006;291(5):H2192-8.
- Cerbai E, Mugelli A. I(f) in non-pacemaker cells: role and pharmacological implications. *Pharmacol Res*. 2006;53(5):416-23.
- Cerbai E, Pino R, Porciatti F, Sani G, Toscano M, Maccherini M, et al. Characterization of the hyperpolarization-activated current, I(f), in ventricular myocytes from human failing heart. *Circulation*. 1997;95(3):568-71.
- Cerbai E, Barbieri M, Mugelli A. Occurrence and properties of the hyperpolarization-activated current If in ventricular myocytes from normotensive and hypertensive rats during aging. *Circulation*. 1996;94(7):1674-81.
- Stilli D, Sgoifo A, Macchi E, Zaniboni M, De lasio S, Cerbai E, et al. Myocardial remodeling and arrhythmogenesis in moderate cardiac hypertrophy in rats. *Am J Physiol Heart Circ Physiol*. 2001;280(1):H142-50.

27. Hoppe UC, Jansen E, Südkamp M, Beuckelmann DJ. Hyperpolarization-activated inward current in ventricular myocytes from normal and failing human hearts. *Circulation*. 1998;97(1):55-65.
28. Michael G, Xiao L, Qi XY, Dobrev D, Nattel S. Remodelling of cardiac repolarization: how homeostatic responses can lead to arrhythmogenesis. *Cardiovasc Res*. 2009;81(3):491-9.
29. Powell KL, Jones NC, Kennard JT, Ng C, Urmaliya V, Lau S, et al. HCN channelopathy and cardiac electrophysiologic dysfunction in genetic and acquired rat epilepsy models. *Epilepsia*. 2014;55(4):609-20.
30. Fenske S, Krause SC, Hassan SI, Becirovic E, Auer F, Bernard R, et al. Sick sinus syndrome in HCN1-deficient mice. *Circulation*. 2013;128(24):2585-94.
31. Brundel BJ, Shiroshita-Takeshita A, Qi X, Yeh YH, Chartier D, van Gelder IC, et al. Induction of heat shock response protects the heart against atrial fibrillation. *Circ Res*. 2006;99(12):1394-402.
32. Miyazaki S, Kuwahara T, Kobori A, Takahashi Y, Takei A, Sato A, et al. Prevalence, electrophysiological properties, and clinical implications of dissociated pulmonary vein activity following pulmonary vein antrum isolation. *Am J Cardiol*. 2011;108(8):1147-54.
33. von Bary C, Weber S, Dornia C, Eissnert C, Fellner C, Latzin P, et al. Evaluation of pulmonary vein stenosis after pulmonary vein isolation using a novel circular mapping and ablation catheter (PVAC). *Circ Arrhythm Electrophysiol*. 2011;4(5):630-6.
34. Ehrlich JR, Cha TJ, Zhang L, Chartier D, Melnyk P, Hohnloser SH, et al. Cellular electrophysiology of canine pulmonary vein cardiomyocytes: action potential and ionic current properties. *J Physiol*. 2003;551(Pt 3):801-13.
35. Chen YJ, Chen SA, Chang MS, Lin CI. Arrhythmogenic activity of cardiac muscle in pulmonary veins of the dog: implication for the genesis of atrial fibrillation. *Cardiovasc Res*. 2000;48(2):265-73.
36. Nattel S, Maguy A, Le Bouter S, Yeh YH. Arrhythmogenic ion-channel remodeling in the heart: heart failure, myocardial infarction, and atrial fibrillation. *Physiol Rev*. 2007;87(2):425-56.
37. Marionneau C, Brunet S, Flagg TP, Pilgram TK, Demolombe S, Nerbonne JM. Distinct cellular and molecular mechanisms underlie functional remodeling of repolarizing K<sup>+</sup> currents with left ventricular hypertrophy. *Circ Res*. 2008;102(11):1406-15.
38. Plotnikov AN, Bucchi A, Shlapakova I, Danilo P Jr, Brink PR, Robinson RB. HCN212-channel biological pacemakers manifesting ventricular tachyarrhythmias are responsive to treatment with I(f) blockade. *Heart Rhythm*. 2008;5(2):282-8.
39. Kuwabara Y, Kuwahara K, Takano M, Kinoshita H, Arai Y, Yasuno S, et al. Increased expression of HCN channels in the ventricular myocardium contributes to enhanced arrhythmicity in mouse failing hearts. *J Am Heart Assoc*. 2013;2(3):e000150.
40. Li JY, Wang HJ, Xu B, Wang XP, Fu YC, Chen MY, et al. Hyperpolarization activated cation current (I(f)) in cardiac myocytes from pulmonary vein sleeves in the canine with atrial fibrillation. *J Geriatr Cardiol*. 2012;9(4):366-74.
41. Riccioni G. Ivabradine: recent and potential applications in clinical practice. *Expert Opin Pharmacother*. 2011;12(3):443-50.
42. Swedberg K, Komajda M, Böhm M, Borer JS, Ford I, Dubost-Brama A, et al. Ivabradine and outcomes in chronic heart failure (SHIFT): a randomised placebo-controlled study. *Lancet*. 2010;376(9744):875-85.
43. Fox K, Ford I, Steg PG, Tendera M, Robertson M, Ferrari R; BEAUTIFUL investigators. Heart rate as a prognostic risk factor in patients with coronary artery disease and left-ventricular systolic dysfunction (BEAUTIFUL): a subgroup analysis of a randomised controlled trial. *Lancet*. 2008;372(9641):817-21.
44. Semelka M, Gera J, Usman S. Sick sinus syndrome: a review. *Am Fam Physician*. 2013;87(10):691-6.
45. Chimenti C, Russo MA, Carpi A, Frustaci A. Histological substrate of human atrial fibrillation. *Biomed Pharmacother*. 2010;64(3):177-83.
46. Piccini JP, Hammill BG, Sinner MF, Jensen PN, Hernandez AF, Heckbert SR, et al. Incidence and prevalence of atrial fibrillation and associated mortality among Medicare beneficiaries, 1993-2007. *Circ Cardiovasc Qual Outcomes*. 2012;5(1):85-93.
47. Li YD, Ji YT, Zhou XH, Jiang T, Hong YF, Li JX, et al. Effects of ivabradine on cardiac electrophysiology in dogs with aged-related atrial fibrillation. *Med Sci Monit*. 2015;21:1414-20.
48. Li H, Zhou Y, Jiang B, Zhao X, Li X, Yang X, et al. Dual effects of amiodarone on pacemaker currents in hypertrophied ventricular myocytes isolated from spontaneously hypertensive rats. *Clin Exp Pharmacol Physiol*. 2014;41(9):698-707.