



Acta Universitaria

ISSN: 0188-6266

actauniversitaria@ugto.mx

Universidad de Guanajuato

México

Alaniz-Andrade, Azucena Lucero; Letechipía de León, Consuelo; Ramírez-Santoyo, Rosa  
María; Guzmán-Moreno, Jesús; Vidales-Rodríguez, Luz Elena

Arsenic tolerance in bacterial cultures isolated from metal contaminated soil

Acta Universitaria, vol. 27, núm. 3, mayo-junio, 2017, pp. 9-18

Universidad de Guanajuato

Guanajuato, México

Available in: <http://www.redalyc.org/articulo.oa?id=41652062002>

- How to cite
- Complete issue
- More information about this article
- Journal's homepage in redalyc.org

redalyc.org

Scientific Information System

Network of Scientific Journals from Latin America, the Caribbean, Spain and Portugal

Non-profit academic project, developed under the open access initiative

# Arsenic tolerance in bacterial cultures isolated from metal contaminated soil

Tolerancia al arsénico en cultivos bacterianos aislados de suelo contaminado con metales

Azucena Lucero Alaniz-Andrade\*, Consuelo Letechipía de León\*\*, Rosa María Ramírez-Santoyo\*, Jesús Guzmán-Moreno\*, Luz Elena Vidales-Rodríguez\*<sup>‡</sup>

## ABSTRACT

Several centuries of uninterrupted mining activities in Zacatecas, Mexico has caused soil pollution with toxic metals and metalloids such as arsenic. In this study, the arsenic-tolerance of ten bacterial isolates from a metal contaminated site were analyzed, and high tolerance was observed in both solid (40 mM – 300 mM of sodium dihydrogen arsenate and 4 mM – 25 mM of sodium arsenite) and liquid media (7.2 mM and 11.3 mM arsenite). The arsenic tolerant isolates were identified by biochemical and 16S rRNA-encoding gene amplification analysis as members of the *Bacillus*, *Micrococcus*, and *Acinetobacter* genus. A study of resistance to antibiotics revealed a high prevalence of resistance to beta-lactams and moderate prevalence to nitrofurantoin, vancomycin, and ceftriaxone, suggesting that the antibiotic multiresistance of the isolates is probably related to arsenic tolerance through a plasmid or chromosomally encoded resistance mechanism.

## RESUMEN

Las actividades mineras realizadas durante varios siglos de manera ininterrumpida en el estado de Zacatecas han generado un problema de contaminación de suelos con metales y metaloides como el arsénico. En este estudio, se analizó la tolerancia al arsénico de diez aislados bacterianos observándose una alta tolerancia en medio sólido (40 mM – 300 mM de arseniato y 4 mM – 25 mM de arsenito) y en medio líquido (7.5 mM – 12 mM de arsenito). Los aislados tolerantes a arsénico fueron identificados mediante análisis bioquímico y amplificación del gen ribosomal 16S (rDNA 16S) como miembros de los géneros *Bacillus*, *Micrococcus* y *Acinetobacter*. Un estudio de resistencia a antibióticos reveló alta prevalencia de la resistencia a beta-lactámicos y moderada a nitrofurantoina, vancomicina y ceftriaxona, sugiriendo que la multiresistencia a antibióticos de estos aislados podría estar relacionada a la tolerancia al arsénico mediante un mecanismo de resistencia presente en plásmidos o en el cromosoma.

Recibido: 29 de febrero de 2016

Aceptado: 2 de marzo de 2017

### Keywords:

Arsenic tolerance; antibiotic resistance; *Bacillus*; *Micrococcus*; *Acinetobacter*.

### Palabras clave:

Tolerancia a arsénico; resistencia a antibióticos; *Bacillus*; *Micrococcus*; *Acinetobacter*.

### Cómo citar:

Alaniz-Andrade, A. L., Letechipía de León, C., Ramírez-Santoyo, R. M., Guzmán-Moreno, J., & Vidales-Rodríguez, L. E. (2017). Arsenic tolerance in bacterial cultures isolated from metal contaminated soil. *Acta Universitaria*, 27(3), 9-18. doi: 10.15174/au.2017.1189

## INTRODUCTION

Environmental pollution with metals has become a major threat to human health (Nies, 1999). Arsenic (As) is a highly toxic and carcinogenic metalloid that is widely distributed in the environment as a result of geogenic and anthropogenic activities (Cullen & Reimer, 1989). The most common oxidation states of arsenic in ecosystems are the pentavalent arsenate As[V] and trivalent arsenite As[III], the latter form being the more toxic (Oremland & Stolz, 2003; Rosen, 2002). Despite its toxicity, the ancient and constant exposure of bacteria to arsenic has led to the microbe colonization of arsenic-rich environments throughout the development of metabolism coupled biotransformation processes, i.e. reduction, oxidation and methylation (Bentley & Chasteen, 2002; Muller *et al.*, 2007; Osborne & Erlich, 1976; Rosen, 2002; Silver & Phung, 2005) that affects geochemistry, speciation, and toxicity of this element (Islam *et al.*, 2004; Oremland & Stolz, 2003). Due to the ability of bacteria

\* Unidad Académica de Ciencias Biológicas, Universidad Autónoma de Zacatecas. Avenida Preparatoria s/n, Col. Agronómica II, Zacatecas, Zac., México, C.P. 98066. Tel.: 01 492 9211326. E-mail: luzelenavr@uaz.edu.mx

\*\* Unidad Académica de Ciencias Nucleares, Universidad Autónoma de Zacatecas. Ciprés 10, Col. La Peñuela, Zacatecas, Zac., México, C.P. 98060. Tel.: 01 492 922 7043

<sup>‡</sup> Corresponding author.

to metabolize highly toxic arsenic compounds into a less toxic form (Oremland, Stolz & Hollibaugh, 2004; Weeger *et al.*, 1999), the isolation and study of arsenic-resistant bacteria is attractive for the establishment of processes to ameliorate the bioavailability of arsenic in contaminated soil and water. Many reports of arsenic resistant bacterial isolates from soil and water include a diversity of bacterial genera such as *Microbacterium*, *Alcaligenes*, *Staphylococcus*, *Pseudomonas*, *Corynebacterium*, *Xanthomonas*, *Acinetobacter*, *Flavimonas*, *Micrococcus*, *Bacillus*, *Aeromonas*, *Enterobacter* and *Agrobacterium* (Achour, Cordi, Poupin, Bauda, & Billard, 2010; Goswami *et al.*, 2015; Majumder, Ghosh, Saha, Kole & Sarkar, 2013; Mateos, Ordóñez, Letek & Gil, 2006; Mokashi & Paknikar, 2002; Nagvenkar & Ramaiah, 2010; Novick & Roth, 1968; Osborne & Erlich, 1976; Pepi *et al.*, 2007; Salmassi *et al.*, 2002; Selvi *et al.*, 2014) and some extremophiles (Baker-Austin *et al.*, 2007; Branco, Chung, & Morais, 2008; Bruneel *et al.*, 2003; Chen & Shao, 2009; Gihring, Druschel, McCleskey, Hamers & Banfield, 2001) some of which are proposed for use in bioremediation (Banerjee, Datta, Chattopadhyay & Sarkar, 2011; Goswami *et al.*, 2015; Mateos, *et al.*, 2006).

In Mexico, metal contaminated environments are widely distributed as a consequence of about five centuries of uninterrupted mining activities. However, the study of bacterial diversity is limited in mining sites. The aim of this study was the isolation and characterization of autochthonous arsenic-resistant bacteria with potential application in the biotransformation of arsenic compounds to less toxic ones.

## MATERIALS AND METHODS

### Bacterial isolates

Arsenic tolerance was evaluated in 28 bacterial cultures isolated from soil samples collected from an area located in Guadalupe, Zacatecas, Mexico, which has been previously reported as a metal-contaminated site (Santos *et al.*, 2006).

### Analysis of arsenic tolerance in bacterial isolates

To test the bacterial tolerance to arsenic, different ranges of arsenite and arsenate concentrations were proven based on the levels of bacterial tolerance previously reported (Abbas *et al.*, 2014; Pepi *et al.*, 2007; Selvi *et al.*, 2014). To establish the bacterial tolerance to arsenate in solid media, the Minimum Inhibitory Concentration

(MIC) was determined. Briefly, 28 bacterial isolates were inoculated onto LB agar plates supplemented with increasing concentrations of sodium dihydrogen arsenate ( $\text{NaH}_2\text{AsO}_4$ ) (1 mM to 300 mM), and incubated at 37 °C for 24 h to 48 h to obtain visible bacterial colonies. The lowest concentration of metal salt that prevented growth on the plates was recorded as the MIC. The isolates that showed the highest MICs to sodium dihydrogen arsenate (AsT2, AsT5, AsT15, AsT21, and AsT23) were also analyzed to determine its tolerance to arsenite in solid media by the establishment of MIC at increasing concentrations (1 mM to 27 mM) of sodium arsenite ( $\text{NaAsO}_2$ ). Arsenite tolerance in liquid media was determined by assessing the MIC values (concentration at which bacterial growth was fully inhibited by arsenic ions), as follows: overnight cultures were diluted 1:100 into fresh LB media supplemented with increasing concentrations of  $\text{NaAsO}_2$  (2.0 mM, 4.0 mM, 6.0 mM, 8.0 mM, 10.0 mM, and 12.0 mM) and grown at 37 °C with shaking at 200 rpm for 8 h (time at which the end of exponential growth was reached in cultures), then the optical density was spectroscopically determined at 600 nm ( $\text{OD}_{600\text{nm}}$ ). At least two independent experiments were carried out.

### Molecular identification of As-tolerant isolates

Genomic DNA from Arsenite Tolerant (AsT) isolates was extracted from stationary-phase cultures according to a protocol previously described for gram-positive bacteria (Cutting & Vander Horn, 1990). Amplification of the 16S rDNA gene was accomplished by PCR with the high-fidelity Platinum *Pfx* DNA-polymerase (Invitrogen, Carlsbad, CA) following the manufacturer's recommendations, and using 0.3  $\mu\text{M}$  of the primers f D1 (5' AGAGTTTGATCCTGGCTCAG 3' corresponding to *E. coli* 16S rDNA gene bases 1 to 20) and rP2 (5' ACGGCTACCTTGTTACGACTT 3' corresponding to *E. coli* 16S rDNA gene bases 1487 to 1507) (Weisburg, Barns, Pelletier, & Lane, 1991), and 50 ng of DNA template. PCR was performed for 30 cycles, each consisting of a 30 s denaturation step at 94 °C, a 30 s annealing step at 50.9 °C, and a 1 min extension step at 68 °C. The PCR products were purified with the High Pure PCR Product Purification Kit (Roche, Mannheim, Germany) and directly sequenced with the same primers used in PCR. The 16S rDNA assembled sequences of the AsT isolates were deposited in the GeneBank of the National Center for Biotechnology Information (NCBI) with the following accession numbers: KX866673 (AsT2), KX866674 (AsT5), KX866675 (AsT13), KX866676 (AsT14), KX866677 (AsT15), KX866678 (AsT18), KX866679 (AsT21), KX866681

(AsT23), KX866680 (AsT25), and KT717629 AsT27 (the latter previously accessed as PbT5 for its Pb tolerance). The sequences were compared to the 16S ribosomal RNA sequences database of the GeneBank of the National Center for Biotechnology Information server, and sequence alignment was performed using the BLASTN program with a MEGABLAST algorithm for highly similar sequences and default parameters to determine the sequence identity percentages. The determination of bacterial genus (identity  $\geq 97\%$ ) and species (identity  $\geq 99\%$ ) was based on the parameters previously described (Drancourt *et al.*, 2000). The 16S rDNA sequences were subjected to pairwise and multiple sequence alignment using the ClustalW program, and the molecular evolutionary analyses were conducted using MEGA6 (Tamura, Stecher, Peterson, Filipski & Kumar, 2013). The sequences used as references for the construction of the phylogenetic tree and their respective accession numbers (gi) for NCBI database are presented below: (631252148) *Acinetobacter lwoffii* strain JCM 6840, (645320411) *Acinetobacter haemolyticus* strain ATCC 17906, (645320413) *Acinetobacter johnsonii* strain ATCC 17909, (636558862) *Bacillus simplex* strain LMG 11160, (631251529) *Bacillus simplex* strain NBRC 15720, (343201410) *Bacillus simplex* strain DSM 1321, (651343563) *Micrococcus luteus* strain NCTC 2665, (636560518) *Micrococcus yunnanensis* strain YIM 65004 and (343205881) *Micrococcus endophyticus* strain YIM 56238.

### Morphological and biochemical characterization of As-tolerant strains

Phenotypic analysis of the AsT isolates was based on colony characteristics, Gram staining and biochemical properties such as motility,  $H_2S$  acid production, and acetoin production from glucose, use of citrate and mannitol as carbon source, and presence of enzymatic activities such as urease, lysine, and ornithine decarboxylases, tryptophanase, cytochrome oxidase, catalase and amylase (MacFaddin, 1984). The identification was established according to the Bergey's Manual of Determinative Bacteriology (Holt, Rieg, Sneath, Staley & Williams, 1994).

### Antimicrobial susceptibility of As-tolerant bacteria

Bacterial resistance to different antibiotics was tested by the disk diffusion method and interpreted according to manufacturer's instructions for BBL Antibiotic SensiDiscs (Becton Dickinson Microbiology Systems, Cockeysville, MD) or Multidiscos\* (BIO RAD, Mexico).

## RESULTS AND DISCUSSION

### Tolerance to arsenate ( $AsO_4$ )<sup>3-</sup> and arsenite ( $AsO_3$ )<sup>3-</sup> ions in bacterial strains isolated from metal contaminated soil

Due to the constant exposure to arsenic concentrations in metal-contaminated sites, bacteria have developed adaptive strategies to overcome the toxicity of arsenic. In this work, we found a prevalence of gram-positive and bacilli shape in a group of 28 bacterial strains isolated from contaminated soil with arsenic. This results are consistent with previous reports that indicate a high prevalence of gram-positive species of the *Arthrobacter*, *Bacillus*, and *Micrococcus* genera, although gram-negative genera such as *Pseudomonas*, *Flavobacterium*, *Acinetobacter*, *Agrobacterium*, and *Novcardia* can also be found (Alexander, 1980; Anderson & Cook, 2004; Grant, 1982; Jackson, Harrison & Dugas, 2005). The analysis of arsenic tolerance of 28 bacterial isolates revealed that MICs of sodium arsenate ( $NaH_2AsO_4$ ) and sodium arsenite ( $NaAsO_2$ ) in LB agar were in a range of 0 mM and 300 mM and 4 mM to 25 mM, respectively (tables 1 and 2), five of this isolates (AsT2, AsT5, AsT15, AsT21, and AsT23) showed a high tolerance to arsenate and arsenite when compared with those reported in genera such as *Exiguobacterium*, *Aeromonas*, *Bacillus*, *Pseudomonas*, *Escherichia*, *Acinetobacter* with CMIs of 10 mM to 275 mM of sodium arsenate As(IV) and 20 mM of sodium arsenite As(III) (Anderson & Cook, 2004; Jackson *et al.*, 2005).

The isolates with MICs  $\geq 40$  mM of  $NaH_2AsO_4$  — AsT2 (40 mM), AsT5 (300 mM), AsT13 (60 mM), AsT14 (200 mM), AsT15 (300 mM), AsT18 (300 mM), AsT21 (50 mM), AsT23 (300 mM), AsT25 (300 mM) and AsT27 (40 mM) — were selected to analyze tolerance to arsenite ions ( $AsO_3$ )<sup>3-</sup>. The MICs values were determined between 4 and 25 mM of  $NaAsO_2$  (table 2) and isolates with the highest tolerance to both arsenate ( $AsO_4$ )<sup>3-</sup> and arsenite ( $AsO_3$ )<sup>3-</sup> ions (AsT2, AsT5, AsT15, AsT21, and AsT23) in solid media, were selected to analyze growth in liquid cultures supplemented with increasing concentrations of sodium arsenite ( $NaAsO_2$ ). The AsT23 isolate, with a MIC value of 12 mM of sodium arsenite, showed a higher tolerance to the toxic effects of arsenite ions ( $AsO_3$ )<sup>3-</sup> in liquid media (figure 1e) with respect to isolates AsT2 (7.5 mM), AsT5 (7.5 mM), AsT15 (8 mM) and AsT21 (10 mM) and those previously reported (Banerjee *et al.*, 2011). MICs values indicate that whereas the toxic effects of arsenite ions ( $AsO_3$ )<sup>3-</sup> over bacterial growth in both liquid and solid media were similar in the AsT2 and AsT5, the isolates

AsT15, AsT21, and AsT23 were more sensitive to arsenite ions in liquid media than in solid media (table 2 and figure 1), particularly, AsT15 showed a significant decrease of arsenite tolerance in liquid (MIC 8.0 mM) when compared with tolerance in solid media (MIC 25 mM) (table 2 and figure 1), this results suggest that in the isolates AsT2 and AsT5 the arsenite tolerance is probably related to the presence of an arsenite oxidase (encoded by *aso* genes) that functions as an initial electron donor in aerobic resistance to arsenite allowing the isolates to grow in liquid media under aerobic conditions. On the other hand, the tolerance of AsT15,

AsT21, and AsT23 to arsenite in liquid media probably decreased due to the absence of arsenite oxidase in the isolates, whereupon tolerance may be mediated by alternative arsenic-resistance mechanisms (Silver & Phung, 2005). However, future studies are required to establish the mechanisms involved in arsenic tolerance of the AsT. Thus, the high tolerance observed indicate that the AsT isolates analyzed in this work are potential candidates for the study of molecular mechanisms involved in tolerance/resistance to arsenic, as well as to evaluate the ability to bioaccumulate or biotransform toxic arsenic compounds to less toxic ones.

**Table 1.**  
Tolerance of bacterial isolates to sodium arsenate ( $\text{NaH}_2\text{AsO}_4$ ).

Isolate	MIC $\text{NaH}_2\text{AsO}_4$ (mM)	Isolate	MIC $\text{NaH}_2\text{AsO}_4$ (mM)
AsT-1	20	AsT-15	300*
AsT-2	40*	AsT-16	2.5
AsT-3	4.5	AsT-17	4.0
AsT-4	1.0	AsT-18	300*
AsT-5	300*	AsT-19	4.0
AsT-6	5.0	AsT-20	2.0
AsT-7	2.5	AsT-21	50*
AsT-8	20	AsT-22	4.0
AsT-9	1.5	AsT-23	300*
AsT-10	2.5	AsT-24	1.0
AsT-11	0	AsT-25	300*
AsT-12	0	AsT-26	4.0
AsT-13	60*	AsT-27	40*
AsT-14	200*	AsT-28	4.0


(\*) Highest MIC strains selected for biochemical and 16s gene sequencing identification and analyses of arsenite-tolerance.  
Source: Author's own elaboration.

**Table 2.**  
Tolerance of bacterial isolates to sodium arsenite ( $\text{NaAsO}_2$ ).

Isolate	MIC $\text{NaAsO}_2$ (mM)
AsT-2	8.0*
AsT-5	8.0*
AsT-13	4.0
AsT-14	7.5
AsT-15	25.0*
AsT-18	5.0
AsT-21	13.0*
AsT-23	15.0*
AsT-25	5.0
AsT-27	7.5

(\*) Isolates with the highest MIC were selected to analyze the effects of arsenite ions ( $\text{AsO}_3$ )<sup>3-</sup> over bacterial growth in liquid culture.  
Source: Author's own elaboration.





**Figure 1.** Effect of arsenite ions on growth of different bacterial strains exposed to increasing concentrations of  $\text{NaAsO}_2$  in liquid culture. a) *Bacillus simplex* AsT2; b) *Micrococcus luteus* AsT5; c) *Bacillus simplex* AsT15; d) *Bacillus simplex* AsT21; e) *Micrococcus* sp. AsT23. Overnight cultures were inoculated into fresh LB broth with increasing amounts of  $\text{NaAsO}_2$  and incubated for 8 h at 37 °C and 200 rpm; the  $\text{O.D.}_{600\text{nm}}$  was measured. Data are the average of two independent experiments.

Source: Author's own elaboration.

### Identification of As-tolerant bacteria such as *Bacillus*, *Micrococcus*, and *Acinetobacter* species

Morphological and biochemical characteristics such as cocci shape, positive Gram staining, non-fermenter of glucose and mannitol, as well as the presence of catalase and oxidase enzymatic activities identified the isolates AsT5 and AsT23 as members of the *Micrococcus* genus based on the biochemical characteristics

reported for this genus (table 3) (Holt *et al.*, 1994; Kocur, Kloos & Schleifer, 2006). The isolate AsT13 was identified as a member of the *Acinetobacter* genus based on its non-motil gram-negative cocci shape, aerobic, and non-fermenter metabolism, and the absence of oxidase activity in agreement with characteristics reported for this genus (Hernández Torres, García Vásquez, Yagüe & Gómez Gómez, 2010; Holt *et al.*, 1994) (table 3). On the other hand, isolates AsT2, AsT14, AsT15, AsT18, AsT21, AsT25 and AsT27 were identified as members of the *Bacillus* genus based on

its *bacilli* shape, positive gram staining, ability to produce endospores, aerobic facultative growth and the presence of catalase and amylase enzymatic activities (Holt *et al.*, 1994) (table 3).


We conducted molecular identification based on amplification and sequence analysis of the 16S rDNA ribosomal gene of the arsenic-tolerant isolates that corroborates with phenotypic identification, this analysis identified the AsT2, AsT14, AsT15, AsT18, AsT21, AsT25, and AsT27 isolates as *Bacillus simplex* (99% identity), the AsT13 was identified as *Acinetobacter lwoffii* (99% identity) and the AsT5 and AsT23 isolates as *Micrococcus luteus* (99% identity) and *Micrococcus* sp. (89% identity), respectively. The phylogenetic analysis

grouped the AsT isolates into three different phyla that include firmicutes (*Bacillus* species), actinobacteria (*Micrococcus* species), and proteobacteria (*Acinetobacter* species) (figure 2), in accordance with the wide distribution of arsenic resistance in bacteria. The identification showed a high prevalence of the *Bacillus* genus, which is widely distributed in natural environments (Castillo, Sosa & Scorza, 2004; Felske, Heyrman, Balcaen & De Vos, 2003). Moreover, the results are in agreement with previous studies that report the isolation of bacteria from arsenic contaminated soil that pertain to *Bacillus*, *Pseudomonas*, *Acinetobacter*, *Artrobacter*, and *Micrococcus* genus (Archour, Bauda & Billard, 2007; Megharaj, Avudainayagam & Naidu, 2003), the results indicate that arsenic tolerance is widely distributed between these genus in arsenic contaminated environments.

**Table 3.**  
Morphology and Biochemical characteristics of arsenic-tolerant soil bacteria.

Strain	Morphology					Biochemical properties										
	Colony	Gram stain	Spore former	Motility	H <sub>2</sub> S production	Carbohydrate metabolism				Enzymatic Activities						
						Acid from glucose	Voges Proskauer	Mannitol	Citrate	Urease	Lysine Decarboxylase	Tryptophanase	Ornithine decarboxylase	Cytochrome Oxidase	Catalase	Amylase
AsT-2 ( <i>Bacillus simplex</i> )	Yellow, smooth, round, convex. 4 mm - 6 mm diameter	Bacilli Gram positive	+	+	-	+	-	-	+	-	+	-	-	-	+	-
AsT-5 ( <i>Micrococcus luteus</i> )	Yellow, smooth, round 2 mm - 3 mm diameter	Cocci Gram positive	-	-	-	-	-	-	-	-	+	+	-	+	+	+
AsT-13 ( <i>Acinetobacter lwoffii</i> )	White, creamy, smooth, round, convex 2 mm - 3 mm diameter	Cocci Gram negative	-	-	-	-	-	-	-	-	+	+	-	-	+	-
AsT-14 ( <i>Bacillus simplex</i> )	Irregular, gray, umbonate, flat 4 mm - 6 mm diameter	Long bacilli Gram positive	+	+	-	+	-	-	-	-	-	-	-	+	+	+
AsT-18 ( <i>Bacillus simplex</i> )	Irregular, white, umbonate, flat. 4 mm - 6 mm diameter	Long bacilli Gram positive	+	+	-	+	-	-	-	-	-	-	-	-	+	+
AsT-21 ( <i>Bacillus simplex</i> )	Irregular, white, umbonate, flat 4 mm - 6 mm diameter	Bacilli Gram positive	+	+	-	+	-	-	-	-	+	-	-	+	+	+
AsT-23 ( <i>Micrococcus</i> sp.)	Punctiform, round, light yellow, smooth 1 mm - 3 mm diameter	Cocci Gram positive	-	-	-	-	-	-	-	-	+	-	-	+	+	+
AsT-25 ( <i>Bacillus simplex</i> )	Regular, white, flat, creamy, 4 mm - 6 mm diameter	Bacilli Gram positive	+	+	-	+	-	-	-	-	-	-	-	+	+	+
AsT-27 ( <i>Bacillus simplex</i> )	Regular, cream, flat, shiny, smooth 5 mm - 7 mm diameter	Bacilli Gram positive	+	+	-	+	-	-	+	-	+	-	-	-	+	-

Source: Author's own elaboration.



**Figure 2.** Molecular phylogenetic analysis of AsT isolates. The evolutionary history was inferred by using the Maximum Likelihood method based on the Tamura-Nei model (Tamura & Nei, 1993). The tree with the highest log likelihood (-3200.2978) is shown. The percentage of trees in which the associated taxa clustered together is shown next to the branches. Initial tree(s) for the heuristic search were obtained by applying the Neighbor-Joining method to a matrix of pairwise distances estimated using the Maximum Composite Likelihood (MCL) approach. The tree is drawn to scale, with branch lengths measured in the number of substitutions per site. The analysis involved 19 nucleotide sequences. All positions containing gaps and missing data were eliminated. There were a total of 961 positions in the final dataset. Evolutionary analyses were conducted in MEGA6 (Tamura *et al.*, 2013).

Source: Author's own elaboration.

The identification of the *Acinetobacter* species is consistent with the normal habitat of the most species of this genus, which is widely distributed in different water and soil environments, even in soils with organic pollutants and heavy metals such as arsenic (Archour *et al.*, 2007; Cai, Liu, Rensing & Wang, 2009). However, some species such as *A. baumannii* and *A. johnsonii*, *A. haemolyticus* and *A. lwoffii*, also present as a normal part of the skin microbiota, mucous membranes, respiratory secretions, urine and other clinical samples, and are commonly associated with opportunistic infections especially in patients with a compromised immune systems in hospital environments (Barbe *et al.*, 2004; Guardabassi, Dalsgaard & Olsen, 1999; Hernández Torres *et al.*, 2010). On the other hand, it has been reported that members of the *Acinetobacter* genus have great potential in biotechnological applications, degradation of hydrocarbons in diesel contaminated soils, and bioremediation (Gallego, Loredó, Llamas, Vázquez & Sánchez, 2001; Koma *et al.*, 2001) and *Micrococcus luteus* has been widely used in biotechnological biodegradation and/or bioremediation

processes because it has been shown that this organism is capable of using some contaminants as a food source (Kayode-Isola, Eniola, Olayemi & Igumnugbemi, 2008). Therefore, the results obtained in this work are the basis for future studies that are needed to establish the potential of these arsenic-tolerant isolates to be used in biotechnological processes.

### Antibiotic resistance of Arsenic-tolerant bacteria

The analysis of antibiotic resistance revealed a moderate prevalence of resistance (in at least 50% of isolates) to beta-lactamics such as cefotaxime, ceftazidime, cefuroxime, dicloxacillin and penicillin, whereas resistance to antibiotics such as vancomycin (glycopeptide), nitrofurantoin (sulfamide), and ceftriaxone (cephalosporin) was observed in a 30% to 40% of the arsenic-tolerant isolates. In contrast, a low prevalence of resistance (0% to 10% of isolates) to ampicillin, levofloxacin, cephalothin, chloramphenicol, netilmicin, cefepime, sulfamethoxazole, pefloxacin, tetracycline, ciprofloxacin, and



erythromycin was observed (table 4). The isolation of an *Acinetobacter lwoffii* strain (opportunistic pathogen) resistant to several antibiotics at the sampling site, which is currently urbanized, is indicative of the distribution of multi-resistant pathogens to antibiotics in natural environments, which have also developed a high tolerance and/or resistance to toxic metals. The above result is supported by previous studies that show an association between plasmid-mediated metal tolerance and multiple antibiotic resistance (Calomiris, Armstrong & Seidler, 1984; Tewari, Ramteke, Tripathi, Kumar & Garg, 2013; Timoney, Port, Giles & Spanier, 1998). Therefore, the results contribute to the understanding of the public health problem in Zacatecas related to multi-resistance to antibiotics by human and animal pathogens.

## CONCLUSIONS

The identification and characterization of autochthonous bacteria with high tolerance to arsenic conducted in this study constitutes the first approach to investigate the mechanisms involved in arsenic resistance and its relationship with antibiotic multiresistance to contribute to the discovery of soil microbes with potential applications for biotransformation and waste treatment of arsenic-polluted sites.

## ACKNOWLEDGEMENTS

This work was supported by FOMIX CONACYT-ZAC (Grant ZAC-2013-C01-202597) and the SEP-PROMEP Program.

## REFERENCES

- Abbas, S. Z., Riaz, M., Ramzan, N., Zahid, M. T., Shakoory, F. R., & Rafatullah, M. (2014). Isolation and characterization of arsenic-resistant bacteria from wastewater. *Brazilian Journal of Microbiology*, 45(4), 1309-1315.
- Achour, A. R., Bauda, P., & Billard, P. (2007). Diversity of arsenite transporter genes from arsenic-resistant soil bacteria. *Research Microbiology*, 158(2), 128-137.
- Achour, A. R., Cordi, A., Poupin, P., Bauda, P., & Billard, P. (2010). Characterization of the ars Gene Cluster from Extremely Arsenic-Resistant *Microbacterium* sp. strain A33. *Applied and Environmental Microbiology*, 76(3), 948-955.
- Alexander, M. (1980). *Introducción a la microbiología del suelo* (2nd. ed). México: Libros y Editoriales S.A.
- Anderson, C. R., & Cook, G. M. (2004). Isolation and characterization of arsenate-reducing bacteria from arsenic-contaminated sites in New Zealand. *Current Microbiology*, 48(5), 341-347.
- Baker-Austin, C., Dopson, M., Wexler, M., Sawers, R. G., Stemmler, A., Rosen, B. P., & Bond, P. L. (2007). Extreme arsenic resistance by the acidophilic archaeon *Ferroplasma acidarmanus* Fer1. *Extremophiles*, 11(3), 425-434.
- Banerjee, S., Datta, S., Chattopadhyay, D., & Sarkar, P. (2011). Arsenic accumulating and transforming bacteria isolated from contaminated soil for potential use in bioremediation. *Journal of Environmental Science and Health Part A*, 46(14), 1736-1747.
- Barbe, V., Vallenet, D., Fonknechten, N., Kreimeyer, A., Oztas, S., Labarre, L., Cruveiller, S., Robert, C., Duprat, S., Wincker, P., Ornston, L. N., Weissenbach, J., Marlière, P., Cohen, G. N., & Médigue, C. (2004). Unique features revealed by the genome sequence of *Acinetobacter* sp. ADP1, a versatile and naturally transformation-competent bacterium. *Nucleic Acids Research*, 32(19), 5766-5779.

**Table 4.**  
Antimicrobial susceptibility of arsenic-tolerant soil bacteria.

	Susceptibility of bacterial isolates to antibiotics																				
	1 <sup>a</sup>	2 <sup>b</sup>	3 <sup>c</sup>	4 <sup>c</sup>	5 <sup>c</sup>	6 <sup>c</sup>	7 <sup>c</sup>	8 <sup>d</sup>	9 <sup>c</sup>	10 <sup>c</sup>	11	12 <sup>b</sup>	13 <sup>f</sup>	14 <sup>e</sup>	15 <sup>c</sup>	16 <sup>c</sup>	17 <sup>b</sup>	18 <sup>c</sup>	19 <sup>g</sup>	20 <sup>c</sup>	h
AsT2 ( <i>B. simplex</i> )	S	S	S	R	S	S	S	S	R	R	R	S	R	S	S	S	S	S	S	R	S
AsT5 ( <i>M. luteus</i> )	S	S	S	S	S	S	S	S	S	R	R	R	R	S	S	S	S	S	S	S	S
AsT13 ( <i>A. lwoffii</i> )	S	S	R	S	S	S	S	S	S	S	R	S	R	S	S	S	S	R	S	S	R
AsT14 ( <i>B. simplex</i> )	S	S	S	R	S	S	S	S	R	R	S	S	S	S	S	S	S	S	S	R	S
AsT15 ( <i>B. simplex</i> )	S	S	S	R	S	S	R	S	R	R	R	S	S	S	S	S	S	R	S	S	S
AsT18 ( <i>B. simplex</i> )	S	S	S	R	S	S	S	S	R	S	R	S	R	S	R	S	S	S	R	R	S
AsT21 ( <i>B. simplex</i> )	R	S	S	R	S	S	S	S	R	R	R	S	R	S	S	S	S	S	S	S	S
AsT23 ( <i>Micrococcus</i> sp.)	S	S	S	S	S	S	S	S	S	R	R	R	S	S	S	S	S	S	S	S	S
AsT25 ( <i>B. simplex</i> )	S	S	S	S	S	S	S	S	R	R	R	S	R	S	S	S	S	R	S	R	R
AsT27 ( <i>B. simplex</i> )	S	S	S	S	S	S	S	S	S	S	S	S	S	S	S	S	S	R	S	S	R

1: Ampicillin; 2: Levofloxacin; 3: Cephalothin; 4: Cefotaxime; 5: Col; 6: Netilmicin; 7: Cefepime; 8: Trimeth/Sulpha; 9: Ceftazidime; 10: Cefuroxime; 11: Dicloxacillin; 12: Pefloxacin; 13: Penicillin; 14: Tetracyclin; 15: Amikacin; 16: Gentamicin; 17: Ciprofloxacin; 18: Vancomycin; 19: Erythromycin; 20: Ceftriaxone.  
R: resistant; S:susceptible. Antibiotic concentrations: <sup>a</sup>: 10 µg; <sup>b</sup>: 5 µg; <sup>c</sup>: 30 µg; <sup>d</sup>: Trimethoprim/Sulphamethoxazole 25 µg; <sup>e</sup>: 1 µg; <sup>f</sup>: 10 IU; <sup>g</sup>: 15 µg; <sup>h</sup>: 300 µg. (BBL Antibiotic SensiDiscs, BD. Gram Positive and Gram Negative II Multidisc, BIO-RAD).  
Source: Author's own elaboration.

- Bentley, R., & Chasteen, T. G. (2002). Microbial methylation of metalloids: arsenic, antimony, and bismuth. *Microbiology Molecular Biology Reviews*, 66(2), 250-271.
- Branco, R., Chung, A. P., & Morais, P. V. (2008). Sequencing and expression of two arsenic-resistance strain *Ochrobactrum tritici* SC1124T. *BMC Microbiology*, 8(1), 95-107.
- Bruneel, O., Personné, J. C., Casiot, C., Leblanc, M., Elbaz-Poulichet, F., Mahler, B. J., Le Flèche, A., & Grimont, P. A. (2003). Mediation of arsenic oxidation by *Thiomonas* sp. in acid-mine drainage (Carnoules, France). *Applied Microbiology*, 95(3), 492-499.
- Cai, L., Liu, G., Rensing, C., & Wang, G. (2009). Genes involved in arsenic transformation and resistance associated with different levels of arsenic-contaminated soils. *BMC Microbiology*, 9(1), 4-14.
- Calomiris, J., Armstrong, J. L., & Seidler, R. J. (1984). Association of metal tolerance with multiple antibiotic resistance of bacteria isolated from drinking water. *Applied and Environmental Microbiology*, 47(6), 1238-1242.
- Castillo, C., Sosa, B., & Scorza, J. (2004). Evaluación de la termorresistencia en metabolitos antifúngicos, producidos por esporulados del género *Bacillus*. *Revista de la Sociedad Venezolana de Microbiología*, 24(1-2), 65-67.
- Chen, S., & Shao, Z. (2009). Isolation and diversity analysis of arsenite-resistance bacteria in communities enriched from deep-sea sediments of the Southwest Indian Ocean Ridge. *Extremophiles*, 13(1), 39-48.
- Cullen, W. R., & Reimer, K. J. (1989) Arsenic speciation in the environment. *Chemical Reviews*, 89(4), 713-764.
- Cutting, S. M., & Vander Horn, P. B. (1990). Genetic analysis. In: Harwood, C. R. & Cutting, S. M. (eds.) *Molecular biological methods for Bacillus* (pp. 27-74). Sussex: John Wiley & Sons.
- Drancourt, M., Bollet, C., Carlioz, A., Martelin, R., Gayral, J. P., & Raoult, D. (2000). 16S Ribosomal DNA Sequence Analysis of a Large Collection of Environmental and Clinical Unidentifiable Bacterial Isolates. *Journal of Clinical Microbiology*, 38(10), 3623-3630.
- Felske, A. D., Heyman, J., Balcaen, A., & De Vos, P. (2003). Multiplex PCR screening of soil isolates for novel *Bacillus*-related lineages. *Journal of Microbiology Methods*, 55(2), 447-58.
- Gallego, J. L., Loredó, J., Llamas, J. F., Vázquez, F., & Sánchez, J. (2001). Bioremediation of diesel-contaminated soils: evaluation of potential *in situ* techniques by study of bacterial degradation. *Biodegradation*, 12(5), 325-335.
- Gihring, T. M., Druschel, G. K., McCleskey, R. B., Hamers, R. J., & Banfield, J. F. (2001). Rapid arsenite oxidation by *Thermus aquaticus* and *Thermus thermophilus*: field and laboratory investigations. *Environmental Sciences and Technology*, 35(19), 3857-3862.
- Grant, W. (1982). *Microbiología ambiental*. Zaragoza: Acribia.
- Goswami, R., Mukherjee, S., Rana, V. S., Saha, D. R., Raman, R., Padhy, P. K., & Mazumder, S. (2015). Isolation and Characterization of Arsenic-Resistant Bacteria from Contaminated Water-Bodies in West Bengal, India. *Geomicrobiology Journal*, 32(1), 17-26.
- Guardabassi, L., Dalsgaard, A., & Olsen, J. E. (1999). Phenotypic characterization and antibiotic resistance of *Acinetobacter* spp. isolated from aquatic sources. *Journal of Applied Microbiology*, 87(5), 659-667.
- Hernández Torres, A., García Vázquez, E., Yagüe, G., & Gómez Gómez, J. (2010). *Acinetobacter baumannii* multiresistente: situación clínica actual y nuevas perspectivas. *Revista Española de Quimioterapia*, 23(1), 12-19.
- Holt, J. G., Rieg, N. R., Sneath, P. H. A., Staley, J. T., & Williams, S. T. (1994). *Bergey's Manual of Determinative Bacteriology* (9th ed.). Maryland: Williams and Wilkins.
- Islam, F. S., Gault, A. G., Boothman, C., Polya, D. A., Chamok, J. M., Chatterjee, D., & Lloyd, J. R. (2004). Role of metal-reducing bacteria in arsenic release from Bengal delta sediments. *Nature*, 430(6995), 68-71.
- Jackson, C. R., Harrison, K. G., & Dugas, S. L. (2005). Enumeration and characterization of culturable arsenate resistant bacteria in a large estuary. *Systematic and Applied Microbiology*, 28(8), 727-734.
- Kayode-Isola, T. M., Eniola, K. I. T., Olayemi, A. B., & Igunnugbemi, O. (2008). Response of Resident Bacteria of a Crude Oil-Polluted River to Diesel Oil. *American-Eurasian Journal of Agronomy*, 1(1), 6-9.
- Kocur, M., Kloos, W. E., & Schleifer, K. H. (2006). *The Genus Micrococcus*. In: M., Dworkin, S., Falkow, E., Rosenberg, K. H., Schleifer, & E., Stackebrandt (Eds). (pp. 961-971). The Prokaryotes New York: Springer.
- Koma, D., Hasumi, F., Yamamoto, E., Ohta, T., Chung, S. Y., & Kubo, M. (2001). Biodegradation of long-chain n-paraffins from waste oil of car engine by *Acinetobacter* sp. *Journal of Bioscience and Bioengineering*, 91(1), 94-96.
- MacFaddin, J. F. (1984). *Pruebas bioquímicas para la identificación de bacterias de importancia clínica*. Ciudad de México: Ed. Médica Panamericana S.A.
- Majumder, A., Ghosh, S., Saha, N., Kole, S. C., & Sarkar, S. (2013). Arsenic accumulating bacteria isolated from soil for possible application in bioremediation. *Journal of Environmental Biology*, 34(5), 841-846.
- Mateos, L. M., Ordóñez, E., Letek, M., & Gil, J. A. (2006). *Corynebacterium glutamicum* as a model bacterium for bioremediation of arsenic. *International Microbiology*, 9(3), 207-215.
- Megharaj, M., Avudainayagam, S., & Naidu, R. (2003). Toxicity of hexavalent chromium and its reduction by bacteria isolated from soil contaminated with tannery waste. *Current Microbiology*, 47(1), 51-4.
- Mokashi, S. A., & Paknikar, K. M. (2002). Arsenic (III) oxidizing *Microbacterium lacticum* and its use in the treatment of arsenic contaminated ground-water. *Letters in Applied Microbiology*, 34(5), 258-262.
- Muller, D., Médigue, C., Koechler, S., Barbe, V., Barakat, M., Talla, E., Bonnefoy, V., Krin, E., Arsène-Ploetze, F., Carapito, C., Chandler, M., Courmoyer, B., Cruveiller, S., Dossat, C., Duval, S., Heymann, M., Leize, E., Lieutaud, A., Lièvremon, D., Makita, Y., Mangenot, S., Nitschke, W., Ortet, P., Perdrial, N., Schoepp, B., Siguier, P., Simeonova, D. D., Rouy, Z., Segurens, B., Turlin, E., Vallenet, D., Van Dorselaer, A., Weiss, S., Weissenbach, J., Lett, M. C., Danchin, A., & Bertin, P. N. (2007). A tale of two oxidation states: Bacterial colonization of arsenic-rich environments. *PLoS Genetics*, 3(4)e53, 518-530.

- Nagvenkar, G. S., & Ramaiah, N. (2010). Arsenite Tolerance and Biotransformation Potential in Estuarine Bacteria. *Ecotoxicology*, 19(4), 604-613.
- Nies, D. H. (1999). Microbial heavy metal resistance. *Applied Microbiology and Biotechnology*, 51(6), 730-750.
- Novick, R. P., & Roth, C. (1968). Plasmid-linked resistance to inorganic salts in *Staphylococcus aureus*. *Journal of Bacteriology*, 95(4), 1335-1342.
- Oremland, R. S., & Stolz, J. F. (2003). The ecology of arsenic. *Science*, 300(5621), 939-944.
- Oremland, R. S., Stolz, J. F., & Hollibaugh, J. T. (2004). The microbial arsenic cycle in Mono Lake, California. *FEMS Microbiology and Ecology*, 48(1), 15-27.
- Osborne, F. H., & Erlich, H. L. (1976). Oxidation of arsenite by a soil isolate of *Alcaligenes*. *Applied Bacteriology*, 41(2), 295-305.
- Pepi, M., Volterrani, M., Renzi, M., Marvasi, M., Gasperini, S., Franchi, E., & Focardi, S. E. (2007). Arsenic-resistant bacteria isolated from contaminated sediments of the Orbetello Lagoon, Italy, and their characterization. *Journal of Applied Microbiology*, 103(6), 2299-2308.
- Rosen, B. P. (2002). Biochemistry of arsenic detoxification. *FEBS Letters*, 529(1), 86-92.
- Salmassi, T. M., Venkateswaren, K., Satomi, M., Nealson, K. H., Newman, D. K., & Hering, J. G. (2002). Oxidation of arsenite by *Agrobacterium albertimagni*, AOL15, sp nov., isolated from Hot Creek, California. *Geomicrobiology*, 19(1), 53-66.
- Santos-Santos, E., Yarto-Ramírez, M., Gavilán-García, I., Castro-Díaz, J., Gavilán-García, A., Rosiles, R., Suárez, S., & López-Villegas, T. (2006). Analysis of arsenic, lead and mercury in farming areas with mining contaminated soils at Zacatecas, Mexico. *Journal of the Mexican Chemical Society*, 50(2), 57-63.
- Selvi, M. S., Sasikumar, S., Gomathi, S., Rajkumar, P., Sasikumar, P., & Govindan, S. (2014). Isolation and characterization of arsenic resistant bacteria from agricultural soil, and their potential for arsenic bioremediation. *International Journal of Agricultural Policy and Research*, 2(11), 393-405.
- Silver, S., & Phung, L. T. (2005). A bacterial view of the periodic table: genes and proteins for toxic inorganic ions. *Journal of Industrial Microbiology and Biotechnology*, 32(11), 587-605.
- Tamura, K. & Nei, M. (1993). Estimation of the number of nucleotide substitutions in the control region of mitochondrial DNA in humans and chimpanzees. *Molecular Biology and Evolution*, 10(3), 512-526.
- Tamura, K., Stecher, G., Peterson, D., Filipski, A., & Kumar, S. (2013). MEGA6: Molecular Evolutionary Genetics Analysis version 6.0. *Molecular Biology and Evolution*, 30(12), 2725-2729.
- Tewari, S., Ramteke, P. W., Tripathi, M., Kumar, S., & Garg, S. K. (2013). Plasmid mediated transfer of antibiotic resistance and heavy metal tolerance in thermotolerant water borne coliforms. *African Journal of Microbiology Research*, 7(2), 130-136.
- Timoney, J. F., Port, J., Giles, J., & Spanier, J. (1998). Heavy-metal and antibiotic resistance in the bacterial flora of sediments of New York Bight. *Applied and Environmental Microbiology*, 36(3), 465-472.
- Weeger, W., Lievremon, D., Perret, M., Lagarde, F., Hubert, J. C., Leroy, M., Lett, M. C. (1999). Oxidation of arsenite to arsenate by a bacterium isolated from an aquatic environment. *Biometals*, 12(2), 141-149.
- Weisburg, W. G., Barns, S. M., Pelletier, D. A. & Lane, D. J. (1991). 16S Ribosomal DNA Amplification for Phylogenetic Study. *Journal of Bacteriology*, 173(2), 697-703.