



Revista Mexicana de Biodiversidad

ISSN: 1870-3453

falvarez@ib.unam.mx

Universidad Nacional Autónoma de México
México

García-Mateos, Rosario; Ibarra-Estrada, Emmanuel; Nieto-Angel, Raúl
Antioxidant compounds in hawthorn fruits (*Crataegus* spp.) of Mexico
Revista Mexicana de Biodiversidad, vol. 84, núm. 4, 2013, pp. 1298-1304
Universidad Nacional Autónoma de México
Distrito Federal, México

Available in: <http://www.redalyc.org/articulo.oa?id=42529675036>

- How to cite
- Complete issue
- More information about this article
- Journal's homepage in redalyc.org

redalyc.org

Scientific Information System
Network of Scientific Journals from Latin America, the Caribbean, Spain and Portugal
Non-profit academic project, developed under the open access initiative



Antioxidant compounds in hawthorn fruits (*Crataegus* spp.) of Mexico

Compuestos antioxidantes en frutos de tejocote (*Crataegus* spp.) de México

Rosario García-Mateos^{1,2✉}, Emmanuel Ibarra-Estrada³ and Raúl Nieto-Angel²

¹Departamento de Preparatoria Agrícola, Universidad Autónoma Chapingo. México-Texcoco Km 38.5, 56230 Chapingo, Estado de México. México.

²Instituto de Horticultura, Departamento de Fitotecnia, Universidad Autónoma Chapingo. México-Texcoco Km 38.5, 56230 Chapingo, Estado de México. México.

³Postgrado de Botánica, Colegio de Postgraduados, Campus Montecillo. Carretera México-Texcoco Km 36.5, 56230 Montecillo, Estado de México, México.

✉ rosgar08@hotmail.com

Abstract. The content of phytochemicals associated with the antioxidant activity of the fruits of species of hawthorn (*Crataegus* spp.; Rosacea) located in Mexico is unknown. The objective of the present study was to evaluate the content of phenolic compounds, flavonoids, vitamin C and the antioxidant activity in a selection of Mexican hawthorn species. A quantification was made of total phenols, flavonoids and vitamin C (expressed on mg of phenol, quercetin and ascorbic acid per 100 g of fresh weight, respectively), in 10 g of fruits selected from each genotype; a total of 20 genotypes were sampled, these located in the germplasm bank of the Universidad Autónoma Chapingo, Mexico. The antioxidant activity was evaluated by the DPPH method, expressed as mean inhibitory concentration (IC₅₀). Content of total phenols, flavonoids and vitamin C cannot be associated with the origin and species of the samples. Some genotypes from the state of Chiapas could be considered to have a higher potential for commercial use and consumption due to their nutraceutical quality. Most of the fruits of the 20 genotypes of hawthorn presented a content of phenolic compounds higher than that described for other fruits (lychee fruits, peaches and strawberries); these nutraceutical characteristics provide an added value to the fruit.

Key words: Mexican hawthorn species, antioxidants, phenolic compounds, flavonoids, vitamin C.

Resumen. El contenido de fitoquímicos asociado con la actividad antioxidante de los frutos de especies de tejocote (*Crataegus* spp.; Rosacea) localizadas en México es desconocido. El objetivo del presente estudio fue evaluar el contenido de compuestos fenólicos, flavonoides, vitamina C y la actividad antioxidante, en una selección de especies de tejocote mexicano. La cuantificación de fenoles totales, flavonoides y vitamina C (expresada en mg de ácido gálico, quercetina y ácido ascórbico por 100 gr de peso fresco, respectivamente) en 10 gr de frutos seleccionados de cada genotipo, se muestrearon 20 genotipos localizados en el banco de germoplasma de la Universidad Autónoma Chapingo, México. La actividad antioxidante se evaluó de acuerdo al método del DPPH y se expresó como concentración media inhibitoria (CI₅₀). No se observó una relación del contenido de fenólicos, flavonoides y vitamina C con el origen y la especie de las muestras. Algunos genotipos del estado de Chiapas por su calidad nutracéutica podrían ser consideradas para uso comercial y consumo. La mayoría de los 20 genotipos de tejocote presentaron un contenido de compuestos fenólicos más alto que el descrito para otros frutos (lichi, durazno y fresa), estas características proporcionan un valor agregado a la fruta.

Palabras clave: especies de tejocote mexicanas, antioxidantes, compuestos fenólicos, flavonoides, vitamina C.

Introduction

Recently in Mexico there has been a growing interest in knowledge and management of underutilized fruits, also known as minor, secondary or alternative fruits, such as the case of hawthorn fruits (*Crataegus* spp.) (Nieto-Ángel, 2007). There are descriptions of 150-200 species of this

genus (*Crataegus*) in the world, approximately 13 have been identified in the north and center of Mexico, Phipps (1997) indicates 9 endemic species (Phipps, 1997;1998). Nieto-Ángel (2007) mentions that there is great diversity and variability in its genotypic and phenotypic characteristics. Hawthorn belongs to the Rosacea family, is located principally in cold and temperate climates (Nieto-Ángel, 2007). *Tejocote* is the most widely used term and comes from the nahuatl language, in which *tetl-xocotl* means wild or hard sour fruit; the Nahoas (ancestors of the mexicas)

referred to them as *texococuahutl*, which means the tree of Indian apple (Martínez, 1994) depending on the region where it is located, the tejocote has adopted different common names (Martínez, 1994).

In Mexico it is mainly used as animal feed, ornamental plant; in agro-industry it is used to make regional sweets, in the preparation of punches and conserves because of its high pectin content; it has a high demand mainly in the south-southeast-central region of Mexico in the traditional festivals of "All Saints" because it is put on the table as an offering and consumed as fruit (offerings and piñatas) (Nieto-Ángel, 2007). In traditional Mexican medicine the flowers, leaves, root and the fruits have numerous uses (Martínez, 1994; Ody, 1994).

There are a number of medicinally active phytochemicals that have been isolated from hawthorn plants with most of the data generated in studies of those species that are native to Europe and Asia. Little is known about the North American and Canadian (Edwards et al., 2012), and more specifically about the Mexican *Crataegus* species. One possible barrier to chemical studies of *Crataegus*, has been the perception that Mexican *Crataegus* are taxonomically problematic (Phipps, 1997). The high contents of phenolic compounds such as flavonoids, proanthocyanidins, catechins, phenolic acids, essential oils and terpenoids (Bahorum et al., 1994; Edwards et al., 2012; García-Mateos et al., 2012) explain their use as natural therapy for the treatment of neurodegenerative diseases, in some types of cancer, in the affection of the immunological system and cardiovascular disorders (Craig, 1999; Chang et al., 2002; Cui et al., 2006). Hawthorn extracts exert a wide range of pharmaceutical properties, especially on the cardiovascular system, including cardiotonic, antiarrhythmic, hypotensive, hypolipidemic, and antioxidant activities (Craig, 1999; Barceloux, 2008; Arrieta et al., 2010).

The wide diversity and genotypic variability that exists in the Mexican hawthorn species demands the characterization of its fruits and the determination of its antioxidant properties to be recommended as a food of high nutraceutical value that permits a more efficient agro-industrial use and provides new economic alternatives for the producer. The nutraceutical content in products (jellies, preserves and candied fruit) made from the fruit is unknown. The objective of the present study was to evaluate the content of phenolic compounds, flavonoids, vitamin C and the antioxidant activity in a selection of Mexican hawthorn species.

Materials and methods

Collection of plant material. The present investigation was carried out with hawthorn fruits from the central

plateau and the south of Mexico located in the hawthorn germplasm bank of the Universidad Autónoma Chapingo, located at 19°29' N, 98°53' W, at 2 240 m (García, 1981). The climate is C(Wo) (w)b (I')g, rainy moderate temperate and the driest of sub-humid climates, with rains in summer; the mean annual temperature is 17.8° C and rainfall is approximately 644.8 mm annually. Ten grams of physiologically mature fruits were randomly selected from each genotype. A total of 20 genotypes were sampled; they were originally collected from 3 states of the center and south of Mexico (Puebla, State of Mexico and Chiapas).

Preparation of the extract. One gram of fresh fruit pulp was weighed; each sample was dissolved in 25 ml of ethanol at 95% v/v. After 24 h the volume was adjusted to 25 ml with ethanol at 80% v/v, and the mixture was centrifuged at 1 409 g.

Quantification of total phenols. Quantification was made according to the method proposed by Waterman and Mole (1994). A mixture of phosphowolframic and phosphomolybdic acids in basic medium is used as reactive, and reduced by oxidizing the phenolic compounds, originating blue oxides of wolframic and of molybdenum. For the analysis, 0.5 ml of ethanolic extract were taken, 10 ml of a solution of Na₂CO₃ was added at 10% p/v, was homogenized and the mixture was incubated in darkness at 38° C for 15 min. One ml of the mixture was taken, 3 ml of water was added along with 1 ml of the reactive of Folin Ciocalteu:water (1:1). The mixture was left to set for 15 min in darkness. Finally, the absorbance reading was taken at 600 nm in a Genesys 10s spectrophotometer. The concentration was obtained from a standard curve ($y = 0.0014x$; $R^2 = 0.997$) prepared with gallic acid. Total phenol values are expressed in mg equivalent of gallic acid per 100 g of fresh weight. Each analysis was done in triplicate.

Quantification of flavonoids. One aliquot of 0.5 ml of the supernatant of the ethanolic extract was previously prepared; 1.5 ml of ethanol at 95% v/v was added, along with 0.1 ml of a solution of AlCl₃ at 10% p/v, 0.1 ml of solution of 1 M of potassium acetate and 2.8 ml of distilled water. The mixture was incubated in darkness for 30 min. Absorbance was read in a Genesys 10s spectrophotometer at a wave length of 415 nm. For the quantification, a standard curve was made ($y = 0.0122x - 0.0067$; $R^2 = 0.965$) based on the flavonol quercetin (Chang et al., 2002). Flavonoid values are expressed in mg equivalent of quercetin per 100 g of fresh weight.

Quantification of vitamin C. Quantification of vitamin C was carried out through the determination of ascorbic acid (vitamin C). To a 1 g of sample, 3 ml of metaphosphoric acid at 3% v/v was added, and then the mixture was macerated for 3 min and was filtered. One ml of the

filtrate was taken and the volume brought to 10 ml with the solution of metaphosphoric acid at 3% v/v. Two ml of the extract was taken and 2 ml of the buffer solution, pH= 4 (glacial acetic acid: sodium acetate 5%, 1:1) was added, along with 3 ml of dichloroindophenol and 15 ml of xylene, and the mixture was agitated vigorously. The organic phase was separated and dried with the addition of anhydrous Na_2SO_4 , the mixture was filtered and absorbance was read in a Genesis 10s spectrophotometer at a wavelength of 520 nm. From the standard curve ($y = -2.666x + 0.567$; $R^2 = 0.995$) the concentration of ascorbic acid present in each sample was obtained by means of the following equation: $\text{mg total ascorbic acid/mg} = (C \times V \times 100) / (A \times P)$; where: C= ascorbic acid in the sample, V= final volume, A= ml of aliquot of the solution taken, P= weight or volume of the sample. The concentration of vitamin C was expressed in mg equivalent of ascorbic acid per 100 g of fresh weight.

Evaluation of antioxidant activity. The analysis was performed using the free radical DPPH method (2,2-diphenyl-1-picrylhydrazyl, Sigma-Aldrich), described by Amico et al. (2008). The antioxidant capacity of the samples was observed at 516 nm from the gradual color change of the DPPH (purple) to reduced-DPPH (yellow) (Cotelle et al., 1996). For the analysis, 10 g of sample were macerated (hawthorn fruit pulp) in methanol (GR) for 48 h, the mixture was decanted and the plant residue was placed once more in methanol for the same period of time. The extracts were reunited and the dissolvent was evaporated in a Büchi evaporator. From the highest concentration of the methanolic extract (stock solution) the following dissolutions were prepared in methanol to obtain the concentrations of 0.2, 0.15, 0.1, 0.05 mg ml^{-1} . As references the concentrations 0.1, 0.001 and 0.0001 mg ml^{-1} of quercetin and ascorbic acid were prepared separately. To each concentration of the extracts (1 ml) and of the references 3 ml were added separately of the solution of DPPH (0.1 mM). The mixtures were left at room temperature during 30 min and later the absorbance readings were made at 516 nm. The percentage of DPPH was determined through the formula: $\% \text{ DPPH} = (A_{\text{control}} - A_{\text{sample}}) / A_{\text{control}} \times 100$; where A_{control} is the absorbance of the control (DPPH 0.1 mM) and A_{sample} is the absorbance obtained after 30 min of each sample with DPPH 0.1 mM. The antioxidant activity of the samples was determined through the calculation of the mean inhibitory concentration (IC_{50}), which is the concentration required by the sample to decrease the absorbance of DPPH to 50%. The low absorbance of the reaction mixture indicated high antioxidant activity.

To construct the standard curve ($y = 9.393x + 0.006$; $R^2 = 0.999$) of DPPH, 3.93 mg were weighed and dissolved

in 100 ml of methanol to obtain a concentration of 0.1 mM. From this solution the different concentrations were prepared: 0.01, 0.02, 0.04, 0.06, 0.08 and 0.1 mM of DPPH. Absorbance was measured at 516 nm in a Genesis 10s spectrophotometer; the readings were taken by triplicate and methanol (GR) was used as reagent blank.

Statistical analysis. A Pearson's correlation and analysis of variance (Anova) were carried out, along with the comparison of means of Tukey ($p \leq 0.05$), using the program Statistical Analysis System (SAS, version 8.0) according to a completely randomized experimental design, where each selected genotype was considered as treatment of which 3 replicates were made.

Results

Statistical analysis showed significant differences ($p \leq 0.05$) in the concentration of phenolic compounds among the 20 genotypes of fruits of different species and origin (Fig. 1); the same tendency was observed in the content of flavonoids and vitamin C (ascorbic acid) (Figs. 2, 3). Genotypes 27, 51 and 55 were the ones that presented the highest concentration of phenols, genotypes 27 and 51 have a common geographic origin (Chiapas) in contrast to genotype 55 (Puebla). The lowest concentration was found in genotypes 2 (*Crataegus rosei*) and 5 (*Crataegus aurescens*), both have as origin the state of Puebla. Genotypes 25 and 27 presented the highest content of flavonoids, similarly, the content of these metabolites was also lower in genotypes 2 and 5. Results showed that the content of total phenols and flavonoids can not be associated with the origin and species of the samples (Fig. 1). Genotypes 2, 26 and 66 presented the highest concentration of vitamin C (Fig. 3).

The greatest antioxidant activity (media inhibitory concentration) = 0.6 - 0.10 $\mu\text{g ml}^{-1}$ was significantly higher




Figure 1. Phenolics content of 20 selected genotypes of hawthorn fruits of the germplasm bank of the Universidad Autónoma de Chapingo, Mexico.




Figure 2. Flavonoids content of 20 selected genotypes of *Crataegus* fruits.




Figure 3. Ascorbic acid content of 20 selected genotypes of hawthorn fruits.

($p \leq 0.05$) in the genotypes 18, 22, 51 and 100 than in the genotypes 26 and 27 (Table 1), due to the fact that a lower concentration of sample was required to inhibit in 50% of the solution (0.1 mM) of the radical DPPH (Table 1). The references were quercetin (flavonoid) and ascorbic acid (vitamin C), which presented an antioxidant activity that was higher (IC_{50} of 0.00024 and of 0.00043 $\mu\text{g ml}^{-1}$, respectively) than that found in the methanolic extracts of the fruits of the 20 genotypes selected for the study.

Discussion

Results showed that the content of phenolic compounds was higher than those of flavonoids in the 20 genotypes studied (Edwards et al., 2012; García-Mateos et al., 2012). Table 2 shows a positive correlation between the level of 0.01 total phenols and flavonoids, namely that with increasing phenol content also increases the level of flavonoids in fruits. Bajorum et al. (1994) observed the

same tendency in the extracts of *Crataegus monogyna*. A small proportion of the content of phenolic compounds corresponds to flavonoids, the difference could be due to the presence of procyanidins and phenolic acids, as occurs in other plant species (Cui et al., 2006). Cui et al. (2006) point out a high content of polyphenols in the fruits of *Crataegus pinnatifida* due to the presence of procyanidins (19.7%) of phenols (1.27%; chlorogenic acid) and of flavonoids (0.48%). Svedström et al. (2002a; b) describe the presence of oligomeric procyanidins (condensed tannins) in leaves (1.58% dry weight), in flowers (1.15%) and the highest concentration detected was in fruits (0.15% dry weight) of *Crataegus laevigata*.

For the 20 selected genotypes, the concentrations of phenols observed were found in the interval 52-558 mg in 100 g^{-1} f. w. Edwards et al. (2012) describe 1.84-248.18 mg g^{-1} f. w. on fruits of several species of *Crataegus*. Georgé et al. (2005) point out that the concentration of phenols that they quantified in lychee fruits, peaches and strawberries (> 180 mg of gallic acid equivalent in 100 g f. w.) is lower in comparison with some vegetables (celery, artichokes and Brussels sprouts, > 250 mg equivalent of gallic acid in 100 g f. w.). In the present study most of the hawthorn samples surpassed the highest limit. Therefore, it is considered that they present a high concentration of phenols, but particularly some from Chiapas.

The content of phenolic compounds maintains a low positive correlation with the content of flavonoids (Table 2). In this respect, Bignami et al. (2003) mention that the ecological factors do not influence the concentration of phenolic compounds in the species that were analyzed; the authors point out that the variations of these metabolites may be of genetic origin. However, the environmental factors seem to have an influence on the presence of these metabolites, since Kirakosyan et al. (2003) mention that stress from both cold and drought increased the concentration of the polyphenols and of some flavonoids, as well as the antioxidant activity of the extracts of *C. laevigata* and *C. monogyna*. The phenolic composition of fruits is determined by genetic and environmental factors but may be modified by oxidative reactions during processing and storage (Robards et al., 1999).

In the present study no relationship was found in the genotypes of the antioxidant activity with respect to the content of flavonoids (negative correlation) (Table 2), probably because the flavonoids may be found as glycosides, not studied in the present work. Kirakosyan et al. (2003) indicate that the antioxidant activity is principally found associated to the content of flavonoids, but when these metabolites are found as aglycones, they present higher activity than their respective glycosides due

Table 1. Antioxidant activity of 20 selected genotypes of hawthorn fruits of the germplasm bank of the Universidad Autónoma Chapingo, Mexico.

Genotype	Species	State	Antioxidant activity ($IC_{50} = \mu g\ ml^{-1}$)
2	<i>C. rosei</i> Eggl <i>rosei</i>	Puebla	0.16 e-i
5	<i>C. aurescens</i> Phipps	Puebla	0.20 d-f
8	<i>C. tracyi</i> Ashe ex Eggl	Chiapas	0.11 g-j
16	<i>C. tracyi</i> Ashe ex Eggl	Chiapas	0.13 f-j
18	<i>C. tracyi</i> Ashe ex Eggl	Chiapas	0.08 i-j
19	<i>C. baroussana</i> Eggl	Chiapas	0.18 d-h
22	<i>C. rosei</i> Eggl <i>parryana</i>	Chiapas	0.06 j
25	<i>C. cuprina</i> Phipps	Puebla	0.13 f-j
26	<i>C. gracillior</i> Phipps	Chiapas	0.22 c-e
27	<i>C. gracillior</i> Phipps	Chiapas	0.27 b-c
50	<i>C. greggiana</i> Eggl	Edo. de México	0.46 a
51	<i>C. mexicana</i> Moc y Sessé	Chiapas	0.16 e-i
55	<i>C. mexicana</i> Moc y Sessé	Puebla	0.10 h-j
63	<i>C. mexicana</i> Moc y Sessé	Edo. de México	0.35 b
66	<i>C. gracillior</i> Phipps	Edo. de México	0.14 e-j
69	<i>C. cuprina</i> Phipps	Chiapas	0.20 d-g
72	<i>C. baroussana</i> Eggl	Chiapas	0.35 b
83	<i>C. tracyi</i> Ashe ex Eggl	Chiapas	0.20 d-g
86	<i>C. mexicana</i> Moc y Sessé	Puebla	0.35 b
100	<i>C. mexicana</i> Moc y Sessé	Puebla	0.10 h-j
CV.			15.00

*Data with different letters in a column are statistically different ($p \leq 0.05$). IC_{50} = mean inhibitory concentration; concentration required by the sample to inhibit 50% of DPPH after 30 min.

Table 2. Correlation analysis between characters phenolic compounds, flavonoids, ascorbic acid and antioxidant activity in hawthorn fruits of different genotypes

Character	Flavonoids		Vitamin C		Antioxidant activity	
Phenolic compounds	0.53123	0.01 ^z	-0.17369	0.46	-0.05720	0.81
Flavonoids			-0.39661	0.08	-1.13107	0.58
Vitamin C					0.18412	0.43

Pearson's correlation coefficient and statistical significance between 4 variables. ^zSignificance level according to the Tukey test ($p \leq 0.05$).

to a structure-activity relationship (Robards et al., 1999). Those structures as quercetin (flavonoid), that present hydroxyl groups in positions 3' and 4' of the B ring and OH in C-3 permit stable structures that efficiently capture free radicals, a requirement for maximum antioxidant capacity (Bors et al., 1990; Cotellet et al., 1996; Robards

et al., 1999). This information may explain the moderate antioxidant activity found in the extracts of the present study. In a previous study García-Mateos et al. (2012) identified 4 glycosides of quercetin in hawthorn flowers from 6 species located in the same germplasm bank as the fruits.

Due to the above, it is affirmed that the antioxidant activity of fruits and vegetables is not only associated with the presence of the phenolic compounds and flavonoids, but also is attributed to other metabolites such as carotenoids and vitamin C (Delgado-Vargas and Paredes-López, 2003; Georgé et al., 2005; Materska and Perucka, 2005; Brat et al., 2007). Nieto-Ángel (2007) mentions that the fruits from 16 genotypes from the germplasm bank have a high content of vitamin C (38.6-49.7 mg 100 g⁻¹ f. w.). In the present study, for the 20 selected genotypes, the concentrations of vitamin C observed were found in the interval 27.51-84.15 mg equivalent of ascorbic acid in 100 g⁻¹ f. w. These concentrations were higher than those described by Nieto-Ángel (2007) and genotypes 26 and 51 (origin Chiapas) and 66 (origin Puebla) presented the highest concentrations. However, no correlation was found for this variable with antioxidant activity (Table 2).

Scalbert and Williamson (2003) mention that phenols are the most abundant antioxidants in foods, and that the daily intake is equivalent to a gram of this type of metabolite, and represents 10 times more than the intake of vitamin C, 100 times more than that of vitamin E and 500 times more than that of carotene. Therefore, according to Scalbert and Williamson (2003) the phenolic compounds are the nutraceuticals that contribute the most to antioxidant activity of the fresh foods that are consumed, which makes it possible to infer that the antioxidant activity in some hawthorn fruits is due to the high content of phenols and in others possibly to a synergetic effect of vitamin C and of the flavonoids, as pointed out by Georgé et al. (2005).

The presence of components with medicinal properties gives importance to the exploitation of hawthorn fruits in the field of herbal preparations and food with medicinal and nutraceutical interest.

The fruits of most of the 20 selected genotypes of hawthorn species from the germplasm bank of the Universidad Autónoma Chapingo presented a higher content of phenolic compounds than that described for other fruits. Content of phenols, flavonoids and vitamin C cannot be associated with the origin and species of the genotypes. The fruits of genotypes 18 and 22 presented the highest antioxidant activity. The highest values of the content of phenolic compounds were identified in the fruits of genotypes 27, 51 and 55 and in genotypes 26, 51 and 66 the highest content of vitamin C was observed. Some of the genotypes from the state of Chiapas could be considered to have a higher potential for commercial use and consumption due to their nutraceutical quality. Phytochemicals were evaluated and provide evidence of the antioxidant activity of the fruits; their presence explains some of the medicinal properties attributed to species from the Mexican *Crataegus*. It is recommended

to evaluate the nutraceuticals content in products made of hawthorn fruits (jellies, preserves and candied fruit) due to the cultural consumption activity in our country.

Literature cited

- Amico, V., R. Chillemi, S. Mangiafico, C. Spatafora and C. Tringali. 2008. Polyphenol-enriched fractions from Sicilian grape pomace: HPLC-DAD analysis and antioxidant activity. *Bioresource Technology* 99:5960-5966.
- Arrieta, J., D. Siles-Barrios, J. García-Sánchez, B. Reyes-Trejo and M. E. Sánchez-Mendoza. 2010. Relaxant effect of the extracts of *Crataegus mexicana* on guinea pig tracheal smooth muscle. *Pharmacognosy Journal* 2:40-46.
- Bahorum, T., F. Trotin, J. Pommery, J. Vasseur and M. Pinkas. 1994. Antioxidant activities of *Crataegus monogyna* extracts. *Planta Medica* 60:323-328.
- Bignami, C., M. Paolucci, A. Scossa and G. Bertazza. 2003. Preliminary evaluation of nutritional and medicinal components of *Crataegus azarolus* fruits. *Acta Horticulturae* 597:95-100.
- Barceloux, D. G. 2008. Hawthorn (*Crataegus* Species). In *Medical toxicology of natural substances: foods, fungi, medicinal herbs, plants, and venomous animals*, D. G. Barceloux (ed.). John Wiley and Sons, New York. p. 510-513.
- Bors, W., W. Heller, C. Michel and S. Manfred. 1990. Flavonoids as antioxidants: determination of radical-scavenging efficiencies. *Methods in Enzymology* 186:343-355.
- Brat, P., S. Georgé, A. Bellany, L. Du Chautaut, L. Mennen, A. Scalbert and M. J. Amiot-Carlin, 2007. Determination of the polyphenol content of fruit and vegetables. Establishment of a database and estimation of the polyphenol intake. In *Proceedings of the First International Symposium on Human health effects of fruits and vegetables*. August, 2005, Québec, Canada. International Society for Horticultural Science, Leiden, Nederland. p. 61-69.
- Chang, C., M. Yang, H. Wen and J. Chern. 2002. Estimation of total flavonoids content in propolis by two complementary colorimetric methods. *Journal Food and Drug Analysis* 10:176-182.
- Chang, Q., Z. Zuo, F. Harrison, M. S. Chow and J. 2002. Hawthorn. *Clinical Pharmacology* 42:605-612.
- Cotelle, N., J. Jery, J. Wallet and E. M. Gaydou. 1996. Antioxidant properties of hidroxy-flavones. *Free Radical Biology and Medicine* 20:35-43.
- Craig, W. J. 1999. Health-promoting properties of common herbs. *The American Journal of Clinical Nutrition* 70 (suppl.):491S-499S.
- Cui, N., K. Nakamura, S. Tian, H. Kayahara and Y. Tian. 2006. Polyphenolic content and physiological activities of chinese hawthorn extracts. *Bioscience Biotechnology and Biochemistry* 70:2948-2956.
- Delgado-Vargas, F. and O. Paredes-López. 2002. Natural colorants for food and nutraceutical uses. CRC Press, Boca Raton. 344 p.
- Edwards, J. E., P. N. Paula, N. Brown, N. Talent, T. A. Dickinson

- and P. R. Shipley. 2012. A review of the chemistry of the genus *Crataegus*. *Phytochemistry* 79:5-26.
- García, E. 1981. Modificaciones al sistema de clasificación climática de Köpen. Edit. UNAM, México, D. F. 246 p.
- García-Mateos, R., L. Aguilar-Santelises, M. Soto-Hernández, R. Nieto-Angel and G. Kite. 2012. Compuestos fenólicos totales, flavonoides y actividad antioxidante en las flores de *Crataegus* spp. *Agrociencia* 46:651-662.
- Georgé, S., P. Brat, P. Alter and M. J. Amiot-Carlín. 2005. Rapid determination of polyphenols and vitamin C in plants derived products. *Journal of Agricultural and Food Chemistry* 53:1370-1373.
- Kirakosyan, A., E. Seymour, P. B. Kaufman, S. Warber, S. Bolling and S. C. Chang. 2003. Antioxidant capacity of polyphenolic extracts from leaves of *Crataegus laevigata* and *Crataegus monogyna* (Hawthorn) subjected to drought and cold stress. *Journal of Agricultural and Food Chemistry* 51:3973-3976.
- Materska, M. and I. Perucka. 2005. Antioxidant activity of the main phenolic compounds isolated from hot pepper fruits (*Capsicum annuum* L.). *Journal of Agricultural and Food Chemistry* 53:1750-1756.
- Martínez, M. 1994. Catálogo de nombres vulgares y científicos de plantas mexicanas. Fondo de Cultura Económica, México, D. F. 1247 p.
- Nieto-Ángel, R. 2007. Colección, conservación y caracterización del tejocote (*Crataegus* spp.). In *Frutales nativos, un recurso fitogenético de México*, R. Nieto-Ángel (ed.). Universidad Autónoma Chapingo-Servicio Nacional de Inspección y Certificación de Semillas (Snics-Sagarpa), Chapingo, México. p. 25-107.
- Ody, P. 1993. The herb society's complete medicinal herb. Dorling Kindersley Publishing. New York. 192 p.
- Phipps, J. B. 1997. Monograph of northern Mexican *Crataegus* (Rosaceae, Subfam. Maloideae). *SIDA Botanical Miscellany* 15:1-94.
- Phipps, J. B. 1998. Introduction to the red fruited hawthorns (*Crataegus*, Rosaceae) of western North America. *Canadian Journal of Botany* 76:1863-1899.
- Robards, K., D. Prenzler, G. Tucker, P. Swatsitang and W. Glover. 1999. Phenolic compounds and their role in oxidative processes in fruits. *Food Chemistry* 66: 401-436.
- Scalbert, A. and G. Williamson. 2000. Dietary intake and bioavailability of polyphenols. *Journal of Nutrition* 130:2073S-2085S.
- Svedström, U., H. Vuorela, R. Kostianen, K. Huovinen, I. Laakso and R. Hiltunen. 2002. High-performance chromatographic determination of oligomeric procyanidins from dimers up to the hexamer in hawthorn. *Journal of Chromatography A* 968:51-60.
- Svedström, U., H. Vuorela, R. Kostianen, J. Tuominen, J. Kokkonen, J. Rauha, I. Laakso and R. Hiltunen. 2002. Isolation and identification of oligomeric procyanidins from *Crataegus* leaves and flowers. *Phytochemistry* 60:821-825.
- Waterman, P. G. and S. Mole. 1994. Analysis of phenolic plant metabolites. *Methods in Ecology*. Blackwell Scientific Publications, Oxford. 238 p.