



Revista Mexicana de Biodiversidad

ISSN: 1870-3453

falvarez@ib.unam.mx

Universidad Nacional Autónoma de México
México

Montoya-Mendoza, Jesús; Jiménez-Badillo, María de Lourdes; Salgado-Maldonado, Guillermo
Helminths of *Ocyurus chrysurus* from coastal reefs in Veracruz, Mexico
Revista Mexicana de Biodiversidad, vol. 85, núm. 3, septiembre-, 2014, pp. 957-960
Universidad Nacional Autónoma de México
Distrito Federal, México

Available in: <http://www.redalyc.org/articulo.oa?id=42532096016>

- How to cite
- Complete issue
- More information about this article
- Journal's homepage in redalyc.org

redalyc.org

Scientific Information System
Network of Scientific Journals from Latin America, the Caribbean, Spain and Portugal
Non-profit academic project, developed under the open access initiative



Research note

Helminths of *Ocyurus chrysurus* from coastal reefs in Veracruz, Mexico

Helmintos de *Ocyurus chrysurus* de los arrecifes de Veracruz, México

Jesús Montoya-Mendoza^{1✉}, María de Lourdes Jiménez-Badillo¹ and Guillermo Salgado-Maldonado²

¹Laboratorio de Evaluación y Manejo de Recursos Pesqueros, Instituto de Ciencias Marinas y Pesquerías, Universidad Veracruzana. Hidalgo 617, Col. Río Jamapa, 94290 Boca del Río, Veracruz, México. Boca del Río, Veracruz, Mexico.

²Laboratorio de Helmintología, Instituto de Biología, Universidad Nacional Autónoma de México. Apartado postal 70-153, 04510 México, D. F., Mexico.

✉jesusmontoya@itboca.edu.mx

Abstract. This study is a survey of helminths of the yellowtail snapper, *Ocyurus chrysurus*, collected in the islands de Enmedio and Anegada de Afuera reefs, Veracruz, Mexico. Nineteen species were found in fish collected in Anegada de Afuera reef: 9 digeneans (7 adults, and 2 metacercariae); 5 nematodes (2 adults and 3 larvae); 3 monogeneans; 1 acanthocephalan (adult) and 1 cestode (larva); while in the Isla de Enmedio reef 15 of these species were recorded. 7 species are new records for this host, increasing from 17 to 24 species known in the Gulf of Mexico and the Caribbean.

Key words: marine fish, parasites.

Resumen. En este estudio se presentan los resultados del análisis helmintológico de la rubia, *Ocyurus chrysurus*, capturados en los arrecifes de las islas de Enmedio y Anegada de Afuera, Veracruz, México. En el arrecife Anegada de Afuera se registraron 19 especies de helmintos: 9 digéneos (7 adultos y 2 metacercarias), 5 nematodos (2 adultos y 3 larvas), 3 monogéneos, un acantocéfalo (adulto) y un céstodo (larva), en tanto, en el arrecife de la Isla de Enmedio se registraron 15 de estas especies. 7 especies son nuevos registros para este hospedero, aumentando de 17 a 24 especies conocidas para la región del golfo de México y el Caribe.

Palabras clave: peces marinos, parásitos.

The biological richness in coastal reefs around the world is remarkable. In a recent series of papers Justine (2010) outlines the richness of helminth parasites found in fish from the Great Australian Barrier Reef, and the continuity of the system enables an homogeneous distribution of hosts and parasites. In America, coral reefs in the Gulf of Mexico also form a barrier running from the Southern Caribbean Sea in Quintana Roo, to the Veracruz coasts (Jordán-Dahlgren and Rodríguez-Martínez, 2003), extending the distribution of species such as the lutjanid fish *Ocyurus chrysurus* (Bloch, 1791). This fish is common in this region, and has a total length ranging from 210 to 600 mm (Gutiérrez-Benítez, 2012). Given the increasing interest in producing fingerlings of this species under controlled conditions, wild specimens were examined to survey their parasitic community, in order to improve management practices. Helminth parasites of *O. chrysurus* have been reported in the Atlantic coast of the US (Linton,

1910; Chandler, 1935; Manter, 1947; Overstreet, 1969); the Atlantic coast of Central America, Panamá and Bimi Islands (Sogandares-Bernal, 1959; Sogandares-Bernal and McAlister-Sogandares, 1961); Jamaica (Nahhas and Carlson, 1944); Belize (Fischthal, 1977), and Cuba (Zhukov, 1976). However, there are no reports for this species in Mexico.

From July 2011 to March 2012, 54 specimens of *O. chrysurus* were examined for helminth parasites. Out of these, 27 fish were captured in the Isla de Enmedio reef (19°16'00" N, 95°56'19" W), and 27 in the Anegada de Afuera reef (19°10'14" N, 95°52'14" W), 10 km away from each other. Fish were caught by hook-and-line and transported alive to the laboratory for immediate helminthological examination. All tissues and organs were reviewed under a stereoscopic microscope. The external examination included skin, scales, fins, gills, eyes, nostrils, mouth and anus. Gills were removed and analyzed separately in Petri dishes filled with seawater, to facilitate the ectoparasites handling. Then, all specimens

Table 1. Prevalence, mean intensity, and site of infection of helminth parasites of the yellowtail snapper, *Ocyurus chrysurus*, from coastal reefs of Veracruz, Mexico. n, number of hosts infected. Abbreviations for life stages are: *mt*, metacercarie; *p*, plerocercoid; *l*, larva; abbreviations for sites are: f, fins; g, gill; gb, gall bladder; i, intestine; ms, mesenteries; s, stomach

Species	CNHE	Site	Reef Isla de Enmedio			Reef Anegada de Afuera		
			n (% prevalence)	Mean intensity (\pm SD)	Range	n (% prevalence)	Mean intensity (\pm SD)	Range
Trematoda								
<i>Aponurus laguncula</i> Looss, 1907	9130, 9131	i, s	1 (3.7)	1 \pm -	1	12 (44.4)	1 \pm 0	1-1
<i>Megasolena</i> sp.*†	9132	i				1 (3.7)	4 \pm -	4
<i>Preptetos trulla</i> Linton, 1907†	9133, 9134	i	17 (62.9)	16.8 \pm 19.5	1-61	11 (40.7)	84.5 \pm 191.3	1-638
<i>Deretrema fusillus</i> Linton, 1910 †‡	9135	i, gb				8 (29.6)	1.6 \pm 1.1	1-4
<i>Siphodera vinalwardsii</i> (Linton, 1901)	9136, 9137	i	1 (3.7)	1 \pm -	1	2 (7.4)	4 \pm 4.2	1-7
<i>Stephanostomum casum</i> (Linton, 1910)	9138, 9139	i	9 (33.3)	2.8 \pm 2.2	1-7	2 (7.4)	1 \pm 0	1-1
<i>Lecithochirium floridense</i> (Manter, 1934)	9140, 9141	i, s	4 (18.5)	1 \pm 0	1-1	8 (29.6)	1.3 \pm 0.5	1-2
Didymozoidae ^{mt}	9142, 9143	i	1 (3.7)	1 \pm -	1	1 (3.7)	3 \pm -	3
Trematoda ^{mt}		f	22 (81.5)	4.1 \pm 1.8	1-10	14 (51.9)	2.6 \pm 1.2	1-6
Monogenea								
<i>Euryhaliotrema torquacirrus</i> (Zhukov, 1976) †	9144, 9145	g	17 (63)	6.6 \pm 6.1	1-24	20 (74)	3.8 \pm 2.7	1-11
<i>Haliotrematoides heteracantha</i> (Zhukov, 1976) †	9146, 9147	g	8 (29.6)	3.3 \pm 4.1	1-12	3 (11.1)	3.7 \pm 0.6	3-4
<i>Microcotyloides incisa</i> (Linton, 1910) *	9148, 9149	g	7 (25.9)	6.6 \pm 6.1	1-16	22 (81.5)	3.2 \pm 2.6	1-11
Cestoda								
Tetraphyllidea ^p	9150, 9151	i	7 (25.9)	6 \pm 6.9	1-17	9 (33.3)	3.8 \pm 3.2	1-8
Nematoda								
<i>Dichelyne bonacii</i> González-Solis, Argáez-García and Guillén-Hernández, 2002*	9152, 9153	i	3 (11.1)	1 \pm 0	1-1	7 (25.9)	2.1 \pm 0.8	1-6
<i>Hysterothylacium reliquens</i> (Norris and Overstreet, 1975) *	9154, 9155	i, ms	1 (3.7)	2 \pm -	2	4 (14.8)	1 \pm 0	1-1
<i>Contracaecum</i> sp. ^l *	9156, 9157	i, ms	3 (11.1)	1 \pm 0	1-1	9 (33.3)	1.3 \pm 0.5	1-2
<i>Hysterothylacium</i> sp. ^l	9158, 9159	i	2 (7.4)	2 \pm 0	2-2	6 (22.2)	1.3 \pm 0.9	1-3
<i>Spirocerca</i> sp.*	9160	i				1 (3.7)	1 \pm -	1
Acantocephala								
<i>Rhadinorhynchus dujardini</i> Golvan, 1969*†‡	9161	i				1 (3.7)	1 \pm -	1
Total helminth species			15			19		

* New host records.

† New record for Mexico.

‡ New record for the Gulf of Mexico.

were dissected to perform internal examinations, including mesenteries, liver, kidneys and gonads. The entire digestive system was placed in Petri dishes with saline solution at 0.75% for examination in detail. All parasites found were directly fixed in hot ($\approx 90^{\circ}\text{C}$) 4% formalin, and preserved in 70% alcohol.

For the taxonomical study, trematodes, monogeneans, cestodes and acanthocephalans were treated with Mayer paracarmine, Gömöri trichrome and/or Erlich's hematoxyline staining techniques, based on the following steps: dehydration in alcohol series, clarification with clove oil, and mounting in total preparations with Canada balsam. Nematodes were studied in temporary slides, and clarified with pure glycerin to be later preserved in 70% alcohol. Some monogeneans specimens were fixed in temporary preparations with ammonium picrate, in order to analyze their sclerotized structures (Vidal-Martínez et al., 2001). Voucher parasite specimens were deposited at the Colección Nacional de Helmintos (CNHE), Instituto de Biología, UNAM, México D. F. Prevalence (percentage of infected hosts) and mean intensity of infection (mean number of parasites per infected fish), were calculated following Bush et al. (1997).

Total length of 27 *O. crysurus* examined from Isla de Enmedio reef ranged from 233 to 450 mm (306 ± 730 mm), while the weight range was 110 to 857 g (320.8 ± 237.8 g). In Anegada de Afuera reef, the 27 fish were largest, with measures from 355 to 535 mm (462 ± 560 mm), and from 428 to 1363 g (909.3 ± 308 g). All examined fish were more than 1 year old. The helminth parasites inventory of sampled hosts included 19 species, as follows: 9 are digeneans (7 adults, 2 metacercariae), 5 nematodes (2 adults, 3 larvae), 3 monogeneans, 1 acanthocephalan (adult) and 1 cestode (larva) (Table I). In general, samples from both reefs included the same species, with the exception of 4 species absent in Isla de Enmedio: the trematodes *Deretrema fusillus* and *Megasolena* sp., the nematode *Spirocerca* sp., and the acanthocephalan *Rhadinorhynchus dujardini*, collected in larger fish from the Anegada de Afuera reef.

The monogenean *Euryhalitrema torquacirrus* showed the highest prevalence in both sites (69%). A second monogenean species, *Microcotyloides incisa* exhibits high prevalence (53%), with higher values in Anegada de Afuera reef. Other species with high prevalence in one site or the other were adults of trematodes *Aponurus laguncula* and *Prepetos trulla*. Table I displays prevalence of all remaining species which vary from 3.7% (a single fish infected in the sample) to 33.3%. The highest intensity was for *P. trulla*, with 638 trematodes found in a single fish from Anegada de Afuera reef. This trematode exhibits high prevalence and abundance in both locations. Most

helminths found in this study had been previously registered in other lutjanid fish from the Gulf of Mexico and the Caribbean, including *Lutjanus apodus* (Walbaum, 1792), *L. cyanopterus* (Cuvier, 1828), *L. griseus* (Linnaeus, 1758), *L. purpureus* (Poey, 1866) and *L. synagris* (Linnaeus, 1758) (Vélez, 1987; Bunkley-Williams et al., 1996; Dyer et al., 1998; Fuentes et al., 2003; Cortés et al., 2009; Argáez-García et al., 2010). However, *O. crysurus* is a new host record for 7 species. It should be noted that 1 unidentified metacercariae was also recovered with high prevalences in both localities, but was higher in Isla de Enmedio fish (81.5%), than in those from Anegada de Afuera (51.9%).

These data show that for a highly vagile fish as *O. crysurus*, composition and abundance of helminth parasites in 2 nearby localities are similar; and differences may be explained by intrinsic host factors and local conditions of each reef system, mainly associated to size and feeding habits. A significant underlying factor to the parasites presence may be the relevance of intermediate host distribution and abundance, as fish feed on the most abundant prey. This study reveals, on one hand, that *O. crysurus* in the Veracruz reef systems, have a rich parasitic community, with potential interchange with other sympatric Lutjanid species; and, on the other hand, that parasite transmission strategies in reef conditions are highly effective, as most hosts are infected.

This paper was sponsored by a postdoctoral fellowship supported by Conacyt (scholarship 164382), granted to the first author in the laboratory of Ma. Lourdes Jiménez Badillo. Thanks to Yuri Okolodkov for translation of Russian papers.

Literature cited

- Argáez-García, N., S. Guillén-Hernández and M. L. Aguirre-Macedo. 2010. Intestinal helminths of *Lutjanus griseus* (Perciformes: Lutjanidae) from 3 environments in Yucatán (Mexico), with a checklist of its parasites in the Gulf of Mexico and Caribbean region. *Revista Mexicana de Biodiversidad* 81:903-912.
- Bunkley-Williams, L., W. G. Dyer and E. H. Jr. Williams. 1996. Some aspidogastriid and digenean trematodes of Puerto Rican marine fishes. *Journal of Aquatic Animal Health* 8:887-92.
- Bush, A. O., K. D. Lafferty, J. M. Lotz and A. W. Shostak. 1997. Parasitology meets ecology on its own terms: Margolis et al. revisited. *Journal of Parasitology* 83:575-583.
- Chandler, A. C. 1935. Parasites of fishes in Galveston Bay. *Proceedings of the United States National Museum* 83:123-157.
- Cortés, J., J. Valbuena and G. Manrique. 2009. Nemátodos parásitos de *Lutjanus synagris* (Linnaeus, 1758) y *Lutjanus analis* (Cuvier, 1828) (Perciformes, Lutjanidae) en las zonas

- de Santa Marta y Neguanje, Caribe Colombiano. *Revista Médica Veterinaria y Zootecnia* 56:23-31.
- Dyer, G., H. W. Ernest Jr. and L. Bucybunkley-Williams. 1998. Some digenetic trematodes of marine fishes from Puerto Rico. *Caribbean Journal of Science* 34:141-146.
- Fischthal, J. 1977. Some digenetic trematodes of marine fishes from the Barrier Reef and Reef Lagoon of Belize. *Zoologica Scripta* 6:81-88.
- Fuentes, J. L., R. C. Silva and Y. Reyes. 2003. Parásitos en juveniles de *Lutjanus griseus* (Pisces: Lutjanidae) de la Laguna de la Restinga, Isla de Margarita, Venezuela. *Interciencia* 28:463-468.
- Gutiérrez-Benítez, O. 2012. Aspectos biológico pesqueros de la rubia *O. chrysurus* (Bloch, 1791) en Antón Lizardo, Veracruz, México. Tesis maestría, Instituto de Ciencias Marinas y Pesquerías. Universidad Veracruzana, Boca del Río. 58 p.
- Jordán-Dahlgren, E. and R. E. Rodríguez-Martínez. 2003. The Atlantic coral reefs of México. In *Latin American coral reefs*, J. Cortés J. (ed.). Elsevier Press, Amsterdam. p. 131-158.
- Justine, J. L. 2010. Parasites of coral reef fish: how much do we know? With a bibliography of fish parasites in New Caledonia. *Belgian Journal of Zoology* 140 (Suppl.):155-190.
- Linton, E. 1910. Helminth Fauna of the Dry Tortugas. II Trematodes. Publication of the Carnegie Institute of Washington 133:11-98.
- Manter, H. W. 1947. The digenetic trematodes of marine fishes of Tortugas, Florida. *The American Midland Naturalist* 38:257-426.
- Nahhas, F. M. and K. Carlson. 1994. Digenetic trematodes of marine fishes of Jamaica, West Indies. Department of Biological Sciences, University of the Pacific, Stockton, CA, USA. Publication 2. 60 p.
- Overstreet, R. M. 1969. Digenetic trematodes of marine teleost fishes from Biscayne Bay, Florida. *Tulane Studies in Zoology and Botanic* 15:119-176.
- Sogandares-Bernal, F. 1959. Digenetic trematodes of marine fishes from the Gulf of Panama and Bimini, British West Indies. *Tulane Studies in Zoology* 7:70-117.
- Sogandares-Bernal, F. and L. McAlister-Sogandares. 1961. Nine digenetic trematodes of marine fishes from the Atlantic coast of Panama. *Tulane Studies in Zoology* 8:141-153.
- Vélez, E. I. 1987. Sobre la fauna de tremátodos en peces marinos de la familia Lutjanidae en el Mar Caribe. *Actualidades Biológicas* 16:70-84.
- Vidal-Martínez, V. M., M. L. Aguirre-Macedo, T. Scholz, D. González-Solís and E. Mendoza-Franco. 2001. Atlas of the helminth parasites of cichlid fish of Mexico. Academia, Praga. 185 p.
- Zhukov, E. V. 1976. New monogenean species of the genus *Haliotrema* Johnston and Tiegs, 1922, from the Gulf of Mexico fishes of the Fam. Lutjanidae, in *Fauna, systematics and phylogeny of Monogenoidea*. Proceedings, Institute of Biology and Pedology, Far-East Science Centre, Academy of Sciences of the U. S. S. R., New Series 35:33-47 (in Russian).