



Revista Mexicana de Biodiversidad

ISSN: 1870-3453

[falvarez@ib.unam.mx](mailto:falvarez@ib.unam.mx)

Universidad Nacional Autónoma de  
México  
México

Gallas, Moisés; Calegaro-Marques, Cláudia; Bencke-Amato, Suzana  
Supplemental observations on the morphology of *Procamallanus* (*Spirocamallanus*) *hilarii*  
(Nematoda: Camallanidae) parasitizing two species of *Astyanax* (Characiformes:  
Characidae) and ecological analyses  
Revista Mexicana de Biodiversidad, vol. 86, núm. 3, septiembre, 2015, pp. 590-596  
Universidad Nacional Autónoma de México  
Distrito Federal, México

Available in: <http://www.redalyc.org/articulo.oa?id=42542746006>

- How to cite
- Complete issue
- More information about this article
- Journal's homepage in [redalyc.org](http://redalyc.org)

[redalyc.org](http://redalyc.org)

Scientific Information System

Network of Scientific Journals from Latin America, the Caribbean, Spain and Portugal

Non-profit academic project, developed under the open access initiative



## Taxonomy and systematics

# Supplemental observations on the morphology of *Procamallanus* (*Spirocamallanus*) *hilarii* (Nematoda: Camallanidae) parasitizing two species of *Astyanax* (Characiformes: Characidae) and ecological analyses

## *Observaciones suplementarias sobre la morfología de Procamallanus (Spirocamallanus) hilarii (Nematoda: Camallanidae) parasitando dos especies de Astyanax (Characiformes: Characidae) y análisis ecológico*

Moisés Gallas\*, Cláudia Calegare-Marques, Suzana Bencke-Amato

Laboratório de Helmintologia, Departamento de Zoologia, Instituto de Biociências, Universidade Federal do Rio Grande do Sul, 91501-970 Porto Alegre, Rio Grande do Sul, Brazil

Received 8 September 2014; accepted 13 May 2015

Available online 1 September 2015

### Abstract

*Procamallanus* (*Spirocamallanus*) *hilarii* parasitizes several fish species but only few studies on morphology and ecology have been published. In this study we describe the morphology of *P. (S.) hilarii* parasitizing *Astyanax* aff. *fasciatus* and *Astyanax jacuhiensis*, and compare ecological parameters of their populations. Seventy *A. aff. fasciatus* and 60 *A. jacuhiensis* individuals were captured in Lake Guaíba, Rio Grande do Sul State, Brazil, between March 2012 and December 2013. The specimens of *P. (S.) hilarii* presented an oval oral opening with 6 cuticular elevations and 2 groups of papillae (one internal and one external) in the anterior region of the body. Males have 8 pairs of sessile caudal papillae (4 pre-cloacal and 4 post-cloacal), spicules with distal bifurcation and lack caudal alae. The prevalence of *P. (S.) hilarii* was higher in *A. aff. fasciatus* (69%) than in *A. jacuhiensis* (18%), whereas there was no difference in mean intensity (respectively, 2 and 5) and mean abundance (respectively, 1.6 and 0.9) of infection between the 2 host species. We suggest that the consumption of copepods, intermediate hosts of *P. (S.) hilarii*, may explain this difference.

All Rights Reserved © 2015 Universidad Nacional Autónoma de México, Instituto de Biología. This is an open access item distributed under the Creative Commons CC License BY-NC-ND 4.0.

**Keywords:** Spiruridae; Taxonomy; Morphology; Characids; Neotropical region

### Resumen

*Procamallanus* (*Spirocamallanus*) *hilarii* (*P. (S.) hilarii*) parasita varias especies de peces, pero son escasos los estudios publicados sobre su morfología y ecología. En este estudio describimos la morfología de *P. (S.) hilarii* que parasitan las mojaras *Astyanax* aff. *fasciatus* y *A. jacuhiensis* y comparamos diferentes parámetros ecológicos de sus poblaciones. Entre marzo de 2012 y diciembre de 2013, capturamos 70 individuos de *Astyanax* (*A.*) aff. *fasciatus* y 60 de *A. jacuhiensis* en el Lago Guaíba, estado de Rio Grande do Sul, Brazil. Los ejemplares de *P. (S.) hilarii* presentaron una abertura oral de forma oval con 6 elevaciones cuticulares y 2 grupos de papilas (una interna y otra externa) en la región anterior del cuerpo. Los machos presentaron 8 pares de papilas caudales sésiles (4 precloaca y 4 poscloaca), espículas con bifurcación distal y ausencia de ala caudal. La prevalencia de *P. (S.) hilarii* fue mayor en *A. aff. fasciatus* (69%) que en *A. jacuhiensis* (18%). Sin embargo, no existieron diferencias en la intensidad promedio (2 y 5 respectivamente) y la abundancia promedio (1.6 y 0.9 respectivamente) de infección entre las 2 especies hospederas. Sugerimos que el consumo de copépodos, hospederos intermediarios de *P. (S.) hilarii*, podría explicar esta diferencia.

Derechos Reservados © 2015 Universidad Nacional Autónoma de México, Instituto de Biología. Este es un artículo de acceso abierto distribuido bajo los términos de la Licencia Creative Commons CC BY-NC-ND 4.0.

**Palabras clave:** Spiruridae; Taxonomía; Morfología; Carácidos; Región neotropical

\* Corresponding author.

E-mail address: [mgallas88@gmail.com](mailto:mgallas88@gmail.com) (M. Gallas).

Peer Review under the responsibility of Universidad Nacional Autónoma de México.

## Introduction

*Procamallanus* Baylis, 1923, a genus of nematode parasites of freshwater and marine fishes (Moravec, Salgado-Maldonado, & Caspeta-Mandujano, 2000), was originally proposed to include *Cucullanus laeviconchus* Wedl, 1862, a parasite of freshwater siluriform fishes based on the morphology of the buccal capsule (Baylis, 1923). Olsen (1952) proposed *Spirocamallanus* for grouping the species that had spiral thickenings in the buccal capsule, whereas Moravec and Sey (1988) considered it as a subgenus of *Procamallanus* based on the presence of buccal capsule with inner spirals in males and females. Gibbons (2010) has recently considered the latter proposition valid and grouped 6 additional subgenera within *Procamallanus*.

Neotropical species of *Procamallanus* belong to the subgenera *Procamallanus* Baylis, 1923, *Spirocamallanus*, and *Denticamallanus* Moravec and Thatcher, 1997 (Luque, Aguiar, Vieira, Gibson, & Santos, 2011; Moravec, 1998). Luque et al. (2011) listed all 22 species of *Procamallanus* so far recorded in Brazil. Vaz and Pereira (1934) reported *Procamallanus hilarii* Vaz and Pereira, 1994, parasitizing the small intestine of *Salminus hilarii* Valenciennes, 1849 from the Santo Amaro Dam, São Paulo State, Brazil. Later, this was classified in the subgenus *Spirocamallanus* by presenting oral capsule with inner spirals in both sexes (Moravec & Sey, 1988). This species was redescribed by Pinto and Noronha (1976) and Rodrigues, Pinto, and Noronha (1991); however, in both studies the authors used the original figures of Vaz and Pereira (1934), thereby missing many morphological details, such as cephalic characters and posterior extremity of males.

In Brazil, *P. (S.) hilarii* was recorded in the states of Ceará in *Astyanax bimaculatus* (Linnaeus, 1758) and *Steindachnerina elegans* (Steindachner, 1874) (Pereira, Dias, & Azevedo, 1936), Rio de Janeiro in *A. bimaculatus* and *Astyanax paraguayae* (Eigenmann, 1908) (Abdallah, Azevedo, & Luque, 2004), São Paulo in *A. bimaculatus* (Kloss, 1966; Kohn & Fernandes, 1987) and *Salminus hilarii* Valenciennes, 1850 (Vaz & Pereira, 1934), as well as unidentified minnows (Pinto & Noronha, 1976), Paraná in *Astyanax fasciatus* (Cuvier, 1819) (Kohn, Fernandes, Pipolo, & Godoy, 1988; Pinto & Noronha, 1976) and unidentified minnows (Pinto & Noronha, 1976), and Rio Grande do Sul in *Acestrorhynchus microlepis* (Schomburgk, 1841), *Astyanax jacuhiensis*, *Astyanax* aff. *fasciatus*, *Hoplias lacerdae* Miranda Ribeiro, 1908, *Hoplias malabaricus*, and *Rhamdia quelen* (Quoy and Gaimard, 1824) (Kohn, Fernandes, Pipolo, & Godoy, 1989). This nematode also was recorded in *Trichomycterus piuræ* (Eigenmann, 1922) in Peru (Freitas & Ibañez, 1970) and in *Astyanax abramis*, *A. bimaculatus*, *A. fasciatus*, *Hoplias malabaricus*, *Leporinus obtusidens* (Valenciennes, 1836), *Oligosarcus jenynsii* (Günther, 1864), *Pimelodus albicans* (Valenciennes, 1840), and *Salminus brasiliensis* (Cuvier, 1816) in Argentina (Ramalho, 1997, 2005). Luque et al. (2011) also cited the characid *Oligosarcus macrolepis* Steindachner, 1877 as a host of this species, but the collection site was not reported. While Lima et al. (2003) and Buckup, Menezes, and Ghazzi (2007) report that *O. macrolepis* occurs in the Jequitinhonha river basin in Bahia and Minas Gerais, these states

were not surveyed by Luque et al. (2011). Their collections were restricted to the states of Ceará, Rio de Janeiro, São Paulo, Paraná, and Rio Grande do Sul.

*Procamallanus (S.) hilarii* was recorded in *A. aff. fasciatus* and *A. jacuhiensis* by Kohn et al. (1989), but a comparison of its ecological parameters between these host species is missing. The present study aims to fulfill this gap by providing a detailed description of the morphology of *P. (S.) hilarii* from Lake Guaíba, Rio Grande do Sul State, Brazil, and discussing their ecological parameters in light of their trophic relations.

## Materials and methods


Seventy specimens of *A. aff. fasciatus* and 60 of *A. jacuhiensis* were collected with seine nets between March 2012 and December 2013 from Lake Guaíba, in the municipalities of Guaíba (30°08'28"S, 51°18'53"W) and Barra do Ribeiro (30°17'11"S, 51°18'01"W), State of Rio Grande do Sul, Brazil. Fishes were individually packed in plastic sacks, and transported in ice coolers to the Laboratório de Helminologia of the Departamento de Zoologia (DZ), Universidade Federal do Rio Grande do Sul (UFRGS) for necropsy. Hosts were determined according to Lucena, Castro, and Bertaco (2013). Nematodes were collected, fixed, and preserved following Amato and Amato (2010). These were temporarily mounted and cleared in Amann's lactophenol (Humason, 1979). *En face* preparations were mounted in glycerin jelly according to Anderson (1958) to visualize cephalic characters. Specimens were prepared for scanning electron microscopy (SEM) through dehydration in ethanol (70° to 100° GL) and the ethanol was gradually replaced by acetone (10–100%) to critical point dried. Then, they were mounted on stubs, coated with carbon and gold, and examined in a JEOL JSM 6060 at Centro de Microscopia Eletrônica (CME) at UFRGS.

Measurements are shown in micrometers (µm) unless otherwise stated. The parameters reported are the range followed by mean ± standard deviation, and sample size is between parentheses. Photomicrographs were taken with a Zeiss Axiolab microscope with phase contrast. Line drawings were made with a drawing tube mounted on a Nikon E-200 microscope, scanned and prepared using CorelDraw X4® and Adobe's Photoshop® CS2. Ecological parameters follow Bush, Lafferty, Lotz, and Shostak (1997). Voucher helminth specimens were deposited in the 'Coleção Helminológica do Instituto Oswaldo Cruz' (CHIOC), Rio de Janeiro, RJ, Brazil. Voucher host specimens were deposited in the 'Coleção Ictiológica', DZ/UFRGS. The Z-test was used to compare prevalence data between the host species, whereas median intensity and median abundance of infection were compared by the Student *t* test (Zar, 2010).

## Description

*Procamallanus (Spirocamallanus) hilarii* Vaz and Pereira, 1934 (Figs. 1–18)

*Diagnosis* (based on 15 males and 15 females mounted in lactophenol, 2 *en face*, and 5 males prepared for scanning electron microscopy). Spiruroidea, Camallanidae.




Figures 1–6. Incomplete diagrams of *Procamallanus* (*Spirocamallanus*) *hilarii*. (1), Anterior extremity of males. Scale bar = 150  $\mu$ m; (2), en face view. Scale bar = 50  $\mu$ m; (3), posterior extremity of males. Scale bar = 50  $\mu$ m; (4), detail of the ventral view of the spicules. Scale bar = 50  $\mu$ m; (5), vulvar region of females. Scale bar = 100  $\mu$ m; (6), posterior extremity of a young female. Scale bar = 150  $\mu$ m.

Reddish nematodes in life, with transverse cuticular striations. Oval mouth opening surrounded by 6 elevations (1 ventral, 1 dorsal, and 4 lateral), and 8 papillae distributed in 2 circles (4 internal smaller and 4 external larger) (Figs. 2, 8, 9 and 16). Yellowish brown buccal capsule, sclerotized, with 13–16 internal spirals, continuous, some incomplete (Figs. 1, 7, 15 and 16). Basal ring sclerotized between buccal capsule and muscular esophagus (Figs. 1 and 7). Esophagus divided into a shorter muscular and a longer glandular portion (Figs. 1 and 7). Nerve ring anterior to deirids and excretory pore (Figs. 1, 7, 15 and 17). Males ( $N=15$ ). Body 6.34–9.07 mm ( $7.58 \pm 0.89$  mm;  $N=15$ ) long, 0.16–0.22 mm ( $0.18 \pm 0.02$  mm;  $N=15$ ) wide. Buccal capsule 45.00–70.00 ( $53.17 \pm 5.71$ ;

$N=15$ ) long, 52.50–65.00 ( $56.67 \pm 3.74$ ;  $N=15$ ) wide, with 13–16 internal spirals. Basal ring 7.50–12.50 ( $9.67 \pm 1.30$ ;  $N=15$ ) long, 20.00–40.00 ( $32.50 \pm 4.82$ ;  $N=15$ ) wide. Muscular esophagus 340.00–460.00 ( $390.00 \pm 34.11$ ;  $N=15$ ) long, 50–70 ( $59.67 \pm 5.50$ ;  $N=15$ ) and 100–130 ( $113.33 \pm 10.47$ ;  $N=15$ ) wide in the anterior and posterior ends, respectively; glandular esophagus 1.04–1.77 mm ( $1.40 \pm 0.20$  mm;  $N=15$ ) long, 0.09–0.14 mm ( $0.11 \pm 0.01$  mm;  $N=15$ ) wide. Esophagus representing 24% of total body length (TBL), and muscular portion corresponding to 28% of the glandular portion. Nerve ring 162.50–222.50 ( $195.83 \pm 16.63$ ;  $N=15$ ) from anterior extremity. Deirids 185.00–282.50 ( $235.00 \pm 24.68$ ;  $N=12$ ) from anterior extremity. Excretory pore 215.00–352.50







Figures 7–11. Photomicrographies of males of *Procamallanus (Spirocamallanus) hilarii*. (7), Anterior extremity in phase contrast. Scale bar = 100  $\mu\text{m}$ ; (8), *en face* view showing the 6 cuticular elevations (white arrow heads). Scale bar = 50  $\mu\text{m}$ ; (9), *en face* view showing the 4 outer papillae (white arrow heads), amphids (black arrow heads) and the end of the buccal capsule (asterisk). Scale bar = 50  $\mu\text{m}$ ; (10), posterior extremity showing the spicules (black arrow heads). Scale bar = 50  $\mu\text{m}$ ; (11), detail of the ventral view of the spicules showing the distal bifurcation (white arrow heads). Scale bar = 50  $\mu\text{m}$ .

( $279.17 \pm 36.37$ ;  $N = 15$ ) from anterior extremity. Posterior end of the body with 8 pairs of papillae: 4 pre- and 4 post-cloacal (Figs. 3, 10 and 18). Similar spicules, bifurcated distal end, with length of 52.50–77.50 ( $64.83 \pm 5.94$ ;  $N = 15$ ) and 62.5–87.5

( $72.83 \pm 6.54$ ;  $N = 15$ ) (Figs. 3, 4, 10 and 11). Gubernaculum absent. Cloaca 120.00–200.00 ( $162.83 \pm 25.03$ ;  $n = 15$ ) from the posterior extremity, pointed tail (Figs. 3, 10 and 18). Females ( $N = 15$ ). Body 12.92–18.78 mm ( $15.69 \pm 1.93$  mm;



Figures 12–14. Photomicrographies of females of *Procamallanus (Spirocamallanus) hilarii*. (12), Vulvar region with larvae L1 (white arrow heads). Scale bar = 250  $\mu\text{m}$ ; (13), posterior extremity of a gravid female showing larvae L1 (white arrow heads). Scale bar = 100  $\mu\text{m}$ ; (14), larvae stained with rose bengal. Scale bar = 100  $\mu\text{m}$ .



Figures 15–18. Scanning electron microscopy of *Procamallanus (Spirocamallanus) hilarii*. (15), Anterior extremity with deirid (black arrow head) and excretory pore (white arrow head). Scale bar = 50  $\mu$ m; (16), detail of the anterior extremity showing cuticular elevations (asterisks), inner papillae (black arrow heads), outer papillae (white arrow heads) and amphids (black arrows). Scale bar = 10  $\mu$ m; (17), detail of the deirid. Scale bar = 10  $\mu$ m; (18), posterior extremity of a male showing the phasmids (white head arrows). Scale bar = 20  $\mu$ m.

$N=15$ ) long, 0.22–0.44 mm ( $0.34 \pm 0.06$  mm;  $N=15$ ) wide. Buccal capsule 55.00–65.00 ( $60.00 \pm 3.13$ ;  $N=15$ ) long, 60.00–72.50 ( $66.67 \pm 3.62$ ;  $N=15$ ) wide, with 13–16 internal spirals. Basal ring 7.50–12.50 ( $10.67 \pm 1.48$ ;  $N=15$ ) long, 35.00–45.00 ( $39.67 \pm 2.81$ ;  $N=15$ ) wide. Muscular esophagus 370.00–520.00 ( $434.00 \pm 36.41$ ;  $N=15$ ) long, 60.00–80.00 ( $69.33 \pm 4.58$ ;  $n=15$ ) and 120.00–150.00 ( $128.67 \pm 9.9$ ;  $n=15$ ) wide in the anterior and posterior ends, respectively; glandular esophagus 1.15–2.22 mm ( $1.77 \pm 0.26$  mm;  $N=15$ ) long, 0.12–0.17 mm ( $0.14 \pm 0.01$  mm;  $N=15$ ) wide. Esophagus representing 11% of TBL, and muscular portion corresponding to 25% of the glandular portion. Nerve ring 182.50–232.50 ( $207.17 \pm 14.48$ ;  $N=15$ ) from anterior extremity. Deirids 250.00–350.00 ( $285.77 \pm 34.86$ ;  $N=13$ ) from anterior extremity. Excretory pore 280.00–410.00 ( $341.04 \pm 47.07$ ;  $N=12$ ) from anterior extremity. Vulva post-equatorial, 5.48–7.94 mm ( $6.82 \pm 0.76$  mm;  $N=15$ ) from posterior extremity (Figs. 5 and 12). Eggs 7.50–10.00 ( $9.00 \pm 1.29$ ) long and 5.00 to 10.00 ( $7.50 \pm 1.67$ ) wide ( $N=10$ ), with first larval stage 390.00–529.50 ( $514.97 \pm 60.38$ ;

$N=15$ ) long and 25.00–32.50 ( $27.00 \pm 3.02$ ;  $N=15$ ) wide (Figs. 12, 13 and 14). Anus 80.00–140.00 ( $111.33 \pm 16.42$ ;  $N=15$ ) from posterior extremity, pointed tail (Figs. 6 and 13).

#### Taxonomic summary

**Synonyms:** *Procamallanus cearensis* Pereira, Dias and Azevedo, 1936; *Spirocamallanus incarocai* Freitas & Ibañez, 1970.

**Hosts:** *A. aff. fasciatus* and *A. jacuhiensis*.

**Host specimens deposited.** *A. aff. fasciatus*: UFRGS 19121 (male); UFRGS 19122 (female) and *A. jacuhiensis*: UFRGS 19123 (male); UFRGS 19124 (female).

**Localities:** Lake Guaíba, Municipalities of Guaíba and Barra do Ribeiro, State of Rio Grande do Sul, Brazil.

**Site of infection:** stomach and intestine.

**Voucher specimens deposited:** CHIOC n° 35948.

**Ecological parameters:** the prevalence of *P. (S.) hilarii* in *A. aff. fasciatus* (69%) was higher than in *A. jacuhiensis* (18%) ( $Z=5.7355$ ,  $p_{\text{two-tailed}} < 0.0001$ ; Table 1). However, the mean intensity (*A. aff. fasciatus* = 2 helminths/host vs. *A.*



Table 1

Ecological parameters of *Procamallanus* (*Spirocamallanus*) *hilarii* populations and feeding habits of host species.

Host	N host	Diet	Prevalence (%)	Mean intensity (helminths/host)	Mean abundance (helminths/host)	Reference
<i>Astyanax abramis</i>	228	Juveniles: vegetal matter and insects. Adults: microcrustaceans, crustaceans and insects <sup>a</sup>	52	3	–	Ramalho (2005)
<i>Astyanax bimaculatus</i>	124	Vegetal matter, insects and other animals (spiders and fish rests) <sup>b</sup>	45	3	–	Ramalho (2005)
	37		40	2	–	Ramalho (2005)
	23		4	–	–	Kohn and Fernandes (1987)
	40		3	1	<0.1	Abdallah et al. (2004)
<i>Astyanax fasciatus</i>	412	Vegetal matter, insects and some microcrustaceans <sup>c</sup>	56	2	–	Ramalho (2005)
	16		6	1	0.1	Kohn et al. (1988)
<i>Astyanax</i> aff. <i>fasciatus</i>	70	Vegetal matter, insects and some microcrustaceans <sup>c</sup>	69	2	1.6	Present study
	8		25	1	0.3	Kohn et al. (1989)
<i>Astyanax jacuhiensis</i>	17	Vegetal matter and insects <sup>c</sup>	65	2	1.2	Kohn et al. (1989)
	60		18	5	0.9	Present study
<i>Astyanax paraguayae</i>	40	*	5	1	0.1	Abdallah et al. (2004)
<i>Acestrorhynchus macrolepis</i>	11	*	45	3	1.4	Kohn et al. (1989)
<i>Hoplias lacerdae</i>	3	Adults: fish <sup>d</sup>	33	1	0.3	Kohn et al. (1989)
<i>Hoplias malabaricus</i>	6	Juveniles: insects and crustaceans. Adults: fish and shrimp <sup>e</sup>	17	2	0.2	Kohn et al. (1989)
<i>Rhamdia quelen</i>	8	Fish, molluscs, crustaceans and insects <sup>e</sup>	12	2	0.1	Kohn et al. (1989)

<sup>a</sup> Barros (2004).<sup>b</sup> Andrian, Silva, and Peretti (2001).<sup>c</sup> Vilella et al. (2002).<sup>d</sup> Alvim and Peret (2004).<sup>e</sup> Malabarba et al. (2013).

\* Unknown data.

*jacuhiensis* = 5 helminths/host;  $t = -0.87$ ,  $df = 10$ ,  $p = 0.4$ ) and the mean abundance (*A. aff. fasciatus* = 1.6 helminths/host vs. *A. jacuhiensis* = 0.9 helminths/host;  $t = 1.24$ ,  $df = 70$ ,  $p = 0.22$ ) of infection were similar between species (Table 1).

## Remarks

The nematodes found in the stomach and in the anterior and middle portions of the small intestine of *A. aff. fasciatus* and *A. jacuhiensis* were identified as *P. (S.) hilarii* based on the taxonomic keys proposed by Rodrigues et al. (1991) and Moravec (1998) by presenting an oral capsule with inner spirals in both sexes and a glandular esophagus at least 3 times longer than the muscular esophagus.

Species of *Procamallanus* have 6 elevations distributed over the anterior region of the buccal capsule, 8 papillae arranged in 2 circles, and a pair of lateral amphids (Moravec, 1998). Unlike the round oral opening reported for *P. (S.) hilarii* by Moravec (1998), the specimens examined presented an oval oral opening. Ramalho (1997) cited the presence of 3 pairs of small cephalic papillae, 1 lateral and 2 median. In fact, these papillae are the cuticular elevations described by Moravec (1998) and observed in the present study (Figs. 2, 8, 9 and 16).

Morphological differences also were found in the posterior end of males. In species considered synonymous of *P. (S.) hilarii*, *P. cearensis* and *S. incarocai*, Pereira et al. (1936)

and Freitas and Ibañez (1970) mentioned the presence of thin caudal alae, whereas Rodrigues et al. (1991) and Moravec (1998) noted the absence of caudal alae as reported in the present study (Figs. 3, 10 and 18). Freitas and Ibañez (1970) and Ramalho (1997) described 8 pairs of pedunculate papillae, unlike the sessile papillae reported here (Figs. 3, 10 and 18) and in Pereira et al. (1936), Rodrigues et al. (1991), and Moravec (1998). Additionally, although the spicules of the specimens examined in the present study are similar to those described by Pereira et al. (1936), these authors did not report the bifurcation of its distal end. In general, the dimensions of characters were similar to those reported in the literature (Freitas & Ibañez, 1970; Pereira et al., 1936; Pinto & Noronha, 1976; Ramalho, 1997; Rodrigues et al., 1991; Vaz & Pereira, 1934). Male and female body size was an exception. The total length of males and females reported in this study was higher. Only Freitas and Ibañez (1970) reported larger females. These authors also reported that the vulva is closer (2.4 mm) to the posterior end of the worm. Morphological differences may result from intraspecific variations in length, for example, or from difficulties in visualizing structures, as was the case for the caudal papillae and caudal alae. However, the MEV images clarify these discrepancies.

Ecological parameters of *P. (S.) hilarii* populations vary among host species and study (Table 1). For instance, prevalence ranges from 3% in *A. bimaculatus* to 69% in *A. aff. fasciatus*.

These differences suggest that the definitive hosts exploit a variable diet in different environments. In the case of the present study, the differences in prevalence of *P. (S.) hilarii* between the 2 characid hosts likely reflect the differences in their feeding habits. Since the life cycle of *Procamallanus* spp. includes copepods as intermediate hosts (Anderson, 2000), it is possible to suggest that these small crustaceans are more heavily preyed upon by *A. aff. fasciatus* than by *A. jacuhiensis*. It is also interesting that the intensity of infection showed the opposite pattern, being higher in *A. jacuhiensis*, a species with a more herbivorous diet (Villela, Becker, & Hartz, 2002). Together, the data on prevalence and intensity of infection may suggest that the fewer *A. jacuhiensis* individuals that feed on copepods do that at a higher frequency than most *A. aff. fasciatus*. In addition, variations in environmental characteristics can be related to the availability in the environment of infected copepods.

## Acknowledgements

We thank Dra. Ana P. S. Dufech for field assistance, Dr. Philip J. Scholl for reviewing the English, and Centro de Microscopia Eletrônica (UFRGS) for SEM preparations. MG also thanks the Brazilian National Research Council (CNPq) for financial support (process #131068/2012-1) and CCM thanks the Brazilian Higher Education Authority (CAPES) and the Research Foundation of the State of Rio Grande do Sul (FAPERGS) for post-doctoral fellowships (PNPD/CAPES 2010).

## References

- Abdallah, V. D., Azevedo, R. K., & Luque, J. L. (2004). Metazoários parasitos dos lambaris *Astyanax bimaculatus* (Linnaeus, 1758), *A. paraguayensis* Eigenmann, 1908 e *Oligosarcus hepsetus* (Cuvier, 1829) (Osteichthyes: Characidae), do Rio Guandu, Estado do Rio de Janeiro, Brasil. *Revista Brasileira de Parasitologia Veterinária*, 13, 57–63.
- Alvim, M. C., & Peret, A. C. (2004). Food resources sustaining the fish fauna in a section of the Upper São Francisco River in Três Marias, MG, Brazil. *Brazilian Journal of Biology*, 64, 195–202.
- Amato, J. F. R., & Amato, S. B. (2010). Técnicas gerais para coleta e preparação de helmintos endoparasitos de aves. In S. Von Matter, F. C. Straube, I. A. Accordi, V. Q. Piacentini, & J. F. Cândido Jr. (Eds.), *Ornitologia e conservação: ciência aplicada, técnicas de pesquisa e levantamento* (pp. 369–393). Rio de Janeiro: Technical Books.
- Anderson, R. C. (1958). Méthode pour l'examen des Nématodes en vue apicale. *Annales de Parasitologie Humaine et Comparée*, 33, 171–172.
- Anderson, R. C. (2000). *Nematode parasites of vertebrates: their development and transmission*. Wallingford: CABI Publishing.
- Andrian, I. F., Silva, H. B. R., & Peretti, D. (2001). Dieta de *Astyanax bimaculatus* (Linnaeus, 1758) (Characiformes, Characidae), da área de influência do reservatório de Corumbá, Estado de Goiás, Brasil. *Acta Scientiarum, Biological Sciences*, 23, 435–440.
- Barros, S. E. (2004). Alimentación de *Astyanax abramis* (Characiformes: Characidae) en el Embalse Cabra Corral, Salta, Noroeste de Argentina. *Revista AquaTIC*, 20, 88–96.
- Baylis, H. A. (1923). Report on a collection of parasitic nematodes, mainly from Egypt. Part III. Camallanidae, etc. With a note on *Probstmatia* and an appendix on Acanthocephala. *Parasitology*, 15, 24–38.
- Bush, A. O., Lafferty, K. D., Lotz, J. M., & Shostak, A. W. (1997). Parasitology meets ecology on its own terms: Margolis et al. revisited. *The Journal of Parasitology*, 83, 575–583.
- Buckup, P. A., Menezes, N. A., & Ghazzi, M. S. (2007). *Catálogo das espécies de peixes de água doce do Brasil*. Rio de Janeiro: Museu Nacional.
- Freitas, J. F. T., & Ibañez, H. N. (1970). Fauna helmintológica del Peru. Nuevo nematodo camalanideo parasito de pez. *Archivos Peruanos de Patología y Clínica*, 24, 205–210.
- Gibbons, L. M. (2010). *Keys to the nematode parasites of vertebrates* (Vol. supplementary volume). Wallingford: CABI Publishing.
- Humason, G. L. (1979). *Animal tissue techniques*. San Francisco: W. H. Freeman and Company.
- Kloss, G. R. (1966). Helmintos parasitos de espécies simpátricas de *Astyanax* (Pisces, aracidæ). *Papéis Avulsos do Departamento de Zoologia de São Paulo*, 18, 189–219.
- Kohn, A., & Fernandes, B. M. M. (1987). Estudo comparativo dos helmintos parasitos de peixes do Rio Mogi Guassu, coletados nas excursões realizadas entre 1927 e 1985. *Memórias do Instituto Oswaldo Cruz*, 82, 483–500.
- Kohn, A., Fernandes, B. M. M., Pipolo, H. V., & Godoy, M. P. (1988). Helmintos parasitos de peixes das Usinas Hidrelétricas da Eletrosul (Brasil). II. Reservatórios de Salto Osório e de Salto Santiago, Bacia do Rio Iguaçu. *Memórias do Instituto Oswaldo Cruz*, 83, 299–303.
- Kohn, A., Fernandes, B. M. M., Pipolo, H. V., & Godoy, M. P. (1989). List of helminth parasites of fishes from the Passo Fundo reservoir, Uruguay River Basin, Brasil. *Memórias do Instituto Oswaldo Cruz*, 84, 427–428.
- Lima, F. C. T., Malabarba, L. R., Buckup, P. A., Silva, J. F. P., Vari, R. P., Harold, A., et al. (2003). *Genera incertae sedis in Characidae*. In R. E. Reis, S. O. Kullander, & C. J. Ferraris Jr. (Eds.), *Check list of the freshwater fishes of South and Central America* (pp. 106–169). Porto Alegre: EDIPUCRS.
- Lucena, C. A. S., Castro, J. B., & Bertaco, V. A. (2013). Three new species of *Astyanax* from drainages of southern Brazil (Characiformes: Characidae). *Neotropical Ichthyology*, 11, 537–552.
- Luque, J. L., Aguiar, J. C., Vieira, F. M., Gibson, D. I., & Santos, C. P. (2011). Checklist of Nematoda associated with the fishes of Brazil. *Zootaxa*, 3082, 1–88.
- Malabarba, L. R., Carvalho-Neto, P., Betaco, V. A., Carvalho, T. P., Santos, J. F., & Artioli, L. G. S. (2013). *Guia de identificação dos peixes da Bacia do Rio Tramandá*. Porto Alegre: Via Sapiens.
- Moravec, F. (1998). *Nematodes of freshwater fishes of the neotropical region*. Praha: Academia.
- Moravec, F., & Sey, O. (1988). Nematodes of freshwater fishes from North Vietnam. Part I. Camallanoidea and Habronematoidea. *Vestník Československé Společnosti Zoologické*, 52, 128–148.
- Moravec, F., Salgado-Maldonado, G., & Caspeta-Mandujano, J. (2000). Three new *Procamallanus* (*Spirocamallanus*) species from freshwater fishes in Mexico. *The Journal of Parasitology*, 86, 119–127.
- Olsen, L. S. (1952). Some nematodes parasitic in marine fishes. *Publications of Institute of Marine Sciences*, 2, 173–212.
- Pereira, C., Dias, M. M., & Azevedo, P. (1936). Biologia do nematóide *Procamallanus cearensis* n. sp. *Arquivos do Instituto Biológico*, 7, 209–226.
- Pinto, R. M., & Noronha, D. (1976). *Procamallanus* brasileiros (Nematoda, mal-lanoidea): considerações finais, com chave para determinação das espécies. *Memórias do Instituto Oswaldo Cruz*, 74, 323–339.
- Ramallo, G. (1997). *Spirocamallanus hilarii* (Nematoda, Camallanidae) parasito de peces dulceacuícolas del embalse de Termas de Río Hondo, Santiago del Estero, Argentina. *Boletín Chileno de Parasitología*, 52, 67–70.
- Ramallo, G. (2005). *Procamallanus* (*Spirocamallanus*) *hilarii* Vaz and Pereira, 34 (Nematoda, Camallanidae), parasito de peces dulceacuícolas del norte de Argentina. *Acta Zoológica Lilloana*, 49, 137–139.
- Rodrigues, H. O., Pinto, R. M., & Noronha, D. (1991). Key to the species of Brazilian *Procamallanus* with general considerations (Nematoda, mal-lanoidea). *Memórias do Instituto Oswaldo Cruz*, 86, 107–113.
- Vaz, Z., & Pereira, C. (1934). Contribuição ao conhecimento dos nematóides de peixes fluviais do Brasil. *Arquivos do Instituto Biológico*, 5, 87–103.
- Vilella, F. S., Becker, F. G., & Hartz, S. M. (2002). Diet of *Astyanax* species (Teleostei, aracidæ) in an Atlantic Forest River in Southern Brazil. *Brazilian Archives of Biology and Technology*, 45, 223–232.
- Zar, J. H. (2010). *Biostatistical analysis*. New Jersey: Prentice Hall.