



Revista Mexicana de Biodiversidad

ISSN: 1870-3453

[falvarez@ib.unam.mx](mailto:falvarez@ib.unam.mx)

Universidad Nacional Autónoma de  
México  
México

Zavala-Páramo, Alfonso Sinuhé; Oyama, Ken; Mendoza, Eduardo; Zavala-Páramo, María  
Guadalupe; Pollinger, John; Smith, Thomas B.

Genetic variability in captive individuals of the endangered species *Tapirus bairdii* in  
Mexico

Revista Mexicana de Biodiversidad, vol. 88, núm. 2, junio-, 2017, pp. 480-484

Universidad Nacional Autónoma de México  
Distrito Federal, México

Available in: <http://www.redalyc.org/articulo.oa?id=42551127029>

- How to cite
- Complete issue
- More information about this article
- Journal's homepage in [redalyc.org](http://redalyc.org)

[redalyc.org](http://redalyc.org)

Scientific Information System

Network of Scientific Journals from Latin America, the Caribbean, Spain and Portugal

Non-profit academic project, developed under the open access initiative



## Research note

# Genetic variability in captive individuals of the endangered species *Tapirus bairdii* in Mexico

## *Variabilidad genética en individuos en cautiverio en México de la especie en peligro de extinción Tapirus bairdii*

Alfonso Sinuhé Zavala-Páramo<sup>a</sup>, Ken Oyama<sup>a</sup>, Eduardo Mendoza<sup>b,\*</sup>,  
María Guadalupe Zavala-Páramo<sup>c</sup>, John Pollinger<sup>d</sup>, Thomas B. Smith<sup>d</sup>

<sup>a</sup> Escuela Nacional de Estudios Superiores, Universidad Nacional Autónoma de México, Antigua carretera a Pátzcuaro Núm. 8701, Expropiación Petrolera INDECO, Col. Ex Hacienda San José de la Huerta, 58190 Morelia, Michoacán, Mexico

<sup>b</sup> Instituto de Investigaciones sobre los Recursos Naturales, Universidad Michoacana de San Nicolás de Hidalgo, Av. San Juanito Itzicuaró s/n, Col. Nueva Esperanza, 58330 Morelia, Michoacán, Mexico

<sup>c</sup> Centro Multidisciplinario de Estudios en Biotecnología, FMVZ, Universidad Michoacana de San Nicolás de Hidalgo, Km. 9.5 carretera Morelia-Zinapécuaro, 58880 Tarímbaro, Michoacán, Mexico

<sup>d</sup> Center for Tropical Research, Institute of the Environment and Sustainability, Universidad de California, Los Ángeles, Charles E. Young Dr. East 619, 90095, Los Angeles, CA, USA

Received 4 January 2015; accepted 8 December 2016

Available online 2 May 2017

## Abstract

The Baird's tapir (*Tapirus bairdii*) is the largest terrestrial mammal dwelling in the neotropics and the only native representative of the order Perissodactyla. This species is under a great threat due to the impacts of hunting and habitat loss. Therefore, implementation of conservation strategies, *in situ* and *ex situ*, is urgent. However, to improve effectiveness of these strategies knowledge of the species level of genetic variability is needed. We collected samples (hair, feces and blood) from *T. bairdii* individuals kept in captivity in 4 zoos in Mexico and the field (feces) to extract DNA. We analyzed tapir levels of genetic variability focusing on a 511 bp region of the mtDNA D-loop and a set of 6 microsatellite loci. We found a relatively low allelic richness  $R_s = 3.94$  but a relatively high frequency of private alleles (6 out of 24) and a variability of mother lines higher than reported in other studies (7 haplotypes with 15 polymorphic sites,  $H_d = 0.846$  and  $P_i = 0.0128$ ). Our results suggest that there might be a relatively high genetic diversity in tapir population inhabiting southern Mexico.

© 2017 Universidad Nacional Autónoma de México, Instituto de Biología. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

**Keywords:** Mammal conservation; Genetic variability; Conservation genetics; Perissodactyla; Mitochondrial D-loop

## Resumen

El tapir centroamericano (*Tapirus bairdii*) es el mamífero terrestre de mayor talla en el neotrópico y el único representante nativo del orden Perissodactyla. Esta especie se encuentra bajo un fuerte peligro de extinción por la cacería y la destrucción de su hábitat. Resulta urgente la implementación de estrategias de conservación *in situ* y *ex situ* de la especie; sin embargo, se requiere contar con información sobre su variabilidad genética. Obtuvimos ADN de *T. bairdii* a partir de muestras (pelo, excretas y sangre) de individuos en cautiverio en 4 zoológicos de México y excretas recolectadas en el campo. Analizamos los niveles de variabilidad genética en un fragmento de 511 pares de base de la región D-loop del ADN mitocondrial y en microsatélites para 6 loci. Encontramos una riqueza alélica baja ( $R_s = 3.94$ ) pero un número relativamente alto de alelos exclusivos (6 de un total de 24) en microsatélites así como una variabilidad de las líneas maternas más altas que la registrada en otros estudios.

\* Corresponding author.

E-mail address: [mendoza.mere@gmail.com](mailto:mendoza.mere@gmail.com) (E. Mendoza).

Peer Review under the responsibility of Universidad Nacional Autónoma de México.

(7 haplotipos con 15 sitios polimórficos,  $H_d = 0.846$  and  $P_i = 0.0128$ ). Esto sugiere la existencia de una diversidad genética relativamente alta en las poblaciones de tapir que habitan en el sureste de México.

© 2017 Universidad Nacional Autónoma de México, Instituto de Biología. Este es un artículo Open Access bajo la licencia CC BY-NC-ND (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

**Palabras clave:** Conservación de mamíferos; Variabilidad genética; Genética de la conservación; Perissodactyla; D-loop mitocondrial

The Baird's tapir (*Tapirus bairdii*) is the largest terrestrial vertebrate (>250 kg) living in the Neotropics and the only native representative of the Perissodactyla (one of the most threatened orders of mammals) in the Americas (Baillie, Griffiths, Turvey, Loh, & Collen, 2010). This species is classified as endangered due to the extreme reduction of its population and habitat (Schank et al., 2015). Southeast Mexico is one of the most important strongholds for *T. bairdii* but is subjected to an increased pressure due to human activities (Mendoza et al., 2013). The level of endangerment of the species makes strategies for its recovery, such as *ex situ* management, greatly needed; however, its implementation requires data on tapir's genetic variability which is mostly missing (Ruiz-García et al., 2012).

We requested biological samples from 8 zoos in Mexico keeping tapirs (15 individuals) in captivity. We obtained 11 samples (hair, feces and in 1 case blood) from the following zoos: ZooMAT, ZooGuadalajara, ZooLeón and Africam Safari. Four of the samples were from tapirs born in the wild in southern Mexico, 3 samples were from tapirs born within the breeding program of the ZooMAT, and the remaining samples were from tapirs born in USA zoos and donated to Mexico. Tapir hair was collected, wearing gloves, and using a pet brush to later be stored in plastic bags. Feces samples (3–20 ml in volume) were taken following the manual of the genetics committee of the IUCN/SSC Tapir Specialist Group (TSG, 2005). The blood sample ( $\approx 100 \mu\text{l}$ ) was collected from the tapir's back leg using a syringe and stored in an EDTA blood Vacutainer. Twenty-eight feces samples (1–10 ml in volume) were collected in different localities within the range of distribution of Baird's tapir in Chiapas, Mexico. These samples were preserved in 95% ethanol and stored at  $-20^\circ\text{C}$ . Total DNA from hair and blood samples was extracted using the salting-out protocol (Miller, Dykes, & Polesky, 1988). Total DNA from feces was extracted using a variation of the Orsini and Romano-Spica (2001) and Hoshino's and Matsumoto (2004) protocols. Samples (20–40 g) were dried and pulverized to later be incubated in 1.5 ml microtubes containing lysis buffer (200 mM, pH 8.0, Tris HCl, 2% SDS, 4% PVP, 100 mM NaCl, 50 mM EDTA) and 5–10 mg of powdered skim milk at  $60^\circ\text{C}$  using a thermomixer for 15 min. Pulses of 3–5 s at 600 W were applied to the mixture every 5 min in a microwave leaving the container's lid open. Protein, carbohydrates and other residues were separated using a phenol-chloroform protocol (Sambrook & Russell, 2001) 1–2 times up to the solution had a clear coloration. To remove residual salts, the DNA samples were dialyzed in Petri dishes, using 0.025 mm micropore membranes and sterile distilled deionized water. The resulting solution was concentrated using an Eppendorff Vacufuge.

Ten different microsatellites loci previously developed using *Tapirus terrestris* and *T. bairdii* sequences were tested using

DNA feces samples of *T. bairdii* requested to the Los Angeles Zoo. Six of the 10 microsatellites (Table 1) occurred in all the tested individuals thus, were selected for the analyses. For detection analysis primers were marked with 5-FAM fluorochrome (Invitrogen). Each reaction was prepared following the instructions of the platinum Taq DNA polymerase kit (Invitrogen). After preparation, the samples were run in a GeneAmp PCR System 2700 (Applied Biosystems) using the conditions reported for their authors (Table 1).

Allelic richness ( $R_s$ ) was estimated with the softwares GeneAEx version 6.3 (Peakall & Smouse, 2006) and FSTAT 2.9.3 (Goudet, 2001). We estimated the allelic richness of all the captive tapirs in Mexico (samples from tapirs donated by USA zoos and originated from wild populations pooled). Afterwards, we separated these samples to contrast their allelic richness against the corresponding allelic richness of individuals from Costa Rica and Panama (Norton & Ashley, 2004b) applying Wilcoxon tests. We used the software GeneAEx and Genetic Data Analysis GDA (Lewis & Zaykin, 2001) to estimate observed heterozygosity ( $H_o$ ), expected heterozygosity ( $H_e$ ) and to run Hardy–Weinberg (H–G) equilibrium tests. We applied an analysis of molecular variance (Ampova) using the software GeneAEx. We calculated the genetic diversification values of  $F_{IS}$ ,  $F_{IT}$  and  $F_{ST}$  with the software FSTAT ver. 2.9.3 (Goudet, 2001) and applying a Bonferroni correction ( $\alpha = 0.05$ ). The allelic richness found in Mexican tapirs was compared against the values of 6 microsatellites loci reported by Norton and Ashley (2004b) using Anova.

The primers used to amplify DNA sequences of the D-loop region of Baird's tapir were developed at the Center for Tropical Research of the University of California, Los Angeles. These primers developed for the amplification of the D-loop hypervariable region in horse (Bowling, Del Valle, & Bowling, 2000) were aligned with the horse (*Equus caballus*, Genbank NC\_001640) and Brazilian tapir (*T. terrestris*, Genbank: AJ428947) mitochondrial sequences to identify the hypervariable region of interest for Baird's tapir. Two sets of primers were designed from conserved sequences shared by horse and the Brazilian tapir. The effectiveness of the primers was validated by PCR amplification and sequencing of the target D-loop region in 3 Baird's tapir samples obtained from the Los Angeles zoo.

The D-loop hypervariable region of 13 samples was amplified by PCR using a platinum Taq DNA polymerase kit (Invitrogen) in a GeneAmp PCR System 2700 (Applied Biosystems) with the forward TapDLN-F (5'-ATT ACC CTG GTC TTG TAA ACC AG-3') and reverse TapDLN-R (5'-GCC CTG AAG TAA GAA CCA GAT G-3') primers with the following conditions: an initial denaturation at  $95^\circ\text{C}$  for 5 min in a thermocycler, followed by 30 cycles of 1 min at  $95^\circ\text{C}$ , 1 min at  $47^\circ\text{C}$ , 1 min at  $72^\circ\text{C}$ , and a

Table 1  
Microsatellite primers used to analyze genetic variation in samples of *Tapirus bairdii*.

Name	Motif	Size	Reported alleles	Tm	Authors
TtGT137F	(GT)17	253–286 pb	8	60	Gonçalves da Silva, Lalonde, Quse, Shoemaker, and Russello (2010)
TtGT137R					
Tba20F	(AC)15	260–280 pb	8	63	Norton and Ashley (2004a, 2004b)
Tba20R					
Tba23F	(AC)14	243–253 pb	4	65	Norton and Ashley (2004a, 2004b)
Tba23R					
Tter04F	(TG)20	230–264 pb	11	63	Sanches et al. (2009)
Tter04R					
Tter09F	(CAGG)7	167–192 pb	5	59	Sanches et al. (2009)
Tter09R					
Tter14F	(CA)22	174–208 pb	13	59	Sanches et al. (2009)
Tter14R					

final extension at 72 °C for 8 min. The sequences of both DNA strands were obtained using a commercial provider (Macrogen USA). The nucleotide sequences were edited and aligned using BioEdit 7.0 (Hall, 1999) and compared using BLAST (NCBI/Basic local alignment search tool BLAST).

The phylogenetic analysis was conducted using a maximum likelihood (ML) and a Bayesian criteria using softwares MEGA v6 (Tamura et al., 2013) and MrBayes v3.1.2 (Ronquist & Huelsenbeck, 2003). For the ML criterion we used the model T9+G, and a discrete gamma distribution, 5 categories and 10,000 bootstrap replicates following the Find Best Model algorithm. For the Bayesian analysis we generated 1,000 trees using the following settings: ngen = 100,000 and sample freq = 100 criteria. The first 250 trees were burn-in and discarded, and the 50% majority rule consensus tree was generated using the remaining trees. *T. terrestris* (Taxid: 9801) was used as out-group. We calculated haplotype and nucleotide diversity and the number of polymorphic sites using DnaSP v5 (Librado & Rozas, 2009). We built a median joining network using the software NETWORK 4.6.0 (Fluxus Technology Ltd., 2013).

We obtained PCR products of the microsatellites in 9 of the 11 samples. A total of 24 alleles were detected for the 6 loci ranging from 3 to 6 alleles/locus with a mean allelic richness  $R_s = 3.94$  (standard deviation,  $s = 0.97$ ) for all tapir samples pooled together and  $R_s = 3.29$  ( $s = 1.01$ ) when including only samples of tapirs from Mexican populations. These  $R_s$  values were not different of homologue data of tapirs from Panama and Costa Rica ( $F < F\text{-crit}$ ,  $p = 0.38$ ). The observed heterozygosity ( $H_o$ ) for each locus was larger than the expected ( $H_e$ ) and the mean  $F_{ST} = 0.021$  was no different between Mexican and USA individuals. The mean value of  $F_{IS} = -0.295$  across the loci suggested an excess of heterozygosity inside the whole group (Table 2). The examination of the H–W equilibrium across the loci for the whole group showed a deviation in the case of Tter09. Nevertheless, the Fisher's exact test applied to each locus showed no deviations from the H–W equilibrium ( $p > 0.05$ ). Likewise, Fisher's exact test applied to samples from captive individuals in Mexico (from USA zoos and local wild populations) showed no significant deviation from the H–W equilibrium. There were no significant differences among the  $F_{ST}$ ,  $F_{IS}$  and  $Fit$  values in the Amova analysis ( $P$  (random  $\geq$  data).

We found 6 private alleles, 4 of them within samples of Mexican tapirs and 2 more within samples of tapirs donated by USA zoos. We isolated DNA from 21 of the 28 samples of feces from the field and obtained amplicons in 13 out of these 21 samples; 7 of them were within the expected range of weight for the D-loop fragment. Only 3 of these sequences were clear enough to be analyzed (MP3324, MP3335 and MP3338). Therefore, there were only 10 sequences of the control region from individuals in captivity and 3 from samples collected in the field. No sequences of the control region of *T. bairdii* were found in the data bank of the National Center for Biotechnology Information (NCBI). However, the analysis of the 10 *T. bairdii* sequences using BLASTn indicated a similarity of up to 87% with the mtDNA of *T. terrestris* (GeneBank: AJ428947.1) and up to 81% with the mtDNA of *T. indicus* (GeneBank: KJ417810). Moreover, this analysis showed the existence of 7 haplotypes with 15 polymorphic sites,  $H_d = 0.846$  and  $P_i = 0.0128$ . All the nucleotide changes were transitions. The dendrograms generated using maximum likelihood and Bayesian inference showed that samples of tapirs donated by USA zoos were arranged in a monophyletic sub-group whereas samples of tapirs from wild populations in Mexico constituted a paraphyletic group. The divergence in the control region sequences within tapirs from Chiapas was up to 0.021% whereas within tapirs from USA zoos, was 0.005% (Fig. 1).

The haplotype network (Fig. 2) showed in tapirs from Mexico a separation of up to 15 nucleotide changes between mother lines. In comparison, the USA group showed only 5 nucleotide changes between sequences. Therefore, with a lower number of individuals, it was possible to see more diversification among tapirs from Chiapas than the reported by Ruiz-García et al. (2012) analyzing 23 individuals of *T. bairdii* from Panama and Costa Rica. We included only one of the locus reported by Norton and Ashley (2004b) in our study, yet, microsatellite analysis showed a similar value of allelic richness compared to the obtained for Panama and Costa Rica individuals. We did not find significant differences in allelic richness when comparing Mexico/USA and Mexico/total individuals ( $p = 0.05$ ). In agreement with Norton and Ashley (2004a, 2004b), we found that our estimated levels of allelic richness were relatively low. However, our sample size was smaller than the used by these authors, a

Table 2  
Polymorphism of six microsatellite loci across captive individuals of *Tapirus bairdii*.

Locus	Size (pb)	Ar	Ao	Ae	Ho	He	$F_{IS}$	$F_{ST}$
Tter04	212–232	11	3	1.684	0.500	0.406	−0.090	0.000
Tter09	99–139	5	3	2.602	1.000	0.616	−0.525	0.000
Tter14	182–200	13	6	2.733	0.775	0.616	−0.123	0.000
Tba20	224–246	8	4	2.770	0.875	0.638	−0.286	0.133
Tba23	198–232	4	4	2.701	0.900	0.627	−0.435	0.029
TtGt137	222–246	8	4	3.009	0.875	0.656	−0.312	0.111
Mean		8.167	4	2.583	0.821	0.593	−0.295	0.021

Ar, number of alleles previously reported; Ao, observed alleles; Ae, effective alleles; Ho, observed heterozygosity; He, expected heterozygosity;  $F_{IS}$ , differentiation inside the population;  $F_{ST}$ , differentiation between populations.




Figure 1. Consensus tree based on the D-loop sequences. The support values correspond to maximum likelihood/Bayesian trees.

situation that likely affected the possibility to detect any genetic structure by reducing statistical power of Amova and Anova. The combination of the negative values of  $F_{IS}$  and the  $Ho > He$  relationship might be related to the occurrence of a relatively high proportion of private alleles in a small number of individuals. Since these alleles occurred at low frequencies their effect over the  $F_{ST}$  value seemed to be minimal. There might be high variability among individuals but not enough individuals as to generate detectable genetic fixation within the population.

The analysis of the D-loop region showed more polymorphisms between mother lines from Chiapas than between the 2 mother lines of individuals donated by USA zoos. The ancestors of the USA tapirs came from Panama, Honduras, Nicaragua and Costa Rica. We do not know with certainty the origin of the mother lines of tapirs donated by USA zoos but Ruiz-García et al. (2012) found 2 different main lineages when analyzing mitochondrial (*Cyt-b*) sequences: Mexico-Guatemala and Costa Rica-Panama, which were estimated to have diverged about 2.4 million years ago. Likewise, the haplotypes network obtained in the same study showed only 1–2 nucleotide changes between Costa Rica-Panama groups and the nearest hypothetical ancestor, similarly to what we found in this study for tapirs donated by USA zoos.

It stands out that we found a relatively high number of private alleles and polymorphisms between mother lines in samples of captive tapirs originated from wild populations in Mexico. This result opens the possibility to find more genetic variation in a more comprehensive survey of tapir populations occurring in our country. The level of endangerment of Baird's tapir, locally and globally, makes studies to analyze the genetics' spatial variation of their populations urgent to identify conservation hot spots and to set the basis for a sound management plan for the species.

We would like to thank the staff of the zoos which provided the samples for this work in particular to Carlos Guichard Romero, Epigmenio Cruz Aldán and Marco Antonio Benítez. A. S. Zavala-Páramo had a scholarship (412357/262584) from




Figure 2. Median joining network tree based on the D-loop sequences.



Conacyt. Fieldwork was supported by the Latin American Institute of the University of California at Los Angeles.

## References

- Baillie, J. E., Griffiths, J., Turvey, S. T., Loh, J., & Collen, B. (2010). *Evolution lost: status and trends of the world's vertebrates*. London: Zoological Society of London.
- Bowling, A. T., Del Valle, A., & Bowling, M. (2000). A pedigree-based study of mitochondrial D-loop DNA sequence variation among Arabian horses. *Animal Genetics*, 31, 1–7.
- Fluxus Technology Ltd. (2013). Free phylogenetic network software. Retrieved on May 12th, 2016, from: <http://www.fluxus-engineering.com/sharenet.html>
- Gonçalves da Silva, A., Lalonde, D. R., Quse, V., Shoemaker, A., & Russello, M. A. (2010). Genetic approaches refine ex situ lowland tapir (*Tapirus terrestris*) conservation. *Journal of Heredity*, 5, 581–590.
- Goudet, J. (2001). FSTAT, a program to estimate and test gene diversities and fixation indices version 2.9.3. Retrieved on May 22nd, 2015, from: <http://www2.unil.ch/popgen/softwares/fstat.htm>
- Hall, T. A. (1999). BioEdit: a user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. *Nucleic Acids Symposium, Series*, 41, 95–98.
- Hoshino, Y. T., & Matsumoto, N. (2004). An improved DNA extraction method using skim milk from soils that strongly adsorb DNA. *Microbes Environment*, 1, 13–19.
- Lewis, P. O., & Zaykin, D. (2001). Genetic data analysis: computer program for the analysis of allelic data. Ver. 1.0 (d16c). Retrieved on January 18th, 2015, from: <http://lewis.eeb.uconn.edu/lewishome/software.html>
- Librado, P., & Rozas, J. (2009). DnaSP v5: a software for comprehensive analysis of DNA polymorphism data. *Bioinformatics*, 11, 1451–1452.
- Mendoza, E., Fuller, T. L., Thomassen, H. A., Buermann, W., Ramírez-Mejía, D., & Smith, T. B. (2013). A preliminary assessment of the effectiveness of the Mesoamerican Biological Corridor for protecting potential Baird's tapir (*Tapirus bairdii* Gill, 1865) habitat in southern Mexico. *Integrative Zoology*, 1, 35–47.
- Miller, S. A., Dykes, D. D., & Polesky, H. F. (1988). A simple salting out procedure for extracting DNA from human nucleated cells. *Nucleic Acids Research*, 3, 12–15.
- Norton, J. E., & Ashley, M. V. (2004a). Genetic variability and population differentiation in captive Baird's tapirs (*Tapirus bairdii*). *Zoo Biology*, 2, 521–531.
- Norton, J. E., & Ashley, M. V. (2004b). Genetic variability and population structure among wild Baird's tapirs. *Animal Conservation*, 7, 211–220.
- Orsini, M., & Romano-Spica, V. (2001). A microwave-based method for nucleic acid isolation from environmental samples. *Letters in Applied Microbiology*, 33, 17–20.
- Peakall, R., & Smouse, P. (2006). GENALEX 6: genetic analysis in Excel. Population genetics software for teaching and research. *Molecular Ecology*, 6, 288–295.
- Ronquist, F., & Huelsenbeck, J. P. (2003). MRBAYES 3: Bayesian phylogenetic inference under mixed models. *Bioinformatics*, 19, 1572–1574.
- Ruiz-García, M., Vásquez, C., Pinedo-Castro, M., Sandoval, S., Castellanos, A., Kaston, F., et al. (2012). Phylogeography of the Mountain Tapir (*Tapirus pinchaque*) and the Central American Tapir (*Tapirus bairdii*) and the origins of the three Latin-American tapirs by means of mtCyt-B sequences. In K. Ananthawat-Jónsson (Ed.), *Current topics in phylogenetics and phylogeography of terrestrial and Aquatic Systems* (pp. 83–116). INTECH Open Access Publisher. Available: <http://www.intechopen.com/books/current-topics-in-phylogenetics-and-phylogeography-of-terrestrial-and-aquatic-systems/phylogeography-of-the-mountain-tapir-tapirus-pinchaque-and-the-central-american-tapir-tapirus-bairdi>
- Sambrook, J., & Russell, D. W. (2001). *Molecular cloning: a laboratory manual*. New York: Cold Spring Harbor Laboratory Press.
- Sanches, A., Gomes-de Figueiredo, M., Fonseca-Pereira, F., Silveira, L., Jácomo, A. T. A., & Galetti, P. M. (2009). Microsatellite loci isolated from the lowland tapir (*Tapirus terrestris*), one of the largest Neotropical mammal. *Conservation Genetic Resources*, 1, 115–117.
- Schank, C., Mendoza, E., Manolo, J., García, V., Cove, M. V., Jordan, C. A., et al. (2015). Integrating current range-wide occurrence data with species distribution models to map the potential distribution of Baird's Tapir. *Tapir Conservation*, 33, 15–25.
- Tamura, K., Peterson, D., Peterson, N., Stecher, G., Nei, M., & Kumar, S. (2013). Molecular evolutionary genetics analysis, version 6.0. *Molecular Biology and Evolution*, 12, 2725–2729.
- TSG (Tapir Specialist Group). 2005. How to sample dung for genetic analyses. Genetics committee, IUCN/SSC Tapir Specialist Group. Retrieved on August 8th, 2012, from: <http://tapirs.org/committees/genetics/sampledung/sampledung.html>