



Revista Mexicana de Biodiversidad

ISSN: 1870-3453

falvarez@ib.unam.mx

Universidad Nacional Autónoma de
México
México

Prieto-Barajas, Cristina M.; Alfaro-Cuevas, Ruth; Valencia-Cantero, Eduardo; Santoyo,
Gustavo

Effect of seasonality and physicochemical parameters on bacterial communities in two hot
spring microbial mats from Araró, Mexico

Revista Mexicana de Biodiversidad, vol. 88, núm. 3, septiembre, 2017, pp. 616-624

Universidad Nacional Autónoma de México
Distrito Federal, México

Available in: <http://www.redalyc.org/articulo.oa?id=42553212013>

- How to cite
- Complete issue
- More information about this article
- Journal's homepage in redalyc.org

redalyc.org

Scientific Information System

Network of Scientific Journals from Latin America, the Caribbean, Spain and Portugal

Non-profit academic project, developed under the open access initiative



Ecology

Effect of seasonality and physicochemical parameters on bacterial communities in two hot spring microbial mats from Araró, Mexico

Efecto de la estacionalidad y de los parámetros fisicoquímicos sobre las comunidades bacterianas de tapetes microbianos de manantiales termales de Araró, México

Cristina M. Prieto-Barajas, Ruth Alfaro-Cuevas, Eduardo Valencia-Cantero, Gustavo Santoyo*

Instituto de Investigaciones Químico Biológicas, Universidad Michoacana de San Nicolás de Hidalgo, Edificio A1', Ciudad Universitaria, 58030 Morelia, Michoacán, Mexico

Received 7 November 2016; accepted 5 May 2017

Available online 18 August 2017

Abstract

In this study, we explored the diversity of culturable bacterial communities residing in hot springs from Araró, México, and analysed the effect of seasonality and related changes in physicochemical parameters of spring water. Two hot springs with unique features, Tina and Bonita, were analysed. Seventy-nine unique 16S rRNA gene phylotypes were detected, belonging to the bacterial phyla Firmicutes, Proteobacteria, and Actinobacteria. A group of dominant phylotypes of the genus *Bacillus* was recovered in 3 out of 4 of the sampling seasons. Another group of phylotypes was recovered in 2 samplings, while the remaining groups were detected in only 1 season. Ecological indexes for species richness and evenness showed moderate to low diversity in both hot springs, and a Sørensen analysis revealed that the 2 communities shared 64% of their bacterial phylotypes. Physicochemical parameters measured every season showed slight variations, except for temperature and arsenic content. Fluctuations in bacterial composition in the Tina hot spring were correlated mainly with salt content, while diversity in the Bonita hot spring was significantly correlated with temperature, pH, and arsenic content.

© 2017 Universidad Nacional Autónoma de México, Instituto de Biología. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Keywords: Microbial mats; Hot springs; Bacterial diversity; Environmental factors; Arsenic; Thermophilic communities

Resumen

En este estudio se explora el efecto de la estacionalidad y los cambios fisicoquímicos sobre la diversidad cultivable de la comunidad bacteriana de dos manantiales termales de Araró, México. Se encontraron 79 filotipos de 16S rRNA de las divisiones bacterianas Firmicutes, Proteobacteria y Actinobacteria. Un grupo de filotipos del género *Bacillus* fue el más dominante, aislado en 3 de las 4 estaciones muestreadas, otro grupo de filotipos se recuperó en 2 muestreos, mientras que los grupos restantes se detectaron en sólo 1 temporada. Los índices ecológicos para riqueza de especies y equitabilidad muestran una diversidad de moderada a baja en ambos manantiales, mientras que el índice de Sørensen resulta en que ambas comunidades de comparten hasta un 64% del total de los filotipos. Los parámetros fisicoquímicos medidos muestran ligeras variaciones estacionales, excepto la temperatura y el contenido de arsénico. Las fluctuaciones en la composición bacteriana en Tina estuvieron correlacionadas principalmente al contenido salino mientras que en Bonita las variaciones se asociaron con la temperatura, pH y contenido de arsénico.

© 2017 Universidad Nacional Autónoma de México, Instituto de Biología. Este es un artículo Open Access bajo la licencia CC BY-NC-ND (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Palabras clave: Tapetes microbianos; Manantiales termales; Diversidad bacteriana; Factores ambientales; Arsénico; Comunidades termófilas

* Corresponding author.

E-mail address: gsantoyo@umich.mx (G. Santoyo).

Peer Review under the responsibility of Universidad Nacional Autónoma de México.

Introduction

Hot springs can be found in many regions of the world. Each hot spring has unique geophysical and biological characteristics, making them interesting subjects of research with potential value for either biotechnological or ecological purposes (Briggs et al., 2014; Brock, 1997; Klatt et al., 2011; Lau, Aitchison, & Pointing, 2009; Tirawongsaroj et al., 2008). Because of these differences, bacterial diversity in microbial mats is expected to be different among hot springs. Analyses of microbial diversity have been reported for hot springs located in many countries, including the USA (Ward, Ferris, Nold, & Bateson, 1998), Russia (Perevalova et al., 2008), Australia (Kimura et al., 2005), Thailand (Portillo, Sirin, Kanoksilapatham, & González, 2009), Bulgaria (Tomova et al., 2010), Colombia (Bohorquez et al., 2012), Philippines (Huang et al., 2013), Chile (Mackenzie, Pedrós-Alió, & Díez, 2013) and China (Briggs et al., 2014).

The physicochemical parameters of source waters in some hot springs can vary by season while in others, these parameters can be very stable. These parameters can influence the microbial diversity in springs. A few studies have assessed these parameters in association with microbial community composition in hot springs (Briggs et al., 2014; Ferris & Ward, 1997; Mackenzie et al., 2013). For example, microbial diversity was surveyed in 3 hot springs located in La Patagonia, Chile, over 2 seasons. The authors found differences in the microbial communities between the seasons, probably due to temperature variations (Mackenzie et al., 2013). Other factors, such as pH, levels of hydrogen sulphide, and temperature, have been associated with the presence of certain microbial species (Briggs et al., 2014; Purcell et al., 2007; Ward & Castenholz, 2000).

To determine the bacterial diversity in microbial mats, both cultivation-dependent and -independent approaches have been employed, depending on the research goals. A cultivation-independent methodology was used in a recent PhyloChip microarray analysis of the microbial communities inhabiting hot springs of the Tengchong region in China (Briggs et al., 2014). However, this technology is not appropriate for the isolation of thermophilic microorganisms, their genes, or their proteins to determine their potential importance for biotechnology applications (Brock & Freeze, 1969). The 2 approaches can be complementary; however, an initial microbial diversity analysis can be a first step in searching for bacteria with enzymatic activities for industrial application. A good example of this approach is a report by Kanokratana, Chanapan, Pootanakit, and Eurwilaichitr (2004), which first described the biodiversity of bacteria and archaea from hot springs in Thailand. Years after this work, they reported on a functional screening of a metagenomic library derived from sediments of the hot springs, which resulted in the isolation of 2 novel genes encoding an esterase and a phospholipase (Tirawongsaroj et al., 2008).

In the present study, using a culture-dependent approach, we investigated the seasonal variation (Spring, Summer, Fall, and Winter) in bacterial diversity of 2 hot springs located in the geothermal system of Araró, Michoacán, México, which is a system independent from Los Azufres geothermal zone (Brito et al., 2014; Viggiano-Guerra & Gutiérrez-Negrín, 2005). The

physicochemical parameters of both hot springs were assessed and associated with fluctuations in bacterial biodiversity.

Materials and methods

The geothermal system of the Araró region is located in the central part of Mexico, inside the Trans Mexican Volcanic Belt located in Michoacán State. The region is approximately 20 km west of the well-known Los Azufres geothermal field (Fig. 1). The zone known as Zimirao (19°53'54" N, 100°49'50" W) is where most of the hot springs are located. There are about 50 hot springs in the Araró region, and many of them are used for recreational activities; however, the selected hot springs, Tina and Bonita, have relatively little disturbance and are far from the recreational area. Another interesting feature of these hot springs is that Bonita has low water emission and forms colourful microbial mats, while Tina has a constant water effluent and a little running stream, with microbial mats formed along the stream (Fig. 2). Compared to Bonita, Tina presents only a very thin bacterial mat. Therefore, the systems exhibit different features that might influence the communities present in microbial mats.

Four samplings were conducted on February 2nd (Winter), June 5th (Spring), October 6th (Summer), and December 5th (Fall) of 2013. Three samples were directly collected from microbial mats in each of the 2 hot springs at a depth of 30–50 cm from the surface. A criterion for selection of these 2 hot springs was whether the system was closed or open. One spring selected was a “closed system” with low water emission (Bonita), while the other (Tina) was an “open system” with constant water emission (Fig. 2). Microbial mat samples were immediately transported on wet ice to the lab. For water sampling, 500-mL sterile Kinex flasks were used to take water from each hot spring. Water samples were stored in darkness and transported to lab on dry ice.

Physicochemical parameters of water in the hot springs were measured during sampling of biological material. The parameters, including temperature (°C), electrical conductivity, pH, and dissolved oxygen, were measured *in situ* with a Corning® Checkmate™ II modular meter system.

Physicochemical analyses, including faecal coliforms analysis, were performed in collaboration with the National Water Commission (Conagua-México) (Table 1). Arsenic concentrations in the water samples of the Bonita and Tina hot springs were measured by absorption spectroscopy using an atomic absorption spectrometer (Perkin-Elmer Analyst 200) with a hydride generation system. The measurement of fluoride was carried out with a conventional fluorometer.

A small sample (0.1 g) of the microbial mat from each spring was placed into Eppendorf tubes (1.5 mL) with 990 µL of sterile water, vortexed, and plated onto 3 different 10-fold-diluted culture media. These included salted, rich, and poor media (Luria-Bertani, Nutrient Agar (NA), and Minimal Medium (MM), respectively, purchased from Sigma-Aldrich). The LB medium contained (/L) yeast extract (5 g), tryptone (10 g), NaCl (10 g). NA medium contained (/L) peptone (5 g) and beef extract




Figure 1. A map that shows the location of the Araró, Michoacán, México (red star). The Araró hot springs are located within the Trans-Mexican Volcanic Belt, approximately 20 km west from the well-known Los Azufres geothermal field.




Figure 2. A view of the 2 hot springs from Araró. On the left (A) is the so called Tina hot spring, with constant emission of water. Panel B shows a view of the Bonita hot spring, a “closed system” with low emission of water.

Table 1

Physicochemical parameters measured from Araró geothermal zone, Tina and Bonita hot springs, 4 seasons sampled.

Hot spring	Season	T	pH	BOD	TS	EC	Cl	SO ₄	A	Ca	Mg	Na	As
Tina	Winter	63	7.35	0.8	2,423	3,860	977	246	330	188	25	625	4.9
	Spring	66	7.64	0.6	3,722	3,840	960	245	338	207	28	581	4.7
	Summer	78	6.95	0.6	2,616	3,950	976	252	347	209	31	610	6.6
	Autumn	74	7.07	3.0	2,410	3,510	867	228	330	153	45	545	6.1
Bonita	Winter	50	6.75	0.6	2,896	3,650	928	239	310	230	22	533	3.7
	Spring	45	7.03	0.8	3,184	3,801	951	245	337	202	29	586	4
	Summer	50	7.49	0.4	2,398	3,670	917	237	328	187	32	569	2.9
	Autumn	55	7.78	2.6	2,502	3,650	910	239	328	158	48	567	2.7

T, temperature (°C); BOD, biochemical oxygen demand (mg/l); TS, total solids*; EC, electrical conductivity (μmhos/cm); Cl, chloride salts*; SO₄, sulphates*; A, alkalinity*, Ca*, Mg*, Na*, As*; *reported as mg/L.

(5 g). MM medium contained (L) glucose (2 g), NaHPO₄ (6 g), KH₂PO₄ (3 g), NaCl (0.5 g), NH₄Cl (1 g), MgSO₄ (0.5 g), CaCl (0.01 g), tyrosine (0.1 g) and agar (15 g). Most bacterial isolates (90%) grew well and were selected on LB media. Plates were incubated for 72 h at 37 °C and the number of colony forming units (CFU) was calculated per gram of microbial mat.

Randomly picked colonies were subject to further serial dilutions to obtain single isolates. Isolates were ON cultured in 3 ml of medium. Genomic DNA was isolated by following the protocol of the Wizard[®] Genomic DNA Purification Kit and subjected to PCR amplification of their 16S ribosomal subunit gene. Other bacterial isolates (105 from each hot spring) that exhibited an identical fingerprint by RAPDs analysis (Random Amplification of Polymorphic DNA) were assigned to the sequenced phylotypes to report presence/absence during the sampling seasons. The primer employed for the RAPD analysis was OPA02, 5'-TGC CGA GCT G-3' (Samal et al., 2003). Bacterial primers fd1, 5'-CAGAGTTTGATCCTGGCTCAG-3', and rD1, 5'-AAGGAGGTGATCCAGCC-3', were used (Weisburg, Barns, Pelletier, & Lane, 1991) to amplify nearly full-length 16S ribosomal genes. PCR conditions have been previously reported (Hernández-León et al., 2015). All the PCR products were purified and sequenced at the LANGE BIO (Irapuato, Mexico). The 16S rDNA sequences obtained were subjected to homology blast searches against databases and deposited in GenBank (Accession Numbers: KJ801569–KJ801648).

Multiple sequence alignments were generated and a phylogenetic analysis of 16S rRNA gene sequences was carried out using the MEGA 5.0 program (Tamura et al., 2011). All sequences passed quality controls and a cut-off value of 98% similarity was applied. To obtain a confidence value for the aligned sequence dataset, a bootstrap analysis of 1,000 replications was performed. A phylogenetic tree was constructed using a maximum likelihood algorithm.

Diversity indexes, including the Shannon (Shannon, 1948), evenness, and Simpson (Simpson, 1949) indexes, were calculated to infer species richness and abundance. Sørensen similarity index (Sørensen, 1948) was used to compare the similarity of the bacterial communities from Tina and Bonita microbial mats. Rarefaction curves were generated to evaluate sampling efforts. Pearson's coefficient with a 90% confidence value was used to determine significant positive relationships between diversity and any physicochemical parameters (PAST software).

Results

The physicochemical parameters of both hot springs are presented in Table 1. Bonita and Tina hot springs have different superficial temperatures. Bonita temperatures range from 45 °C to 55 °C throughout the year, while in Tina, they vary from 63 °C to 78 °C. A neutral pH was found in both hot springs, with little seasonal variation; however, greater variation was found in Bonita, with a pH range of 6.75–7.78, whereas in Tina it ranged from 6.95 to 7.64. Salt contents, including Ca, Cl, SO₄, Mg, and Na, of the springs also showed variation across seasons and differences between hot springs. Interestingly, arsenic concentration varied little across seasons in each hot spring, but concentrations were much higher in Tina (4.7–6.6 mg/L) than in Bonita (2.7–4 mg/L). Conductivity values were slightly higher for Tina during Winter, Spring, and Summer. Faecal coliforms (NMP/100 m³) were not detected in either hot spring (data from Conagua).

In order to detect the abundance of culturable bacterial cells, the number of CFUs per gram of microbial mat was determined from each hot spring. In general, the number of CFUs was relatively low and fluctuated slightly across seasons. However, it was interesting to note that Bonita showed a 10-fold higher CFU count than the Tina hot spring (1×10^2 CFU/g and 1×10^3 CFU/g, respectively), regardless of the season.

From randomly picked colonies subjected to RAPD analysis, 79 phylotypes (out of 109 bacterial isolates) were detected and subjected to 16S rDNA sequencing. Bacteria from 3 phyla were identified, including species from the genera *Paenibacillus*, *Exiguobacterium*, and *Bacillus* (76%) from the phylum Firmicutes, *Aeromonas* and *Pseudomonas* (18%) from the phylum Proteobacteria, and only *Microbacterium* (6%) from the phylum Actinobacteria. In total, 17 culturable bacterial species were detected in both hot springs. Figure 3 shows the phylogenetic relationships of the bacterial species. Fifty-seven 16S rDNA sequences, including the 17 representative bacterial species shared between both hot springs, were selected for the phylogenetic reconstruction based on length. All of the 16S rDNA sequences displayed close relationships with known bacterial species, particularly with representatives of the *Bacillus* genus. The clades were not associated with either of the hot springs or any other parameter measured.




Figure 3. Phylogenetic tree of the 79 phylotypes identified in both hot springs from Araró. Analysis of the 16S rDNA gene sequences was carried out with the MEGA 5.0 program. The tree was constructed by using the maximum likelihood algorithm with a bootstrap analysis of 1,000 replications.

A dominant group of *Bacillus* was present in at least 3 out of 4 samplings and included the species *Bacillus licheniformis*, *Bacillus cereus*, *Bacillus subtilis*, and *Bacillus pumilus* (Fig. 4). The frequent species were *Bacillus megaterium*, *Bacillus vietnamsensis*, *Bacillus boroniphilus*, *Aeromonas hydrophila*, *Bacillus amyloliquefaciens*, and *Exiguobacterium profundum*, which appeared in 2 sampling seasons, while occasional bacterial species appeared only in 1 sampling and included *Paenibacillus pabuli*, *Paenibacillus favisporus*, *Exiguobacterium sibiricum*, *Bacillus oceanisediminis*, *Pseudomonas psychrotolerans*, *Microbacterium oleivorans*, and *Pseudomonas stutzeri*.

The bacterial communities of the microbial mats from Tina and Bonita hot springs were characterised by a few dominant taxa within the Firmicutes and by low abundance (Fig. 4). The Shannon index showed a similar low diversity of bacterial species for each hot spring in each of the 4 sampling seasons. The highest diversity was observed in the Spring and the lowest diversity was observed in the Fall for both Tina and Bonita. According to the Sørensen index, the 2 hot springs shared up to 64% of their bacterial species. However, this percentage varied, and in some seasons was much lower (Table 2). For example, during the Fall sampling, both springs shared only 16%, opposed to Spring, where they shared almost 50% of bacterial species.

Pearson's correlation analysis suggested that water salts (such as Mg, Na, Cl), biological oxygen demand, and electrical conductivity all had effects on bacterial communities inhabiting Tina hot spring, whereas in Bonita, temperature, pH, and arsenic concentration exerted effects on bacterial diversity (Table 3). These results suggest that Tina and Bonita share some physical and biological features, but are not interconnected subsystems.

Discussion

The present study represents the first exploration of bacterial diversity in 2 of the approximately 50 hot springs located in Araró, Michoacán, which is in the middle of the Trans Mexican Volcanic Belt (Israde-Alcántara & Garduño-Monroy, 1999). Unfortunately, most thermal springs in Araró, and nearby regions, are used for recreational purposes and have been drastically modified, leaving the undisturbed ecosystems unexplored and likely reducing their microbiological diversity and biotechnological potential (López-Sandoval, Montejano, Carmona, Cantoral, & Becerra-Absalón, 2016). Therefore, there is a need for microbial censuses of the few hot springs in Araró that remain undisturbed.

Most recent publications on the biodiversity of microbial mats from hot springs around the world have described studies based on non-culture methods, including denaturing gradient gel electrophoresis (DGGE), PhyloChip microarray, and sequencing of 16S rDNA gene libraries. Some of these studies have analysed the effect of seasonality and physicochemical parameters on microbial diversity (Briggs et al., 2014; Lacap, Barraquío, & Pointing, 2007; Mackenzie et al., 2013). Here, it was our short-term goal to describe the bacterial diversity in microbial mats by first isolating aerobic heterotrophic thermophilic bacteria. Our long-term goal, however, is to obtain easy-culturable bacterial isolates for bioremediation or biodegradation. Araró hot

Table 2
Ecological indexes for each sampling at Araró geothermal zone.

Ecological indexes	Hot spring							
	Bonita				Tina			
	Winter	Spring	Summer	Autumn	Winter	Spring	Summer	Autumn
Shannon	1.91	2.02	1.53	1.07	1.91	1.77	1.75	1.17
Evenness	0.96	0.94	0.77	0.73	0.84	0.84	0.82	0.80
Simpson	0.16	0.14	0.26	0.44	0.17	0.20	0.21	0.37
Sorensen ^a	64%							
Sorensen ^b	Winter 40%		Spring 48%		Summer 32%		Autumn 16%	

^a Sorensen bacterial community shared between Tina and Bonita.

^b Sorensen bacterial communities shared each sampled season.

springs are extreme environments where bacteria with potential for biodegradation may reside. Our results suggest that bacteria belonging primarily to division Firmicutes occur there (Fig. 3). Interestingly, of the 13 species of Firmicutes isolated, only *B. megaterium* (Baker, Gaffar, Cowan, & Suharto, 2001), *E. profundum* (Crapart et al., 2007), and a strain of *Paenibacillus* (Mead et al., 2012) have been isolated from thermal environments. To our knowledge, this is the first time that the remaining species have been identified as inhabitants of hydrothermal systems. *B. subtilis*, *B. cereus*, *B. licheniformis*,

B. pumilus, and *B. megaterium* are common soil inhabitants (Slepecky & Hemphill, 2006), although they have the capacity to grow at high temperatures (Slepecky & Hemphill, 2006; Yakimov, Timmis, Wray, & Fredrickson, 1995). The vast majority (> 90%) of the bacterial isolates from Tina and Bonita hot springs grew optimally at 50 °C (data not shown). Several studies have demonstrated that thermophilic bacilli, grown optimally above 40 °C, possess the potential to degrade or convert environmental pollutants (Margesin & Schinner, 2001). In addition, mat-forming cyanobacterial strains have shown synergistic

Hot spring	Tina				Bonita			
	Sampled seasons							
Bacterial species	W	Sp	Su	A	W	Sp	Su	A
<i>B. licheniformis</i>								
<i>B. cereus</i>								
<i>B. subtilis</i>								
<i>B. pumilus</i>								
<i>B. megaterium</i>								
<i>B. vietnamensis</i>								
<i>B. boroniphilus</i>								
<i>A. hydrophila</i>								
<i>B. amyloliquefaciens</i>								
<i>E. profundum</i>								
<i>P. pabuli</i>								
<i>P. favisporus</i>								
<i>E. sibiricum</i>								
<i>B. oceanisediminis</i>								
<i>P. psychrotolerans</i>								
<i>M. oleivorans</i>								
<i>P. stutzeri</i>								

Figure 4. Occurrence of bacterial species isolated during the 4 seasons (Wi, winter; Sp, spring; Su, summer; Au, Autumn) of the year in the hot spring microbial mats from Araró, México. Bacterial species were classified as dominant (isolated in at least 3 out of 4 sampling seasons), frequent (isolated in 2 sampling seasons) and occasional (isolated only in one sampling season).

Table 3

Correlation between physicochemical parameters and bacterial fluctuations. In bold the higher correlations for each sampled site, Tina and Bonita.

Physicochemical parameters	Pearson correlation coefficients	
	Tina	Bonita
Arsenic	47.4	94.4
Calcium	82.3	87.9
Chlorides	97.9	87.0
DBO	95.4	78.9
Electrical conductivity	92.0	60.4
Magnesium	99.4	80.7
pH	42.5	91.7
Sodium	91.3	6.8
Sulphates	89.8	55.8
Temperature	50.1	90.5
Total solids	29.1	81.8

effects in degrading petroleum compounds like phenanthrene, pristane, n-octadecane, and dibenzothiophene (Abed & Köster, 2005).

Another genus frequently isolated from Tina and Bonita hot springs was *Exiguobacterium*. Strains of this genus have been isolated from many different sources, including glacier ice, hot springs in Yellowstone National Park, as well as other extreme environments (Vishnivetskaya, Kathariou, & Tiedje, 2009). This genus comprises psychrophilic, mesophilic, and moderately thermophilic species (Vishnivetskaya et al., 2009). *E. profundum* was also a frequent isolate. It is described as a halotolerant and moderately thermophilic bacterium, and it has been isolated from hydrothermal vents in the Pacific Ocean at depths of about 2,600 m (Crapart et al., 2007). *Aeromonas hydrophila* is a bacterium that has not been associated with high temperatures (Janda & Abbott, 2010). However, some bacterial strains that are closely related to *Aeromonas salmonicida* (Huang et al., 2011) and *A. sober* have been obtained from hot springs in Tibet (Yim, Hongmei, Aitchison, & Pointing, 2006).

The region of Araró is close to the geothermal zone of Los Azufres, which comprises several hot springs, fumaroles, and boiling mud pools, and was explored in the 1980s for its geothermal potential (Birkle & Merkle, 2000). Recently, the microbial diversity in microbial mats and mud pools from Los Azufres were analysed by T-RFLP and 16S rRNA gene library analyses, and the presence of a few genera, such as *Rhodobacter*, *Acidithiobacillus*, *Thiomonas*, *Desulfurella*, and *Thermodesulfobium*, was reported. The authors were also able to isolate 2 sulphate and sulphur reducers related to *Thermodesulfobium* and *Desulfurella* (Brito et al., 2014). Our results are not comparable to those found by Brito and colleagues, since the thermal systems, project goals, and methodologies to assess the bacterial diversity were very different. Tina and Bonita hot springs exhibit clear water emissions with well-formed, colourful microbial mats, while Brito and colleagues mainly analysed microbial mats from mud pools. This difference supports a previous suggestion that the Araró region represents an independent thermal system from Los Azufres, despite their close proximity (Viggiano-Guerra & Gutiérrez-Negrín, 2005). However, the

microbial diversity from the 2 thermal systems contains strains with potential biotechnological applications.

The Araró hot springs are extreme thermophilic environments with elevated concentrations of salts and As. This is in agreement with diversity indexes that indicate low bacterial diversity. In addition, no significant changes in the microbial communities were detected over the 4 samplings. Our results suggest that the dominant species of Firmicutes inhabit and remain part of microbial mats in these springs. The Sørensen analysis also shows that Tina and Bonita hot springs share 64% of their bacterial species. However, similarity was variable, with some seasons showing similarities below 50%, suggesting that these springs are independent from each other. However, a deeper analysis by PhyloChip or 16S rDNA gene pyrosequencing could reveal a wide panorama of non-culturable microbial communities in both hot springs.

No significant correlation was found between temperature and bacterial diversity in Tina spring. However, for Bonita, our results suggest a significant correlation with this parameter. This is probably due to the stability of the bacterial community in Bonita spring throughout the year, with microbial mats that were maintained at about 1 cm thickness. It is also a closed system, in which the water emission is very low and water remains stagnant, with no entry or exit of water streams except for rainfall. Microbial communities may be affected by light and changes in temperature across seasons ranging from 45 °C to 55 °C and an environment that changes from a mesophilic to a thermophilic one. In the case of Tina, a thermophilic environment is maintained. The influence of temperature on the diversity of microbial communities has been widely reported for other hot springs (Ferris & Ward, 1997; Mackenzie et al., 2013; Purcell et al., 2007; Ward et al., 1998; Yim et al., 2006).

A significant correlation was also found between pH, Mg, and arsenic concentrations and bacterial community composition for Bonita, but not for Tina spring. In contrast, salt concentrations, electrical conductivity, and biological oxygen demand were correlated with bacterial community composition in Tina spring. Some hot springs, including some located in China and Chile (Hou et al., 2013; Mackenzie et al., 2013), have considerable salt quantities. Sulphates, in particular, have been shown to affect archaeal and bacterial communities along with elevated temperatures, so it is suggested that, together, salt concentrations and temperature regulate prokaryotic diversity in thermophilic environments (Purcell et al., 2007).

Previous reports on the water quality of thermal recreational springs in Araró suggest high concentrations of As, with values ranging from 0.01 mg/L to 6.26 mg/L, which exceeded the World Health Organization (WHO) and Mexican regulations for drinking water use and suggested a potential health risk (Vázquez-Vázquez, Cortés-Martínez, & Alfaro-Cuevas, 2015). Preliminary results in our lab show that most of the *Bacillus* strains isolated from Tina and Bonita springs contain putative arsenic resistance (*ars*) genes, which allow the bacteria to survive in elevated As concentrations. Several thermophilic strains of *Bacillus* isolated from Araró hot springs are being evaluated for bioremediation of metal pollutants.

Acknowledgements

To the Consejo Nacional de Ciencia y Tecnología, México (Project No. 169346) and the Coordinación de la Investigación Científica, Universidad Michoacana de San Nicolás de Hidalgo (2014–2015) for financially supporting our research projects. CMPB received a scholarship from Conacyt, México.

References

- Abed, R. M. M., & Koster, J. (2005). The direct of aerobic heterotrophic bacteria associated with cyanobacteria in the degradation of oil compounds. *International Biodeterioration and Biodegradation*, 55, 29–37.
- Baker, G. C., Gaffar, S., Cowan, D. A., & Suharto, A. R. (2001). Bacterial community analysis of Indonesian hot springs. *FEMS Microbiology Letters*, 200, 103–109.
- Birkle, P., & Merkel, B. (2000). Environmental impact by spill of geothermal fluids at the geothermal field of Los Azufres, Michoacán, Mexico. *Water, Air & Soil Pollution*, 124, 371–410.
- Bohorquez, L. C., Delgado-Serrano, L., López, G., Osorio-Forero, C., Klepac-Ceraj, V., Kolter, R., et al. (2012). In depth characterization via complementing culture-independent approaches of the microbial community in an acidic hot spring of the Colombian Andes. *Microbial Ecology*, 63, 103–115.
- Briggs, B. R., Brodie, E. L., Tom, L. M., Dong, H., Jiang, H., Huang, Q., et al. (2014). Seasonal patterns in microbial communities inhabiting the hot springs of Tengchong, Yunnan Province China. *Environmental Microbiology*, 16, 1579–1591.
- Brito, E. M. S., Villegas-Negrete, N., Sotelo-González, A., Caretta, C. A., Goñi-Urriza, M., Gassie, C., et al. (2014). Microbial diversity in Los Azufres geothermal field (Michoacán, México) and isolation of representative sulfate and sulfur reducers. *Extremophiles*, 18, 385–398.
- Brock, T. D. (1997). The value of basic research: discovery of *Thermus aquaticus* and other extreme thermophiles. *Genetics*, 146, 1207–1210.
- Brock, T. D., & Freeze, H. (1969). *Thermus aquaticus* gen. n. and sp. n., a non-sporulating extreme thermophile. *Journal of Bacteriology*, 98, 289–297.
- Crapart, S., Fardeau, M. L., Cayol, J. L., Thomas, P., Sery, C., Ollivier, B., et al. (2007). *Exiguobacterium profundum* sp. nov., a moderately thermophilic, lactic acid-producing bacterium isolated from a deep-sea hydrothermal vent. *International Journal of Systematic and Evolutionary Microbiology*, 57, 287–292.
- Ferris, M. J., & Ward, D. M. (1997). Seasonal distributions of dominant 16S rRNA-defined populations in a hot spring microbial mats examined by denaturing gradient gel electrophoresis. *Applied and Environmental Microbiology*, 63, 1375–1381.
- Hernández-León, R., Rojas-Solis, D., Contreras-Pérez, M., Orozco-Mosqueda, M. C., Macías-Rodríguez, L. I., Reyes-de la Cruz, H., et al. (2015). Characterization of the antifungal and plant growth-promoting effects of diffusible and volatile organic compounds produced by *Pseudomonas fluorescens* strains. *Biological Control*, 81, 83–92.
- Hou, W., Wang, S., Dong, H., Jiang, H., Briggs, B. R., Peacock, J. P., et al. (2013). A comprehensive census of microbial diversity in hot springs of Tengchong, Yunnan Province China using 16S rRNA gene pyrosequencing. *PLoS ONE*, 8, 1–15, e53350.
- Huang, Q., Dong, C. Z., Dong, R. M., Jiang, H., Wang, S., Wang, G., et al. (2011). Archaeal and bacterial diversity in hot springs on the Tibetan Plateau China. *Extremophiles*, 15, 549–563.
- Huang, Q., Jiang, H., Briggs, B. R., Wang, S., Hou, W., Li, G., et al. (2013). Archaeal and bacterial diversity in acidic to circum neutral hot springs in the Philippines. *FEMS Microbiology Ecology*, 85, 452–464.
- Israde-Alcántara, I., & Garduño-Monroy, V. H. (1999). Lacustrine record in a volcanic intra-arc setting: the evolution of the late Neogene Cuitzeo Basin System (Central Western Mexico, Michoacán). *Palaeogeography, Palaeoclimatology and Palaeoecology*, 151, 209–227.
- Janda, J. M., & Abbott, S. L. (2010). The genus *Aeromonas*: taxonomy, pathogenicity, and infection. *Clinical Microbiology Review*, 23, 35–73.
- Kanokratana, P., Chanapan, S., Pootanakit, K., & Eurwilaichitr, L. (2004). Diversity and abundance of Bacteria and Archaea in the Bor Khlung hot spring in Thailand. *Journal of Basic Microbiology*, 44, 430–444.
- Kimura, H., Sugihara, M., Yamamoto, H., Patel, B. K. C., Kato, K., & Hanada, S. (2005). Microbial community in a geothermal aquifer associated with the subsurface of the Great Artesian Basin Australia. *Extremophiles*, 9, 407–414.
- Klatt, C. G., Wood, J. M., Rusch, D. B., Bateson, M. M., Hamamura, N., Heidelberg, J. F., et al. (2011). Community ecology of hot spring cyanobacterial mats: predominant populations and their functional potential. *International Society for Microbial Ecology Journal*, 5, 1262–1278.
- Lacap, D. C., Barraquiu, W., & Pointing, S. B. (2007). Thermophilic microbial mats in a tropical geothermal location display pronounced seasonal changes but appear resilient to stochastic disturbance. *Environmental Microbiology*, 9, 3065–3076.
- Lau, M. C. Y., Aitchison, J. C., & Pointing, S. B. (2009). Bacterial community composition in thermophilic microbial mats from five hot springs in central Tibet. *Extremophiles*, 13, 139–149.
- López-Sandoval, Ó., Montejano, G., Carmona, J., Cantoral, E., & Becerra-Absalón, I. (2016). Diversidad algal de un ambiente extremo: el manantial geotermal Los Hervideros. *Revista Mexicana de Biodiversidad México*, 87, 1–9.
- Mackenzie, R., Pedrós-Alió, C., & Díez, B. (2013). Bacterial composition of microbial mats in hot springs in Northern Patagonia: variations with seasons and temperature. *Extremophiles*, 17, 123–136.
- Margesin, R., & Schinner, F. (2001). Biodegradation and bioremediation of hydrocarbons in extreme environments. *Applied Microbiology and Biotechnology*, 56, 650–663.
- Mead, D. A., Lucas, S., Copeland, A., Lapidus, A., Cheng, J. F., Bruce, D. C., et al. (2012). Complete genome sequence of *Paenibacillus* strain Y4.12MC10, a novel *Paenibacillus lautus* strain isolated from Obsidian Hot Spring in Yellowstone National Park. *Standard in Genomic Sciences*, 6, 366–385.
- Perevalova, A. A., Kolganova, T. V., Birkeland, N. K., Schleper, C., Bonch-Osmolovskaya, E. A., & Lebedinsky, A. V. (2008). Distribution of *Crenarchaeota* representatives in terrestrial hot springs of Russia and Iceland. *Applied and Environment Microbiology*, 74, 7620–7628.
- Portillo, M. C., Sirin, V., Kanoksilapatham, W., & González, J. M. (2009). Differential microbial communities in hot spring mats from western Thailand. *Extremophiles*, 13, 321–331.
- Purcell, D., Sompong, U., Yim, L. C., Barraclough, T. G., Peerapornpisal, Y., & Pointing, S. B. (2007). The effects of temperature, pH and sulphide on the community structure of hyperthermophilic streamers in hot springs of northern Thailand. *FEMS Microbiology Ecology*, 60, 456–466.
- Samal, S., Rout, G. R., Nayak, S., Nanda, R. M., Lenka, P. C., & Das, P. (2003). Primer screening and optimization for RAPD analysis of cashew. *Biologia Plantarum*, 46, 301–302.
- Shannon, C. E. (1948). A mathematical theory of communication. *Bell System Technical Journal*, 28, 42623–42656.
- Simpson, E. H. (1949). Measurement of diversity. *Nature*, 163, 688.
- Slepecky, R. A., & Hemphill, H. E. (2006). The genus *Bacillus* – nonmedical. In M. Dworkin, S. Falkow, E. Rosenberg, K. H. Schleifer, & E. Stackebrandt (Eds.), *Prokaryotes* (pp. 530–562). New York: Springer.
- Sørensen, T. (1948). A method of establishing groups of equal amplitude in plant sociology based on similarity of species and its application to analyses of the vegetation on Danish commons. *Biologiske Skrifter*, 5, 1–34.
- Tamura, K., Peterson, D., Peterson, N., Stecher, G., Nei, M., & Kumar, S. (2011). MEGA5: molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. *Molecular Biology and Evolution*, 28, 2731–2739.
- Tirawongsaroj, P., Sriprang, R., Harnpicharnchai, P., Thongaram, T., Champrada, V., Tanapongpipat, S., et al. (2008). Novel thermophilic and thermostable lipolytic enzymes from Thailand hot spring metagenomic library. *Journal of Biotechnology*, 133, 42–49.
- Tomova, I., Stoilova-Disheva, M., Lyutskanova, D., Pascual, J., Petrov, P., & Kambourova, M. (2010). Phylogenetic analysis of the bacterial community

- in a geothermal spring, Rupi Basin Bulgaria. *World Journal of Microbiology and Biotechnology*, 26, 2019–2028.
- Vázquez-Vázquez, M. J., Cortés-Martínez, R., & Alfaro-Cuevas, R. (2015). Arsenic occurrence and water quality in recreational thermal springs at Araro Mexico. *The International Journal of Science & Technology*, 5, 1–5.
- Viggiano-Guerra, J. C., & Gutiérrez-Negrín, L. C. A. (2005). The geothermal system of Araró Mexico, as an independent system of Los Azufres. In *Proceedings World Geothermal Congress* (pp. 24–29).
- Vishnivetskaya, T. A., Kathariou, S., & Tiedje, J. M. (2009). The *Exiguobacterium* genus: biodiversity and biogeography. *Extremophiles*, 13, 541–555.
- Ward, D. M., & Castenholz, R. W. (2000). Cyanobacteria in geothermal habitats. In B. A. Whitton, & M. Potts (Eds.), *Ecology of Cyanobacteria* (pp. 37–59). Amsterdam: Kluwer Academic Publishers.
- Ward, D. M., Ferris, M. J., Nold, S. C., & Bateson, M. M. (1998). A natural view of microbial biodiversity within hot spring cyanobacterial mat communities. *Microbiology and Molecular Biology Reviews*, 62, 1353–1370.
- Weisburg, W. G., Barns, S. M., Pelletier, D. A., & Lane, D. J. (1991). 16S Ribosomal DNA amplification for phylogenetic study. *Journal of Bacteriology*, 173, 697–703.
- Yakimov, M. M., Timmis, K. N., Wray, V., & Fredrickson, H. L. (1995). Characterization of a new lipopeptide surfactant produced by thermotolerant and halotolerant subsurface *Bacillus licheniformis* BAS50. *Applied and Environment Microbiology*, 61, 1706–1713.
- Yim, L. C., Hongmei, J., Aitchison, J. C., & Pointing, S. B. (2006). Highly diverse community structure in a remote central Tibetan geothermal spring does not display monotonic variation to thermal stress. *FEMS Microbiology Ecology*, 57, 80–91.