



MEDICC Review

ISSN: 1555-7960

editors@medicc.org

Medical Education Cooperation with Cuba
Estados Unidos

Varona, Patricia; Bonet, Mariano; García, René; Chang, Martha; Suárez, Ramón;
Coordinating Group
Implementation of Chronic Disease Risk Factor Surveillance in 12 Cuban Municipalities
MEDICC Review, vol. 16, núm. 1, 2014, pp. 43-47
Medical Education Cooperation with Cuba
Oakland, Estados Unidos

Available in: <http://www.redalyc.org/articulo.oa?id=437542096009>

- How to cite
- Complete issue
- More information about this article
- Journal's homepage in redalyc.org

redalyc.org

Scientific Information System
Network of Scientific Journals from Latin America, the Caribbean, Spain and Portugal
Non-profit academic project, developed under the open access initiative

Implementation of Chronic Disease Risk Factor Surveillance in 12 Cuban Municipalities

Patricia Varona MD MS, Mariano Bonet MD, René García MD MS, Martha Chang MS, Ramón Suárez MD; National–Provincial Coordinating Group*

SUMMARY

Chronic non-communicable diseases have been called the pandemic of the 21st century and constitute a high-priority public health challenge; hence growing interest in chronic disease risk factor surveillance. Cuba is implementing decentralized risk factor surveillance in each of its municipalities as part of a strategy to address non-communicable diseases. Decentralized surveillance with this level of detail and explicitly designed to inform municipal and provincial decisionmaking is unprecedented in Cuba. We describe the methodology for planning and implementing measurement of major risk factors in 12 municipalities in 10 provinces, as part of Cuba's National Surveillance System. The results have facilitated timely use of information and evidence-based decisionmaking at the local level.

KEYWORDS: Surveillance system, non-communicable disease risk factors, health surveys, survey methods, Cuba

INTRODUCTION

Chronic non-communicable diseases (NCD) have been called the pandemic of the 21st century and constitute a high priority for public health. It is vitally important to improve health systems' capacity to respond, making better use of preventive potential by reducing prevalence of specific risk factors and conditions: tobacco and alcohol use, unhealthy diet, physical inactivity, obesity, diabetes, high serum cholesterol, hypertension, and physical and chemical factors in the environment;[1,2] hence growing interest in NCD risk factor surveillance.

Globally, NCDs and their risk factors are important and growing contributors to the burden of morbidity, mortality and disability, and to diminished quality of life for patients and their families. NCDs contribute to 60% of all deaths, and of these, 46% are in the population aged <70 years (the most productive years), to the detriment of countries' development potential; 80% of NCD deaths are in low- and middle-income countries.[3–5]

In the Americas, 62% (3.2 million) of the 5.2 million deaths reported in 2004 resulted from NCDs, mainly cardiovascular diseases, cancer, chronic respiratory diseases and diabetes. Given current demographic, social and economic changes, as well as epidemiological and nutritional shifts, NCD-attributable deaths are expected to increase in the coming years.[2,3,6] At the same time, a growing body of evidence indicates that NCDs are largely preventable, including assessment of effectiveness, cost, acceptability and feasibility of preventive interventions. Thus, it is estimated that improved diet, more physical activity and eliminating tobacco use could prevent 80% of heart disease and 90% of diabetes cases.[7–10]

In Cuba, NCDs are involved in 76% of all deaths and annual NCD-related mortality (642.2 per 100,000 population) is over ten times the combined rates for communicable diseases and maternal, perinatal and nutritional causes (58.7 per 100,000

population); NCDs add greatly to potential years of life lost and impact negatively on individual quality of life and on the Cuban economy.[11]

National policy statements and other documents have expressed awareness of the importance of NCD risk factors since 1981. A commitment to address them was ratified in the 1992 document *Objectives, Principles and Directives for Cuban Public Health for the Year 2000*, which pointed to the need for surveillance and research on the major risk factors associated with population lifestyles. Thus, the first National Survey on Risk Factors and Chronic Diseases was conducted in 1995 to identify prevalence and distribution of the main NCD risk factors in Cuba's urban population aged ≥15 years. Its results provided a baseline that has facilitated trend detection and evaluation of interventions. In 1996 Cuba's Ministry of Public Health (MINSAP, the Spanish acronym) established the National Chronic Non-Communicable Diseases Program, with a strong foundation in primary care, hospitals and provincial hygiene and epidemiology centers. The Program combines comprehensive health care actions and collaboration with other sectors.[12,13]

The Second National Survey on Risk Factors and Chronic Diseases (ENFRENT II, the Spanish acronym) was completed in late 2001. For the first time, it included a section on injuries and illnesses and another on health-related quality of life. In 2005–2006, health projections for 2015 were approved, which emphasized reaching a life expectancy at birth of 80 years. The interventions proposed to meet that goal have been affected in recent years by a series of natural disasters resulting in epidemics of communicable diseases (influenza, dengue and others), which affected the National Program's sustainability.

In 2007, the need to reassess the NCD situation was recognized[14] and the decision taken to implement a different, decentralized surveillance strategy capable of identifying distribution and trends of major risk factors at the municipal level. This in turn would enable closer study, assessment and management of NCDs and their risk factors and identify new opportunities for intervention and future research at the local level. This would all, of course, have to be consistent with Cuba's public health objectives and carried out in line with MINSAP's policy of decentralization.[15,16]

This paper describes planning and implementation of NCD risk factor measurement as part of Cuba's decentralized surveillance system in 12 municipalities during the years 2008–2010.

METHODS

A descriptive cross-sectional study was designed and implemented from 2008 through 2010. The study's target population consisted of persons aged 15–74 years resident in urban and rural areas in 17 selected municipalities throughout Cuba.

Lessons from the Field

Municipalities were selected based on their economic, social or cultural influence on the life of the province, as well as three municipalities with previous NCD-related health intervention projects. The total population in these municipalities is 28.3% of the Cuban population of approximately 11.2 million.[11]

The framework was based on WHO/PAHO *Stepwise Approach to Non-Communicable Disease Risk Factor Surveillance* (STEPS), which can be adapted to specific economic, infrastructure and logistical conditions. Cuba implemented the instrument's three steps: questionnaire (step 1), physical measurements (step 2), and biochemical measurements (step 3).[17] The study was led by the National Hygiene, Epidemiology and Microbiology Institute (INHEM, the Spanish acronym), in conjunction with MINSAP experts in health education and research, and with the National Statistics Bureau. A national coordinating team was formed with expertise in NCD control and population surveys. The process was replicated in each province to ensure appropriate oversight and implementation. This was not possible in every municipality because of reassignment of some personnel to other tasks, in Cuba and abroad.

Use of the STEPS questionnaire permitted comparison of resulting data with those from other countries and also within Cuba over time. The questionnaire was adapted to the Cuban context and expanded to include questions about quality of life. Serum creatinine was added to the biochemical measures to determine renal function.[17]

The instrument included:

- general information—address, health care catchment area;
- sociodemographic information—sex, age, education level completed (no formal schooling, primary school, middle school, trades school, middle-level technical training, high school and university), skin color, marital status, main employment, and wages earned in the previous year;
- behavioral measurements—tobacco and alcohol use; diet; physical activity at work, in commuting and leisure activities;
- personal history of diabetes, high total cholesterol and hypertension;
- family history of diabetes, high cholesterol, hypertension, ischemic or hemorrhagic stroke, heart attack and cancer;
- physical measurements—blood pressure and anthropometrics (weight, height, and waist and hip circumference);
- biochemical measurements—fasting blood glucose, triglycerides, total cholesterol and creatinine;
- health-related quality of life—EuroQol instrument using a visual analogue scale for self report of five aspects of health status (mobility, self care, usual activities, pain or discomfort, and anxiety or depression).[18]

Constraints Due to difficulties obtaining a reliable supply of reagents, biochemical measurements were not taken consistently in all municipalities. There were also challenges to determining wages earned by those surveyed, so that part of the analysis was postponed. Several natural disasters hit Cuba and spawned outbreaks of communicable diseases, preventing simultaneous collection of data in all municipalities. The present paper therefore includes only the 12 municipalities where data analysis was complete at the time of writing. These problems may also influence variation in municipal results, which will require local-level analysis.

IMPLEMENTATION

Study plan and implementation In all provinces, meetings were held to provide detailed explanations of the study and local use of its results to provincial (People's Power president or vice president) and municipal governments. In each municipality, five working groups of health personnel (technicians, nurses, doctors and administrative staff) were constituted, with defined tasks:

- data collection (ten surveyors and two supervisors);
- physical measurements: anthropometrics and blood pressure (ten people, including two supervisors);
- biochemical measurements (three people);
- data management (data entry clerks, epidemiologists and biostatisticians); and
- logistics, including resources and survey organization.

The Provincial Hygiene, Epidemiology and Microbiology Centers (CPHEM, the Spanish acronym), municipal hygiene and epidemiology centers and units, and head nurses in the selected provinces, municipalities and polyclinics worked throughout the organizational phase.

Sample design In each municipality an independent sample was taken, although the design was common to all (except for the municipality of Cienfuegos, which participated in the international project CARMELA [Cardiovascular Risk Factor Multiple Evaluation in Latin America], whose sampling design and basic criteria have been published).[19]

The sample size required was calculated according to expert criteria (see acknowledgments for technical contributions) of approximately 200 individuals in each stratum (age group and sex). To ensure this sample size, allowing for attrition, 220 individuals per stratum were selected. The design, therefore, was equiprobabilistic within each group but not between groups.

Cluster sampling was performed in two stages, stratified by age and sex. Clusters consisted of family doctor-and-nurse practices in the municipality's health areas; units of analysis were the individuals selected in each practice. There were 12 strata: 6 age groups (15–24; 25–34; 35–44; 45–54; 55–64; 65–74 years) for each sex.

In the first stage, family doctor-and-nurse practices were randomly selected from among all practices in each municipality through systematic sampling; in these practices, a census was conducted of all persons in the age groups of interest. The second stage involved a systematic selection of 220 individuals in each stratum, using fractions equal to 220/number of individuals in the stratum, for a final desired sample size of 2640 for each municipality (with the exception of Cienfuegos, noted above).

Since the sampling was complex (multistaged, stratified and non-equiprobabilistic), a weighting factor was required for data analysis. It consisted of the inverse of the probability of selecting an individual and was calculated by dividing, for each age and sex group, the 2008 population in the municipality by the total number of persons in the sample, so that estimates were representative of all persons in that age and sex group in the municipality.[20]

By the time this paper was being prepared, data processing and analysis had concluded in 12 municipalities, whose populations together comprise 20% of the country's total population (Table 1).

Table 1: Sample characteristics by municipality (n = 2640*)

Municipality	Population	Sample achieved n (%)	Sex distribution		Mean age (years) (95% CI)
			Male n (%)	Female n (%)	
Pinar del Río	191,594	2608 (98.8)	1226 (47.0)	1382 (53.0)	41.1 (39.4–42.8)
San Antonio de los Baños	49,644	2614 (99.0)	1265 (48.4)	1349 (51.6)	39.8 (37.5–42.1)
Matanzas	147,708	2640 (100.0)	1320 (50.0)	1320 (50.0)	41.4 (39.5–42.5)
Cienfuegos	32,114	2193* (91.4)	933 (42.5)	1260 (57.5)	44.6 (43.9–45.3)
Aguada	166,651	2178 (82.5)	1074 (49.3)	1104 (50.7)	41.0 (39.9–42.1)
Santa Clara	236,758	2054 (77.8)	854 (41.6)	1200 (58.4)	42.3 (38.5–46.1)
Yaguajay	57,508	2331 (88.3)	1144 (49.1)	1187 (50.9)	41.8 (39.6–44.1)
Morón	139,508	2469 (93.5)	1192 (48.3)	1277 (51.7)	39.8 (37.5–42.2)
Ciego de Ávila	62,672	2456 (93.0)	1184 (48.2)	1272 (51.8)	40.1 (37.7–42.5)
Camagüey	326,211	2552 (96.7)	1238 (48.5)	1314 (51.5)	40.8 (38.4–43.3)
Holguín	334,299	2141 (81.1)	811 (37.9)	1330 (62.1)	40.8 (37.6–43.9)
Santiago de Cuba	495,574	2397 (91.8)	1064 (44.4)	1333 (55.6)	40.4 (39.1–41.8)

*Desired sample for Cienfuegos was 2400[20]

Data collection As explained above, for various reasons, it was not possible to collect data in all municipalities at the same time. Thus, in municipalities in eastern Cuba and one in the central region, the data collection phase extended from 2008 through June 2010.

Before starting the survey proper, a pilot test was conducted in the 10 de Octubre Municipality in Havana Province (then called City of Havana Province, prior to provincial boundary changes in 2010), to estimate questionnaire completion time and assess comprehension and acceptability. Of the 100 people who were randomly selected, 94 agreed to participate and 6 declined, citing time limitations.

Participation in the survey proper involved a 50–60 minute formal interview, during which respondents were asked to go to the measurement center 48 hours later (or at an agreed-upon time) in a fasting state (written instructions provided). Measurement centers were usually family doctor-and-nurse offices, schools or other suitable buildings in the community, chosen to meet minimal conditions of privacy and comfort and for location as close as possible to the study area, that is, the catchment area of the family doctor-and-nurse office.

Family doctors and nurses facilitated recruitment and trained study personnel in each measurement center, took physical measurements per protocol, and drew blood samples for laboratory testing for glucose, creatinine, and total cholesterol and triglycerides.[21]

Ethics All persons selected were visited at home to obtain written consent to participate, following explanation of the purpose of the survey, importance and benefits of participation, absence of risk,

data confidentiality and freedom to opt out without repercussions at any moment should they so desire.

Special care was taken to send participants' test results via their family doctor and nurse; if any changes requiring medical attention were observed, participants were immediately referred for treatment and followup.

Quality control A system of quality control was set in place from the beginning, including detailed protocols and standard procedures for questionnaire administration and anthropometric and laboratory measurements. Three-day sessions were held in 14 provinces to train personnel designated for each of the three functions. A rigorous accreditation step was also set up for quality assurance.

The national coordinating group visited provinces to verify quality on the ground through direct observation of protocol adherence. In two municipalities, the process was interrupted because of organizational problems and reinitiated later when organization met required standards.

Laboratory results were monitored weekly by a national reference laboratory in Havana (at Carlos J. Finlay Biological Products Company). Some 438 health technicians were involved in analyzing samples, which helped increase their scientific–technical preparation in NCD risk factors.

Data management and statistical analysis Data entry used EpiData Version 3.1 software. In each municipality, a supervisor reviewed and then coded data collected daily. At the end of each work day, provincial coordinating team supervisors sent the database by email to biostatisticians in the INHEM national coordinating team, where they again reviewed and corrected the minimum, average and maximum values of each variable. A central database was consolidated for each municipality with an automated validation system for each variable in the questionnaire.

Data were analyzed in each municipality. Means were calculated for continuous variables and percentages for qualitative ones, with 95% confidence intervals, using methods recommended for complex sample analysis, including standard error adjustments.[20]

The population's sociodemographic characteristics are presented in Table 2. This were consistent with 2002 census data.[22]

LESSONS LEARNED

The planning and implementation processes described here were unprecedented in Cuba and enabled decentralization of NCD risk factor surveillance and prompt use of information generated for decisionmaking in municipalities and provinces. From the research team's perspective, this is the most important result obtained.

Lessons from the Field

Table 2: Sociodemographic distribution of sample*

Variable	Municipality											
	PDR n (%)	SAB n (%)	MTZ n (%)	CFG n (%)	AGU n (%)	SCL n (%)	YAG n (%)	MOR n (%)	CAV n (%)	CAM n (%)	HOL n (%)	STG n (%)
Skin Color												
White	2046 (78.8)	1786 (68.5)	2017 (76.4)	1585 (72.3)	1732 (79.6)	1727 (84.8)	1934 (83.0)	2025 (82.6)	2016 (82.6)	1880 (73.7)	1685 (78.8)	689 (28.7)
Mestizo	247 (9.5)	533 (20.4)	341 (12.9)	428 (19.5)	281 (12.9)	202 (9.9)	192 (8.2)	238 (9.7)	236 (9.7)	438 (17.2)	330 (15.4)	1165 (48.6)
Black	306 (11.8)	289 (11.1)	282 (10.7)	180 (8.2)	164 (7.5)	107 (5.3)	205 (8.8)	189 (7.7)	187 (7.7)	234 (9.2)	122 (5.7)	543 (22.7)
Missing data	9	6	0	0	1	18	0	17	17	0	4	0
Total	2608	2614	2640	2193	2178	2054	2331	2469	2456	2552	2141	2397
Education completed												
None	147 (5.6)	45 (1.7)	39 (1.5)	0 (0.0)	60 (2.8)	49 (2.4)	106 (4.5)	131 (5.3)	131 (5.3)	133 (5.2)	82 (3.8)	46 (1.9)
Primary school	346 (13.3)	358 (13.7)	254 (9.6)	194 (8.8)	471 (21.7)	165 (8.1)	300 (12.9)	247 (10.0)	245 (10.0)	253 (9.9)	233 (10.9)	218 (9.1)
Middle school	595 (22.8)	734 (28.1)	645 (24.4)	565 (25.8)	628 (28.9)	520 (25.5)	723 (31.0)	604 (24.5)	600 (24.5)	629 (24.6)	474 (22.2)	493 (20.6)
Trade school	53 (2.0)	56 (2.1)	51 (1.9)	0 (0.0)	67 (3.1)	59 (2.9)	57 (2.4)	39 (1.6)	38 (1.6)	53 (2.1)	29 (1.4)	63 (2.6)
Technical training	377 (14.5)	751 (28.7)	563 (21.3)	376 (17.1)	296 (13.6)	435 (21.3)	409 (17.5)	460 (18.7)	456 (18.6)	307 (12.0)	338 (15.8)	507 (21.1)
High school	639 (24.5)	386 (14.8)	615 (23.3)	662 (30.2)	480 (22.1)	406 (19.9)	498 (21.4)	584 (23.7)	582 (23.8)	902 (35.3)	496 (23.2)	632 (26.4)
University	448 (17.2)	284 (10.9)	473 (17.9)	396 (18.1)	170 (7.8)	407 (19.9)	238 (10.2)	396 (16.1)	396 (16.2)	275 (10.8)	484 (22.7)	438 (18.3)
Missing data	3	0	0	0	6	13	0	8	8	0	5	0
Total	2608	2614	2640	2193	2178	2054	2331	2469	2456	2552	2141	2397
Main employment												
Public sector	1241 (48.1)	1372 (52.6)	1429 (54.1)	1116 (51.0)	945 (43.4)	935 (47.3)	1142 (49.0)	1040 (44.1)	1034 (44.1)	1453 (56.9)	1063 (50.0)	1188 (49.6)
Joint venture	35 (1.4)	58 (2.2)	91 (3.4)	17 (0.8)	38 (1.7)	18 (0.98)	3 (0.1)	62 (2.6)	62 (2.6)	13 (0.5)	53 (2.5)	18 (0.7)
Self employed	115 (4.4)	95 (3.6)	116 (4.4)	98 (4.5)	194 (8.9)	65 (3.3)	357 (15.3)	120 (5.1)	120 (5.1)	96 (3.8)	92 (4.3)	62 (2.6)
Student	233 (9.0)	171 (6.6)	243 (9.2)	160 (7.3)	109 (5.0)	98 (5.0)	91 (3.9)	170 (7.2)	168 (7.2)	172 (6.7)	92 (4.3)	204 (8.5)
Homemaker	297 (11.5)	270 (10.4)	233 (8.8)	307 (14.0)	542 (24.9)	292 (14.8)	354 (15.2)	379 (16.1)	377 (16.1)	391 (15.3)	365 (17.2)	320 (13.3)
Retired	550 (21.3)	557 (21.43)	455 (17.2)	374 (17.1)	303 (13.9)	510 (25.8)	376 (16.1)	477 (20.2)	475 (20.2)	402 (15.7)	418 (19.7)	492 (20.5)
Unemployed	111 (4.3)	84 (3.2)	73 (2.8)	114 (5.2)	46 (2.1)	59 (3.0)	8 (0.3)	111 (4.7)	110 (4.7)	25 (1.0)	41 (1.9)	113 (4.7)
Missing data	26	7	0	7	1	77	0	110	110	0	17	0
Total	2608	2614	2640	2193	2178	2054	2331	2469	2456	2552	2141	2397
Marital Status												
Single	808 (31.1)	572 (22.1)	663 (25.1)	596 (27.2)	464 (21.3)	407 (20.3)	368 (15.8)	634 (26.4)	632 (26.5)	600 (23.5)	400 (18.9)	817 (34.1)
Married	1110 (42.7)	1140 (44.1)	1103 (41.8)	932 (42.5)	1185 (54.4)	1036 (51.6)	982 (42.1)	888 (37.0)	882 (36.9)	1091 (42.8)	953 (44.9)	807 (33.7)
Common law	305 (11.7)	549 (21.3)	516 (19.5)	370 (16.9)	265 (12.2)	289 (14.4)	693 (29.7)	614 (25.6)	612 (25.6)	562 (22.0)	420 (19.8)	543 (22.6)
Widowed	153 (5.9)	127 (4.9)	133 (5.0)	107 (4.9)	115 (5.3)	103 (5.1)	95 (4.1)	119 (5.0)	119 (5.0)	99 (3.9)	103 (4.8)	100 (4.2)
Separated or divorced	222 (8.5)	195 (7.5)	225 (8.5)	188 (8.6)	149 (6.8)	173 (8.6)	193 (8.3)	145 (6.0)	142 (5.9)	200 (7.8)	244 (11.5)	130 (5.4)
Missing data	10	31	0	0	0	46	0	69	69	0	21	0
Total	2608	2614	2640	2193	2178	2054	2331	2469	2456	2552	2141	2397


*Percentages were calculated based on net totals (without missing data)

AGU: Aguada CAM: Camagüey CAV: Ciego de Ávila CFG: Cienfuegos HOL: Holguín MOR: Morón MTZ: Matanzas PDR: Pinar del Río
SAB: San Antonio de los Baños SCL: Santa Clara STG: Santiago de Cuba YAG: Yaguajay

From the outset, participation of local government authorities was decisive in the process and for establishing accountability for progress in measurements, to ensure timely corrections if necessary.

The survey's success depended in part on discussion and analysis of data with senior provincial public health authorities and representatives of provincial and municipal governments, as well as members of the National–Provincial Coordinating Group. The results of these

discussions informed design of a concrete intersectoral action plan with emphasis on a communications strategy. Participating sectors and institutes included those in education, food and nutrition, sports and recreation, culture, higher education, and the media.

This experience demonstrates the usefulness and feasibility of intersectoral work in implementing NCD risk factor surveillance, as has been found in other countries.[23–25] 

REFERENCES

- World Health Organization. Prevention and control of noncommunicable diseases: implementation of the global strategy. Resolution WHA60.23 [Internet]. Geneva: World Health Organization; 2007 [cited 2012 Jan 30]. Available from: http://www.who.int/gb/eb_e_wha60.html
- World Health Organization. Preventing chronic diseases: a vital investment [Internet]. Geneva: World Health Organization; 2005 [cited 2012 Mar 29]. Available from: http://whqlibdoc.who.int/publications/2005/9241563001_eng.pdf
- Pan American Health Organization. Enfermedades No Transmisibles en las Américas: construyamos un futuro más saludable. Washington: Pan American Health Organization; 2011. Spanish.
- Mathers C, Loncar D. Projections of global mortality and burden of disease from 2002 to 2030. PLoS Med [Internet]. 2006 [cited 2012 Feb 3];3:e442. Available from: <http://10.1371/journal.pmed.0030442> PMID:17132052
- World Health Organization. Global health risks: mortality and burden of disease attributable to selected major risks. Geneva: World Health Organization; 2009.
- PAHO/WHO. Regional Health Observatory. Americas 2008 [Internet]. Washington: Pan American Health Organization; 2010 [cited 2012 Jul 5]. Available from: http://new.paho.org/hq/index.php?option=com_content&view=category&layout=blog&id=2395&Itemid=2523&lang=en
- Beaglehole R, Ebrahim S, Reddy S, Voûte J, Leeder S; Chronic Disease Action Group. Prevention of chronic diseases: a call to action. Lancet. 2007 Dec 22;370(9605):2152–7.
- World Health Organization. 2008–2013 Action Plan for the Global Strategy for the Prevention and Control of Noncommunicable Diseases [Internet]. Geneva: World Health Organization; 2008 [cited 2012 May 12]. Available from: http://whqlibdoc.who.int/publications/2009/9789241597418_eng.pdf
- Yusuf S, Hawken S, Ounpuu S, Dans T, Avezum A, Lanas F. Effect of potentially modifiable risk factors associated with myocardial infarction in 52 countries (the INTERHEART study): case-control study. Lancet. 2004 Sep 11–17;364(9438):937–52.
- World Health Organization. Facts related to chronic diseases. Geneva: World Health Organization; c2013 [cited 2006 Jan 20]. Available from: http://www.who.int/topics/chronic_diseases/factsheets/en/index.html
- Ministry of Public Health (CU). Anuario Estadístico 2011. Havana: Ministry of Public Health (CU); 2012. Spanish.
- Ministry of Public Health (CU). Objetivos, Propósitos y Directrices de la Salud cubana para el año 2000. Havana: Ministry of Public Health (CU); 2002. Spanish.
- National Institute of Hygiene, Epidemiology and Microbiology (CU). Informe Técnico. Primera Encuesta Nacional de Factores de Riesgo de Enfermedades no Transmisibles. Havana: National Institute of Hygiene, Epidemiology and Microbiology (CU); 1995. Spanish.
- National Institute of Hygiene, Epidemiology and Microbiology (CU). Informe Técnico. Vigilancia de los Factores de riesgo de enfermedades no transmisibles. Cuba, 2007. Havana: National Institute of Hygiene, Epidemiology and Microbiology (CU); 2007. Spanish.
- National Institute of Hygiene, Epidemiology and Microbiology (CU). Informe Técnico. Segunda Encuesta Nacional de Factores de Riesgo de Enfermedades no Transmisibles y Actividades Preventivas, 2001. Havana: National Institute of Hygiene, Epidemiology and Microbiology (CU); 2001. Spanish.
- Ministry of Public Health (CU). Proyecciones de la Salud Cubana hasta el año 2015. Havana: Ministry of Public Health (CU); 2006. Spanish.
- Programa STEPS Panamericano. Método “paso a paso” de la OPS/OMS para la vigilancia de factores de riesgo para las enfermedades crónicas no transmisibles. Washington: Pan American Health Organization; 2008 [cited 2012 Jan 4]. Available from: <http://www.paho.org/spanish/ad/dpc/nc/panam-steps.htm>. Spanish.
- Herdman H, Badia X, Berra S. El EuroQol-5D: una alternativa sencilla para la medición de la calidad de vida relacionada con la salud en la atención primaria. Aten Primaria. 2001;28(6):425–9. Spanish.
- Silva LC, Benet M, Morejón A, Orduñez P. An Efficient Sampling Approach to Surveillance of Non-Communicable Disease Risk Factors in Cienfuegos, Cuba. MEDICC Rev. 2012;14(4):36–9.
- Cañizares M, Barroso I, Alfonso A, García R, Alfonso K, Chang M, et al. Estimaciones usadas en diseños muestrales complejos: Aplicaciones en la encuesta de salud cubana en el año 2001. Rev Panam de Salud Pública. 2004;15(3):176–84. Spanish.
- O'Brien E, Waber B, Parati G, Staessen J, Myers MG; the European Society of Hypertension Working Group on Blood Pressure Monitoring. Blood pressure measuring devices: recommendations of the European Society of Hypertension. BMJ. 2001 Mar 3;322(7285):531–6.
- National Statistics Bureau (CU). Censo de población y viviendas, 2002. Havana: National Statistics Bureau (CU); 2003. Spanish.
- State and Society Studies Center (AR). Encuesta de Factores de Riesgo para Enfermedades No Transmisibles. Adaptación Transcultural y Validación. Buenos Aires: CEDES/Área de Salud, Economía y Sociedad; 2010. Spanish.
- Ministry of Health (CL). Encuesta Nacional de Salud de Chile 2009, 2010. Tomo1. Santiago de Chile: Ministry of Health (CL), Pontificia Universidad Católica de Chile/Universidad Alberto Hurtado; 2011. Spanish.
- Ministry of Health (CR). Encuesta de factores de riesgo para enfermedades no transmisibles. CARTAGO 2000. Módulo Tabaco y Alcohol. Costa Rica: Ministry of Health (CR), INCIENSA/IAFA; 2002. Spanish.

THE AUTHORS

Patricia Varona Pérez (Corresponding author: patricia.varona@infomed.sld.cu), physician specializing in epidemiology with a master's degree in environmental health. Associate professor and researcher, National Hygiene, Epidemiology and Microbiology Institute (INHEM), Havana, Cuba.

Mariano Bonet Gorbea, physician specializing in epidemiology. Associate professor and researcher, INHEM, Cuba.

René García Roche, physician specializing in medical informatics. Associate professor and adjunct researcher, INHEM, Cuba.

Martha Chang La Rosa, sociologist with a master's degree in public health, INHEM, Cuba.

Ramón Suárez Medina, physician specializing in biostatistics, INHEM, Cuba.

*Ma. Elena Díaz Sánchez PhD, Delia Herrera Travieso MD, Dianelys Pérez Jiménez, Alain Morejón Giraldo MD, Santa Jiménez Acosta MD PhD, Silvia Venero Fernández MD, Yaima Fernández Díaz, Rosamarina García Pérez MD, Geominia Maldonado Castillo MD, Héctor Hernández Hernández MD, Karen Alfonso Sagué MD, Mikhail Benet Rodríguez MD PhD, Orlando Landrove Rodríguez MD, Rodolfo Jova Morel MD, Alicia Almeyda Deulofeu MD, Odalys Urraca Castillo MD, Loida A. Borrego Alvarez MD, José Morales Rigau MD, Arelys Dorta Delgado MD, Dania Pedraza Alejo MD, Heidy Rodríguez MD, Marelys Cendra Asencio MD, Juan Núñez Chacón MD, Luis Benítez Guerrero MD, Mileideis Toselló López MD, Marlenis Brosard Cisneros MD, Isabel Carbonell MD, Francisca D. Gómez Torres

Submitted: September 6, 2012

Approved for publication: December 31, 2013

Disclosures: None