



Lankesteriana International Journal on  
Orchidology

ISSN: 1409-3871

lankesteriana@ucr.ac.cr

Universidad de Costa Rica  
Costa Rica

Wilson, Mark; Jost, Lou

Phylogenetic analysis of the Andean genus *Brachycladium* Luer (syn. *Oreophilus* Higgins  
& Archila) and closely related genera based on nuclear ITS sequencing

Lankesteriana International Journal on Orchidology, vol. 11, núm. 3, diciembre, 2011

Universidad de Costa Rica  
Cartago, Costa Rica

Available in: <http://www.redalyc.org/articulo.oa?id=44339822023>

- How to cite
- Complete issue
- More information about this article
- Journal's homepage in redalyc.org

redalyc.org

Scientific Information System

Network of Scientific Journals from Latin America, the Caribbean, Spain and Portugal

Non-profit academic project, developed under the open access initiative

# PHYLOGENETIC ANALYSIS OF THE ANDEAN GENUS *BRACHYCLADIUM* LUER (SYN. *OREOPHILUS* HIGGINS & ARCHILA) AND CLOSELY RELATED GENERA BASED ON NUCLEAR ITS SEQUENCING

MARK WILSON<sup>1,3</sup> & LOU JOST<sup>2</sup>

<sup>1</sup> Biology Department, The Colorado College, Colorado Springs, CO 80903, USA

<sup>2</sup> Via a Runtun, Baños, Tungurahua, Ecuador

<sup>3</sup> Corresponding author: mwilson@coloradocollege.edu

We previously reported that nrITS sequencing and phylogenetic analysis revealed at least four cryptic species within *Brachycladium nummularium* (syn. *Oreophilus nummularius*). Sequencing of additional species in the genus revealed two major clades within *Brachycladium*, the “nummularium”-complex plus *B. stalactites* and Luer’s section *Amplectentes*. Since the relationship of *Brachycladium* to *Andinia* had been noted earlier, a number of *Andinia* species were also included. Two clades were observed, a clade containing *A. lappacea*, *A. pensilis*, and *A. vestigipetalum* that was closely related to *Brachycladium* and a second, more distantly related clade containing *A. dalstroemii*, *A. pogonion*, and *A. schizopogon*.

Samples of two additional genera, *Masdevalliantha* Szlachetko & Margonska and *Xenosia* Luer, were included in the analysis based upon morphological similarity to *Andinia*. These two genera formed a clade that was sister to both *Brachycladium* and *Andinia*. Only the node subtending clades of all four genera had a high level of bootstrap support, whereas nodes subtending clades of *Brachycladium* plus *Andinia*, or *Brachycladium* alone, had much lower support. If future chloroplast sequencing produces phylogenies congruent with nrITS, this would indicate that *Andinia*, *Brachycladium* (syn. *Oreophilus*), *Masdevalliantha*, and *Xenosia* should all be placed in the same genus.