



Lankesteriana International Journal on
Orchidology

ISSN: 1409-3871

lankesteriana@ucr.ac.cr

Universidad de Costa Rica
Costa Rica

Dalström, Stig

A NEW AND PREVIOUSLY MISIDENTIFIED CYRTOCHILUM (ORCHIDACEAE:
ONCIDIINAE) FROM THE HIGH PLAINS OF CENTRAL ECUADOR

Lankesteriana International Journal on Orchidology, vol. 15, núm. 1, abril, 2015, pp. 101-
105

Universidad de Costa Rica
Cartago, Costa Rica

Available in: <http://www.redalyc.org/articulo.oa?id=44339830008>

- How to cite
- Complete issue
- More information about this article
- Journal's homepage in redalyc.org

redalyc.org

Scientific Information System

Network of Scientific Journals from Latin America, the Caribbean, Spain and Portugal

Non-profit academic project, developed under the open access initiative

A NEW AND PREVIOUSLY MISIDENTIFIED *CYRTOCHILUM* (ORCHIDACEAE: ONCIDIINAE) FROM THE HIGH PLAINS OF CENTRAL ECUADOR

STIG DALSTRÖM

2304 Ringling Boulevard, unit 119, Sarasota FL 34237, USA
Lankester Botanical Garden, University of Costa Rica, Cartago, Costa Rica
National Biodiversity Centre, Serbithang, Bhutan
stigdalstrom@gmail.com

ABSTRACT. A new and previously misidentified, small-flowered *Cyrtochilum* (Orchidaceae: Oncidiinae) from central Ecuador is described, illustrated with a line drawing and photographs, and compared with the species that it has previously been taxonomically mixed-up with by the author. The new species is readily distinguished by the dark yellow, gibbous and carnoselip callus, versus more elongate, longitudinally furrowed and bilobed calli for similar species.

KEY WORDS: *Cyrtochilum*, Orchidaceae, Oncidiinae, new species, Ecuador, Azuay

Introduction. The *Cyrtochilum* species described here has previously been misidentified by the author as “*Cyrtochilum viminale* (Rehb.f.) Dalström” (2001, 2010). However, that name refers to a different species with a different floral morphology. Our new, but well-known species represents an enigmatic paradox. It is very common along roads in the south-central highlands of Ecuador, particularly around the old city of Cuenca. Plants can sometimes be found growing on the road itself, clinging tenaciously to the dirt while the relentless traffic constitutes a permanent threat to crush the plants. Anybody that passes by and has an interest in orchids will sooner or later notice the flowers and stop to investigate, and often make herbarium specimens. The peculiar thing, however, is that our new *Cyrtochilum* is most rarely encountered in herbaria from older collections. The reason for this may have an interesting explanation. It seems plausible that before modern transportation entered the stage (and the Ecuadorian wilderness) this species occurred only as terrestrials in remote, scrub and grass vegetation, not seen by anybody except occasional natives passing by. Professional collectors probably avoided this seemingly ‘empty’ land that did not seem to host any commercial rewards. As roads began to crisscross these relatively flat and engineering friendly areas, however, the disturbed road cuts offered suitable habitats for the orchid.

In addition, passing vehicles loaded with livestock, plants and miscellaneous equipment may have helped dispersing the seeds, with new populations establishing along the roads as a consequence.

TAXONOMIC TREATMENT

Cyrtochilum soennemarkii Dalström *sp. nov.*

TYPE: Ecuador. Azuay, km 52 from Cuenca towards Loja, in full sun along roadside, at 3250 m, 13 Dec. 1982, S. Dalström 354 (holotype: SEL). Figs. 1, 2.

Diagnosis. *Cyrtochilum soennemarkii* (Figs. 1–4) is distinguished by the tall and erect loosely flowered inflorescences, carrying dull brownish to clear yellow flowers commonly with a purplish column and a bright yellow to orange, basally gibbous, carnoselip callus. *Cyrtochilum soennemarkii* differs from the similar but previously misapplied *Cyrtochilum viminale* Rehb.f.) Dalström (Fig. 5) by the gibbous and convex lip with a carnoselip callus, versus a basally straight, slightly elongated, smooth, dorsally furrowed and bifurcated callus of the same ivory white color as the lip lamina for *C. viminale*. *Cyrtochilum soennemarkii* is distinguished from all other yellow-flowered and superficially similar species by the gibbous lip with a carnoselip callus.




FIGURE 1. *Cyrtorchilum soennemarkii*. A. Plant habit. B. Column and lip lateral view. C. Column lateral and ventral views. D. Anther cap dorsal view and pollinia frontal, and lateral view with stipe. E. Flower dissected. Drawn from holotype by Stig Dalström.



FIGURE 2. *Cyrtochilum soennemarkii*. Plant cultivated and photographed by G. Deburghgraeve.



FIGURE 3. *Cyrtochilum soennemarkii*, plant habit, Azogues, Ecuador. Photo by S. Dalström.

Terrestrial herb. *Pseudobulbs* caespitose, ovoid, unifoliate or bifoliate, ca. 5×3 cm, subtended basally by 5 to 6 distichous sheaths, the uppermost foliaceous. *Leaves* subpetiolate, conduplicate, linear elliptic to obovate, acute to obtuse, sometimes apiculate, $22\text{--}34 \times 1.5\text{--}1.7$ cm. *Inflorescences* axillary and arising from the uppermost sheaths, erect to slightly arching, to ca. 80 cm long almost straight panicle, with widely spaced, to ca. 5 cm long and to 5-flowered side-branches; *bracts* appressed, scale-like, $0.2\text{--}1.2$ cm long; *pedicel* with *ovary* $0.5\text{--}1.5$ cm long. *Flowers* campanulate to stellate; *dorsal sepal* dull olive brown on type, but generally yellow and sometimes with purple stains near the base, unguiculate, broadly elliptic to obovate, obtuse, apiculate, ca. 8×4 mm; *lateral sepals* similar in color, slightly oblique, narrowly unguiculate to slightly spatulate, ovate, acute, ca. $10\text{--}11 \times 3.0\text{--}3.5$ mm; *petals* similar in color, broadly unguiculate, obliquely ovate, acute, ca. 8.5×4.0 mm; *lip* similar in color, rigidly fused to the base of the column, then gibbous and recurved away from the column, apically concave, truncate, broadly ovate to indistinctly 3-lobed, lateral lobes rotund, front lobe bluntly obtuse, ca. 7.5×6 mm; *callus* bright yellow, of a fleshy



FIGURE 4. *Cyrtochilum soennemarkii*, flowers of the type specimen. Photo by S. Dalström.

and broad, gibbose, central, longitudinally grooved, fleshy keel, extending from the base up to *ca.* 1/2 of the length of the lamina, ending in a pair of rounded, somewhat diverging angles, with an intermediate apical knob; *column* variably purplish, stocky, dorsally straight, ventrally concave, with concave lateral flanks, ending in a rounded lobe on each side of the stigmatic surface, *ca.* 4 mm long; *anther cap* purplish to yellow, campanulate, rostrate, dorsally lobulate; *pollinarium* of two obovoid, cleft, or folded, pollinia with an ovate, *ca.* 1 mm long stipe on a pulvinate viscidium.

ADDITIONAL MATERIAL SEEN: Ecuador. Cañar, uplands of “Huairacaja”, 10–20 km NE Azogues, 3600 m, 2 Feb. 1945, *W. H. Camp E-1788* (NY). Cañar, near San Marcos, NE Azogues, “1000” m [most certainly 3000 m; author’s note], 13 Apr. 1945, *W. E. Camp E-2602* (AMES). Cañar, Azogues to Taday, 3300 m, 9 Mar. 1992, *S. Dalström 1630* (SEL). Cañar, Azogues—Taday, km 16, 3300 m, 2 Feb. 1988, *U. Molau et al. 2849* (QCA). Azuay, Cuenca—Loja, km 78, 3000 m, 20 Sep. 1980, *C. A. Luer et al. 5507* (SEL). Azuay,



FIGURE 5. *Cyrtochilum viminale*. Plant cultivated and photographed by G. Deburghgraeve.

- Cuenca—Loja, km 52, 3250 m, 13 Dec. 1982, *S. Dalström* 343 (SEL). Azuay, Cuenca—Loja, km 42, 2900 m, 30 Nov. 1984, *C. H. Dodson & B. Malo* 15490 (MO, QCNE, RPSC). Azuay, Cuenca—Loja, km 67, 3380 m, 30 Jul. 1959, *H. G. Barclay & P. Juajibioy* 8573 (AMES, MO). Azuay, Sigüig, 2800 m, 6 May 1981, *J. Kuhn* 44 (SEL); Cuenca—Loja, km 30, 3360 m, 29 Dec. 80, *M. Madison et al.* 7407 (SEL). Azuay, Sigüig—Ludo, km 5, 3100 – 3300 m, 78° 50'W, 3° 05'S, 17 Nov. 1983, *B. Eriksen & B. B. Larsen* 45727 (AAU). Azuay, W Andes of Cuenca, 2400 – 2800 m, *F. C. Lehmann* 8069 (AMES). Azuay, Cuenca—Loja, km 80, 3000 m, 17 Jul. 1977, *C. A. Luer et al.* 1723 (SEL). Azuay, Cuenca—Loja, Paramo de Tinajillos, Sta Rosa, 3000 m, 2 Nov. 1988, *G. Harling & L. Andersson* 25580 (AMES). Azuay, Cuenca—Loja, 65 km S of Cumbe, 2900, 3 Nov. 1988, *G. Harling & L. Andersson* 25622 (AMES). Azuay, Cuenca—Loja, Paramo de Tinajillos, Sta Rosa, 3200, 4 Nov. 1988, *G. Harling & L. Andersson* 25636 (AMES). Azuay, Oña—Rio Yacuambi, 2600 m, 10 Sep. 1945, *F. Prieto P-206* (AMES). Azuay, Sigüig—Gualaquiza, Río Altarurucu, 2800 m, 13 Apr. 1968, *G. Harling et al.* 8285 (AMES, GB). Azuay, Sigüig, 10 km S, 3200 m, 3 Aug. 1975, *C. A. Luer et al.* 406, 421 & 416 (SEL). Azuay, Km 40 S Cuenca, ca. 3600 m, 20 Sept. 1944, *I. L. Wiggins* 10767, 10823 (AMES). Azuay, Cuenca—Loja, km 65 – 70, 3400 m, 3 Jan. 1981, *H. Balslev* 1423 (AMES). Loja, Loja—Cuenca, km 65, at pass above Saraguro, 3100 m, 20 Sep. 1980, *C. H. Dodson et al.* 10500 (SEL, MO). Loja, Loja—Cuenca, 12 km N Saraguro, 3000 m, 3 Jan. 1981, *H. Balslev* 1405 (NY). Loja, Cuenca—Loja, near Saraguro 3000 m, 5 Mar. 1977, *C. A. Luer et al.* 1524 (SEL). Loja, Loja—Cuenca, vicinity of Saraguro, 27 Sep. 1918, *J. N. Rose et al.* 23141 (US). Loja, Saraguro—Tenta, 2700 m, 16 Sep. 1980, *J. Jaramillo* 3828 (AAU, QCA, QCNE). Loja, San Lucas to Oña, 2600, 7 Sep. 1923, *Hitchcock* 21574 (US). Loja, Cuenca—Loja, km 80, 3000 m, 19 Dec. 1957, *C. H. Dodson & G. P. Frymire* 257 (SEL). Loja, Saraguro to Loja, 3100 m, 6 Feb. 1993, *S. Dalström* 1854 (SEL). Loja, Cuenca—Loja, km 140, 2800 m, 10 Feb. 1978, *C. A. Luer et al.* 2504 (SEL). Loja, Km 78 S Cuenca, ca. 3000 m, *C. A. Luer et al.* 5510 (SEL). Morona-Santiago, Sigüig—Chiguinda, 3000 m, 11 Aug. 1990, *A. Hirtz et al.* 5047 (QCNE). Morona Santiago, Gualaceo—Chiguinda, near Chiguinda, 3200 m, Jan. 1989, *A. Hirtz et al.* 4015 (MO).
- DISTRIBUTION AND HABITAT: *Cyrtorchilum soennemarkii* is reported from the high altitude, grassy and shrubby plains of the Ecuadorean provinces of Azuay, Cañar and Loja, at the altitude of 2400 – 3600 m.
- EPONYMY: Named in honor of Jan Sönnemark of Halmstad, Sweden, who contributed substantially to the understanding of the true identity of this species.
- ACKNOWLEDGMENTS. I thank Jan Sönnemark for many years of great companionship in the field.
- LITERATURE CITED
- Dalström, S. (2001). A synopsis of the genus *Cyrtorchilum* (Orchidaceae; Oncidiinae): Taxonomic reevaluation and new combinations. *Lindleyana*, 16(2), 56–80.
- Dalström, S. (2010). *Cyrtorchilum*. In Harling, G. & Persson, C. (Eds.), *Flora of Ecuador No. 87* (pp. 18–198). Sweden: Göteborg University.