



Semina: Ciências Agrárias

ISSN: 1676-546X

semina.agrarias@uel.br

Universidade Estadual de Londrina
Brasil

Silva Bonfim, Dihego; Costa de Siqueira, Jefferson; Delmondes Bomfim, Marcos Antonio;
Barbosa Ribeiro, Felipe; Loiola de Oliveira, Francisco; Cardoso Nagib Nascimento,
Daphinne; de Araujo Melo, Solange

Productive characteristics of meat quails reared in different environments

Semina: Ciências Agrárias, vol. 37, núm. 6, noviembre-diciembre, 2016, pp. 4313-4326

Universidade Estadual de Londrina
Londrina, Brasil

Available in: <http://www.redalyc.org/articulo.oa?id=445748913042>

- How to cite
- Complete issue
- More information about this article
- Journal's homepage in redalyc.org

redalyc.org

Scientific Information System

Network of Scientific Journals from Latin America, the Caribbean, Spain and Portugal

Non-profit academic project, developed under the open access initiative

Productive characteristics of meat quails reared in different environments

Características produtivas de codornas de corte mantidas em diferentes ambientes

Dihego Silva Bonfim^{1*}; Jefferson Costa de Siqueira²;
Marcos Antonio Delmondes Bomfim²; Felipe Barbosa Ribeiro²;
Francisco Loiola de Oliveira³; Daphinne Cardoso Nagib Nascimento⁴;
Solange de Araujo Melo⁵

Abstract

The objective of this study was to evaluate the effects of ambient temperature and consumption on the performance, carcass yields and cuts, organs, and body temperatures of meat quails. The experiment was conducted at the Federal University of Maranhão, campus of Chapadinha-MA, during September to October 2013. From the 14th day of age, 450 meat quails, not sexed, with an average initial weight of 90.4 ± 12.9 g, were distributed in batteries and accommodated in either acclimatized or non-acclimatized rooms, until 42 days of age. The experimental design was completely randomized, with three treatments, and six replicates of 25 birds each. The treatments were CE (acclimatized environment at 26 °C); NE (natural environment), and CECC (acclimatized environment at 26 °C with controlled feed, to keep the same consumption level as that of the birds in NE). The evaluated variables at 28 and 42 days of age were as follows: (1) the intake of feed, weight gain, feed conversion, body weight, and energy efficiency to weight gain, (2) the carcass weight, carcass yields (%) for breast, thigh, drumstick, and wing, and (3) the relative weights (%) of the digestive tract, heart, liver, gizzard, and intestine. In addition, we evaluated the cloacal and surface temperatures (°C), and subsequently, calculated the average and surface body temperatures. The treatment means were compared using the Student-Newman-Keuls test, with 5% level of significance. Compared with the acclimatized ambient environment, the natural environment more favored the performance of quails up to 28 days, but then affected it negatively from that age onward. Carcass yield, organs, and internal temperature of the quails maintained in the acclimatized and natural environments were not influenced.

Key words: *Coturnix coturnix coturnix*. Performance. Relative weights. Temperatures. Yields.

Resumo

Objetivou-se avaliar os efeitos da temperatura ambiente sobre o desempenho, rendimentos de carcaça e cortes, órgãos e temperaturas corporais de codornas de corte. O experimento foi conduzido na

¹ Mestre em Ciência Animal, Centro de Ciências Agrárias e Ambientais, Universidade Federal do Maranhão, CCAA/UFMA, Chapadinha, MA, Brasil. E-mail: dihegozootecnia@hotmail.com

² Profs., Centro de Ciências Agrárias e Ambientais, Universidade Federal do Maranhão, CCAA/UFMA, Chapadinha, MA, Brasil. E-mail: jc.siqueira@ufma.br; mad.bomfim@ufma.br; felipe.barbosa@ufma.br

³ Graduado em Zootecnia, UFMA, Chapadinha, MA, Brasil. E-mail: loiola93@gmail.com

⁴ Doutorado em Ciência Animal, Universidade Federal do Piauí, UFPI, Bom Jesus, PI, Brasil. E-mail: daphinnec@yahoo.com.br

⁵ Profª., UEMA, Balsas, MA, Brasil. E-mail: sol-vet@hotmail.com

* Author for correspondence

Universidade Federal do Maranhão, campus de Chapadinha-MA, no período de setembro a outubro de 2014. A partir do 14º dia de idade, 450 codornas de corte, não sexadas, com o peso inicial médio de $90,4 \pm 12,9$ g, foram distribuídas em baterias, acondicionadas em sala climatizada e não climatizada, até o 42º dia. O delineamento experimental foi o inteiramente casualizado, com três tratamentos e seis repetições de 25 aves. Os tratamentos consistiram em: AC (Ambiente climatizado com temperatura de 26°C); AN (Ambiente natural) e ACCC (Ambiente climatizado com temperatura de 26°C com alimentação controlada para manter o mesmo consumo das aves mantidas no AN). Foi avaliado o consumo de ração, ganho de peso, conversão alimentar, peso vivo, eficiência energética para ganho de peso; o peso de carcaça, rendimentos (%) de carcaça, peito, coxa, sobrecoxa e asa; os pesos relativos (%) do trato digestório, coração, fígado, moela e intestino aos 28 e aos 42 dias. Também foram avaliadas as temperaturas (°C) cloacais e superficiais, e posteriormente, calculadas a superficial média e corporal média. As médias dos tratamentos foram comparadas pelo teste SNK considerando o nível de 5% de significância. O ambiente natural melhorou o desempenho das codornas até os 28 dias, prejudicando a partir dessa idade em relação ao ambiente climatizado. Os rendimentos de carcaça, órgãos e a temperatura interna das codornas mantidas nos ambientes climatizado e natural, não foram influenciados.

Palavras-chave: *Coturnix coturnix coturnix*. Desempenho. Pesos relativos. Rendimentos. Temperaturas.

Introduction

High ambient temperature is one of the factors that have negative effects on feed intake, and consequently, on the productive characteristics of birds, resulting in economic losses to poultry farmers. The reduction of feed intake with increase in the ambient temperature is well-documented for broilers (EL-DEEP et al., 2014), laying hens (OLIVEIRA et al., 2014), turkeys (VELDKAMP et al., 2005), and laying quails (OZBEY; OZCELIK, 2004).

Studies of broilers maintained at high temperatures (32 °C) have allowed us to conclude that approximately 60% of the decrease in performance is indirectly due to a reduction in consumption, the other 40% being the direct result of the effects of temperature (BONNET et al., 1997; GERAERT et al., 1996). In this context, part of the decrease in the birds' performance might be caused by the deficit of energy and nutrients, and the remaining is associated with the increase in energy spent for heat dissipation, which is necessary for the maintenance of homeothermy.

The breeding of meat quails (*Coturnix coturnix coturnix*) has great potential for expansion in Brazil, as it requires only a small area for development, low manpower, and low investment but ensures quick

return of capital. In addition, there is a growing demand for this type of meat in the consumer market, mainly due to the meat's characteristics such as intense color, smoothness, and flavor (PASTORE et al., 2012). Moreover, Sousa et al. (2014) mention that compared with chickens, meat quails have higher thermal comfort temperatures and might be more tolerant to high ambient temperatures because of a greater surface/volume ratio, suggesting greater ability to dissipate the heat generated by their metabolism.

Therefore, the breeding of meat quails can be a very important socio-economic activity, representing an alternative for many farmers in the Northeast region of Brazil, where temperatures remain high during most parts of the year.

Considering that only a few studies have evaluated the effects of ambient temperature on feed intake and productive characteristics of meat quails, the objective of this study was to evaluate the productive characteristics of meat quails reared at different environments, from 14 to 42 days of age.

Materials and Methods

The experiment was carried out from 25th September to 23rd October 2014, at the Agricultural and Environmental Sciences Center of the Federal

University of Maranhão, located in the municipality of Chapadinha (03° 44' 30" S and 43° 21' 33" W), at an altitude of 105 m, as per the software SPRING 4.3.3® (INPE, 2010). According to Köppen's climate classification, the region's climate corresponds to the Aw type, considered as a tropical zone with dry winter (ALVARES et al., 2013).

During the pre-experimental period (1-13 days of age), the quails were kept in brick sheds and housed in metal cages with a wood-shavings floor. The cages were fitted with 60 W incandescent bulbs, for maintaining the ambient temperature between 32 °C and 35 °C.

On the 14th day of age, 450 meat quails (*Coturnix coturnix coturnix*), not sexed, with an average starting weight of 90.4 ± 12.9 g, were randomly distributed in batteries, containing cages of 0.72 m² (0.85 m × 0.85 m), packed in two brickwork rooms with an area of 38 m² (5.0 m × 7.7 m) and side windows. One of the rooms had an air conditioner with a capacity of 24,000 BTU/HR (British Thermal Unit = 252.2 cal) to climatize the room, whereas the other one had no air conditioning, to mimic the conditions of the natural environment.

A completely randomized experimental design was used, with three treatments and six replicates of 25 birds each, totaling 18 experimental units (cages). The treatments were CE (acclimatized environment with temperature of 26 °C); NE (natural environment), and CECC (acclimatized environment with temperature of 26 °C, but with controlled feeding to maintain the same consumption level as the birds kept in NE). CECC was used to isolate the effect of different environments on the feed intake of the birds.

The average, maximum, minimum, and black globe temperatures, as well as the relative humidity inside the premises, were monitored and recorded on a daily basis throughout the experimental period in the different environments (7:00 AM, 1:00 PM, and 7:00 PM), by using thermohygrometers located in the geometric centers of the rooms. To characterize

the different environments, the Index of Globe Temperature and Humidity (IGTU) was calculated, as proposed by Buffington et al. (1981): $IGTU = T_{gn} + 0.36 T_{po} - 330.08$, where T_{gn} = Black Globe Temperature (K) and T_{po} = Dew Point Temperature (K).

Continuous light was used throughout the experimental period (24 h of artificial light), with the help of 40 W fluorescent lamps.

The birds were fed rations based on corn and soybean meal, formulated to meet the nutritional requirements of meat quail during the breeding (1-21 days) and raising (22-42 days) phases, as recommended in the tables for Japanese and European Quails (SILVA; COSTA, 2009) (Table 1).

The quails of all experimental groups had free access to water and feed, except those of the CECC group, which received controlled feed so that they would consume the same amount of feed as that by the quails reared in NE (quantity from the previous day). For this, the feed consumption of the birds reared in NE was recorded daily and the same amount was supplied to the birds of the CECC group, as proposed by Geraert et al. (1996).

Feed intake (FI; g), live weight on the 28th and 42th days (W28 and W42; g), weight gain (WG; g), feed conversion (FC; g/g), and energy efficiency (EEWG; g/Mcal) were quantified for quails of ages 14-28 days and for the cumulative age period of 14-42 days. FI was obtained by the difference between the feed supplied and the remaining quantity in each period, divided by the number of birds, and was corrected according to mortality, considering the date of death as per Sakomura and Rostagno (2007). LW and WG for each period were obtained by weighing the birds in each group at 28 and 42 days of age. FC was calculated as the ratio of FI and WG. EEWG was calculated as the ratio of WG (g) and the consumption of metabolizable energy (Mcal), the latter being calculated as the product of FI (kg) and metabolizable energy of the feed (Mcal/kg).

Table 1. Rations formulated to meet the nutritional requirements of quails in the breeding and raising stages¹.

Ingredients (%)	Breeding (1-21 days)	Raising (22-42 days)
Corn	52.005	60.577
Soybean meal	43.466	36.019
Soya oil	0.000	0.920
Dicalcium phosphate	1.185	0.952
Limestone	1.009	0.826
Sodium chloride	0.376	0.325
DL-methionine (98%)	0.344	0.161
L-threonine (98%)	0.193	0.010
L-Lysine HCl (78.5%)	0.025	0.000
Vitamin compound ²	0.100	0.100
Mineral compound ³	0.050	0.050
Choline chloride (60%)	0.060	0.060
Inert (washed sand)	1.187	0.000
Total	100.00	100.00
Calculated Nutritional Compounds		
Metabolizable energy (kcal/kg)	2.950	3.100
Crude protein (%)	25.00	22.00
Calcium (%)	0.850	0.700
Available phosphorus (%)	0.320	0.270
Sodium (%)	0.170	0.150
Chlorine (%)	0.269	0.239
Potassium (%)	1.091	0.959
Methionine + Digestible cystine (%)	1.040	0.800
Digestible Methionine (%)	0.686	0.477
Digestible Lysine (%)	1.370	1.178
Digestible Threonine (%)	1.040	0.780
Digestible Valine (%)	1.143	1.017
Digestible Isoleucine (%)	1.048	0.914
Digestible Tryptophan (%)	0.243	0.210
Raw fiber (%)	3.255	3.061
Neutral detergent fiber (%)	9.222	9.320
Acid detergent fiber (%)	3.803	3.641

¹ Rations formulated based on composition data of the food and nutritional requirements of heavy quails presented in the Tables for Japanese and European Quails (SILVA; COSTA, 2009). ²Composition/kg of product: vit. A = 12,000,000 U.I.; vit. D3 = 3,600,000 U.I.; vit. E = 3,500 U.I.; vit B1 = 2,500 mg; vit. B2 = 8,000 mg; vit. B6 = 5,000 mg; Pantothenic acid = 12,000 mg; Biotin = 200 mg; vit. K = 3,000 mg; Folic acid = 1,500 mg; Nicotinic acid = 40,000 mg; vit. B12 = 20,000 mg; Se = 150 mg; vehicle q.s.

³Composition/kg of product: Mn = 160 g; Fe = 100 g; Zn = 100 g; Cu = 20 g; Co = 2 g; I = 2 g; vehicle q.s.

On the 28th and 42nd days, two quails weighing close to the average weight of each experimental unit ($\pm 5\%$) were sampled, totaling 36 birds for each age, which were identified and kept under fasting for 6 h, to reduce the contents of the digestive tract.

After fasting, the quails were killed by cervical dislocation, plucked, and eviscerated to obtain the weight of the carcass and offal, following the procedures approved by the Committee of Ethics in Animal Experimentation of the Federal University

of Maranhão (Case No. 23115.002714/2014-74). Next, the carcass was cut (chest, thigh, drumstick, and wing) and the internal organs were separated (heart, liver, gizzard, and intestine [small intestine + ceca + colon]); each part and organ were then individually weighed.

The carcass yield (%) (CY) was determined as the ratio of the weight of the eviscerated carcass (CW) and the weight after fasting. Breast (BY), thigh (TY), drumstick (DSY), and wing (WY) yields, relative weights of the digestive tract (RWDT) (pharynx + esophagus + crop + proventriculus + gizzard + pancreas + liver + intestine) and organs (heart [RWHRT], liver [RWLIV], gizzard [RWGIZ], and intestine [RWINT]) were determined with respect to the CW.

To check the effect of the environment on the body temperatures of quails at 28 and 42 days of age, two birds were randomly sampled from each group, and cloacal (CT) and surface (head, HDT; breast, BT; back BCKT, wing, WT; and legs, LGT) temperatures were measured at the time of day (1 PM) that was considered the most critical, by using clinical digital (Omron MC 245) and infrared (Instrutemp ITTI 380) thermometers, respectively. Based on these data, the average surface temperature was calculated ($AST = 0.12 \cdot WT + 0.03 \cdot HDT + 0.15 \cdot LGT + 0.70 \cdot BCKT$), followed by the average body temperature ($ABT = 0.3 \cdot AST + 0.7 \cdot TC$) (RICHARDS, 1971).

Initially, data for all the assessed variables were subjected to a normality test (Cramer-Von Mises) and a homoscedasticity test (Levene); the values were in agreement to all assumptions, except in the case of FI. The data for this variable did not comply with the assumption of homoscedasticity, as the CECC group received controlled feed with respect to the NE group, thus, showing no significant variation across replicates. As a result, the FI variance analysis was carried out considering the CE and NE groups.

The data for each variable were subjected to variance analysis according to the following statistical model: $Y_{ij} = \mu + A_i + e_{ij}$; with $i = 1, 2, 3$ and $j = 1, 2, 3, 4, 5, 6, \dots$,

where Y_{ij} = FI, LW, WG, FC, EEWG, CW, CY, BY, TY, DSY, WY, RWDT, RWHRT, RWLIV, RWGIZ, RWINT, CLT, HDT, BT, BCKT, WT, LGT, AST, or ABT of the birds kept in the i^{th} experimental group; μ = effect of the general average; A_i = effect of the i^{th} experimental group; and e_{ij} = experimental error associated to j^{th} replicate.

Subsequently, the averages of experimental groups for each variable for the two periods (14-28 and 14-42 days of age) were compared using the Student-Newman-Keuls test (SNK), considering a significance level of up to 5% probability. Statistical analyses were performed using the GLM procedure of the SAS Software 9.0 (2002).

Results and Discussion

During the trial period, the average, maximum, minimum, and black globe temperatures ($^{\circ}\text{C}$) recorded within the acclimatized environment were 26.1 ± 0.62 ; 27.2 ± 0.64 ; 24.8 ± 0.80 , and 27.4 ± 0.64 $^{\circ}\text{C}$, respectively, with an average relative humidity of $69.5 \pm 9.25\%$. In the non-acclimatized environment, these temperatures were 32.2 ± 0.32 ; 34.5 ± 0.61 ; 29.9 ± 0.54 , and 33.7 ± 0.33 $^{\circ}\text{C}$, respectively, with a relative humidity of $56.8 \pm 2.26\%$. The average temperature and relative humidity were 25.8 ± 1.20 $^{\circ}\text{C}$ and $67.5 \pm 3.89\%$ at 7 AM, 25.9 ± 0.98 $^{\circ}\text{C}$ and $67.3 \pm 5.23\%$ at 1 PM, and 25.8 ± 1.15 $^{\circ}\text{C}$ and $66.4 \pm 6.17\%$ at 7 PM, respectively, in the acclimatized environment, and 31.9 ± 1.93 $^{\circ}\text{C}$ and $52.6 \pm 3.01\%$ at 7 AM, 32.8 ± 1.98 $^{\circ}\text{C}$ and $67.3 \pm 5.40\%$ at 1 PM, and 32.2 ± 1.90 $^{\circ}\text{C}$ and $52.4 \pm 3.02\%$ at 7 PM, respectively, in the natural environment.

To characterize the thermal environment by using a single value that represents the impact of the variables affecting the thermal balance of the animal,

Buffington et al. (1981) proposed the Black Globe and Humidity Index (BGHI), which has been used widely, as it incorporates, among other variables, the dry bulb temperature, black globe temperature, and relative humidity. Thus, the BGHI values throughout the experimental period were calculated, and the average values were found to be 75.8 ± 1.36 and 83.0 ± 0.46 in environments with and without air conditioning, respectively. The BGHI was 75.4 ± 1.08 at 7 AM, 75.3 ± 1.45 at 1 PM, and 75.2 ± 1.66 at 7 PM in the climatized environment, and 82.3 ± 1.30 at 7 AM, 83.4 ± 0.91 at 1 PM, and 82.6 ± 1.33 at 7 PM in the natural environment.

Recently, Sousa et al. (2014) studied the effects of different thermal environments on the meat quails' performance from ages 22-35 days, in order to determine the thermal comfort ranges,

expressed in terms of temperature ($^{\circ}\text{C}$) and BGHI. The authors concluded that temperatures between 25.6 and 26.7 $^{\circ}\text{C}$, corresponding to BGHI values between 75.3 and 75.8 , characterize thermal comfort, whereas temperatures between 30.4 and 33.2 $^{\circ}\text{C}$, corresponding to BGHI values between 79.7 and 82.2 , are characteristic of a moderate/severe heat condition. Based on the results of Sousa et al. (2014), the average temperatures of 26.1 and 32.2 $^{\circ}\text{C}$ observed in this study, corresponding to BGHI of 75.8 and 83.0 , would characterize the environment as comfortable and moderately/severely hot, respectively.

FI (g), WG (g), FC (g/g), W28d (g), W42d (g), and EEWG (g/Mcal) of the quails were influenced by the environment ($P < 0.05$), independent of the evaluation period (14-28 and 14-42 days) (Table 2).

Table 2. Feed intake (FI), weight gain (WG), feed conversion (FC), body weight (W28d, W42d), and energy efficiency for weight gain (EEWG) of meat quails kept in different environments for different evaluation periods (14-28 and 14-42 days).

Variable	Treatments			VC ¹	P>F ²
	CE (26 °C)	NE (32 °C)	CECC (26 °C)		
14-28 days					
FI ³ (g)	346.37 ^a	330.50 ^b	331.40	2.31	0.0056
WG (g)	126.71 ^a	120.99 ^b	106.96 ^c	2.37	<0.0001
FC (g/g)	2.73 ^b	2.73 ^b	3.09 ^a	2.42	<0.0001
W28d (g)	215.33 ^a	209.78 ^b	195.25 ^c	1.46	<0.0001
EEWG (g/Mcal)	124.0 ^a	124.2 ^a	109.4 ^b	2.41	<0.0001
14-42 days					
FI ³ (g)	802.06 ^a	761.10 ^b	761.56	2.13	0.0017
WG (g)	201.39 ^a	184.89 ^b	181.70 ^b	2.99	<0.0001
FC (g/g)	3.99 ^b	4.12 ^{ab}	4.19 ^a	2.82	0.0220
W42d (g)	290.02 ^a	273.68 ^b	270.18 ^b	2.00	<0.0001
EEWG (g/Mcal)	81.05 ^a	78.37 ^{ab}	76.97 ^b	2.94	0.0243

Averages followed by the same letters in the same line do not differ by the SNK test ($P > 0.05$).

¹Variation Coefficient (%).

²Significance of the "F" test of the variance analysis.

³For the variable FI, the variance analysis was performed with the CE and NE treatments.

In the initial period (14-28 days), it was observed that the NE birds had lower FI ($P < 0.05$) by 4.58% when than those in CE, resulting

in a decrease ($P < 0.05$) of 4.51% in the WG and 2.58% in W28d. As the reduction in WG was proportional to the reduction in FI, the FC and

EEWG were similar ($P > 0.05$) between these groups.

Considering the 26 °C ambient temperature, it was observed that quails in the CECC group showed a 4.32% lower FI than those of the CE group, resulting in a deterioration ($P < 0.05$) of 15.59% in WG, 13.18% in FC, 9.32% in W28d, and 11.77% in EEWG. As the deterioration observed in the performance variables of the CECC birds was greater than the reduction of the FI, it is possible that the ambient temperature of 26 °C had a negative effect on the performance variables evaluated during 14-28 days of age.

Assuming that temperatures between 25.6 and 26.7 °C are characteristic of thermal comfort and temperatures between 30.4 and 33.2 °C represent moderate/severe heat (SOUSA et al., 2014), it was expected that quails of the CECC group would exhibit higher performance than those of the NE group, as has been noted in broilers (BONNET et al., 1997; GERAERT et al., 1996); however, the situation was precisely the opposite.

Quails from the CECC group consumed the same amount of feed as those in the NE group, however, there was a deterioration of ($P < 0.05$) 11.60% in the WG, 13.18% in FC, 6.92% in W28d, and 11.91% in EEWG, which confirms the fact that the temperature of 26 °C detracted the performance of quails till 28 days of age.

The lower performance of the CECC group compared with the NE group is probably associated with the increased metabolic activity, in order to increase heat production for maintaining thermal homeostasis, causing an increase in the net energy requirements for maintenance, and consequently, reducing the net energy available for gain (JORDÃO FILHO et al., 2011); this is supported by the observed worsening of EEWG. These results suggest that the range of thermal comfort for meat quails for 14-28 days of age is above 26 °C, which differs from the recommendations of Sousa et al. (2014) (25.6-26.7 °C).

During the later accumulated period (14-42 days), there was a decrease ($P < 0.05$) of 5.11% in the FI of quails kept in NE, when compared with those of the CE group, which resulted in a reduction ($P < 0.05$) of 8.19% in WG and 5.63% in W42d, with no effect ($P > 0.05$) on FC and EEWG.

At the ambient temperature of 26 °C, it was observed that quails in the CECC group showed a 5.04% lower FI ($P < 0.05$) than those of the CE group, resulting in a deterioration of 9.77% in WG, 5.01% in FC, 6.84% in W42d, and 5.03% in EEWG.

The reduction in WG due to FI was lower in the accumulated period (9.77%) (14-42 days) than in the initial period (15.59%) (14-28 days), which suggests that after 28 days, the ambient temperature of 26 °C improved the performance of quails with respect to the 32 °C environment. The comparison between the CECC and NE groups reinforces this notion, as the quails from the CECC group showed similar FI, WG, FC, W42d, and EEWG ($P > 0.05$) to those in the NE group, during the accumulated period.

The evaluations in this study were conducted on the same birds, sequentially at 28 and 42 days of age, and therefore, comparisons involving the final period (28-42 days) are questionable from a statistical point of view, because of the residual effect of the initial phase (14-28 days). On the other hand, the results can help clarify the fact that quails of the CECC group showed a higher WG than did those of the NE group after 28 days, as 28-day-old quails of the CECC group showed a lower W28d (195.25 g) ($P < 0.05$) than did those of the NE group (209.78 g); however, after 42 days, the W42d was similar ($P > 0.05$) between these groups (270.18 and 273.68 g, respectively). These results confirm that the 26 °C environment was more suitable than the 32 °C environment for quails from 28 to 42 days of age.

Studies on broilers kept at thermoneutral (22 °C) and stressful (32 °C) environment showed that approximately 60% of the reduction in performance

is indirectly due to the reduction of consumption, while the remaining 40% is related to the direct effects of high temperature (BONNET et al., 1997; GERAERT et al., 1996). In this study, the division of the direct and indirect effects of ambient temperature for meat quails became impossible because the temperature of 26 °C, recommended as thermally comfortable (SOUSA et al., 2014),

lowered the birds' performance in the CECC group compared with the NE group, until 28 days of age, and favored the performance from that age onward.

The quails' CW (g) was affected ($P < 0.05$) by the environment at 28 and 42 days of age, however, CY (%), PR (%), TY (%), DSY (%), and WY (%) were similar ($P > 0.05$) among the experimental groups at both age groups (Table 3).

Table 3. Carcass weight (CW), and carcass (CY), breast (BY), thigh (TY), drumstick (DSY), and wing (WY) yields of meat quails in different environments/feed intake at 28 and 42 days of age.

Variable	Treatments			VC ¹	P>F ²
	CE (26 °C)	NE (32 °C)	CECC (26 °C)		
28 Days					
BW (g)	172.96 ^a	164.11 ^b	159.50 ^b	3.57	0.0042
CY (%)	80.33 ^a	78.21 ^a	80.95 ^a	3.62	0.2612
BY (%)	33.31 ^a	32.75 ^a	33.95 ^a	2.49	0.0891
TY (%)	8.75 ^a	8.95 ^a	8.67 ^a	3.19	0.2592
DSY (%)	12.11 ^a	12.12 ^a	12.09 ^a	3.56	0.9942
WY (%)	8.16 ^a	8.48 ^a	8.37 ^a	4.41	0.3406
42 days					
BW (g)	227.25 ^a	217.31 ^b	216.51 ^b	2.92	0.0225
CY (%)	78.38 ^a	79.45 ^a	80.13 ^a	3.16	0.4936
BY (%)	32.41 ^a	33.56 ^a	32.29 ^a	5.32	0.4434
TY (%)	8.02 ^a	7.95 ^a	8.32 ^a	5.16	0.2882
DSY (%)	12.62 ^a	12.28 ^a	12.41 ^a	5.05	0.6495
WY (%)	7.23 ^a	7.40 ^a	7.32 ^a	4.44	0.7105

Averages followed by the same letters in the same line do not differ by the SNK test ($P > 0.05$).

¹Variation Coefficient (%).

²Significance of the "F" test of the variance analysis.

At 28 days of age, a decrease of 5.12% ($P < 0.05$) in the CW of NE quails was observed, along with a decrease of 7.78% in those from the CECC group, when compared with the CE quails. Although the CW of quails from the CE group (172.96 g) was higher ($P < 0.05$), compared with the NE (164.11 g) and CECC (159.50 g) groups, it was observed that the CY was similar ($P > 0.05$) among the groups.

Similarly, at 42 days, the quails from the NE and CECC groups showed a decrease of 4.37% and 4.73%, respectively, in CW ($P < 0.05$), when

compared with those of the CE group, without any effect ($P > 0.05$) on the CY.

Studies with broilers demonstrated that high temperatures (32 °C) induce a reduction in the size of the most metabolically active organs, in order to reduce the production of body heat, which results in increased carcass ratio:weight after fasting, and consequently, the increase of the carcass yield (OBA et al., 2012; ROSA et al., 2007); however, these changes were not evident in meat quails at 28 or 42 days of age.

Evaluating the effect of environment on carcass characteristics of broilers at 42 days of age, Rosa et al. (2007) observed that an environment at 32 °C resulted in an increase in carcass (2.58%) and thigh + drumstick (2.61%) yield, along with lower breast yield (11.19%), when compared with the group that was kept at 23 °C. Oba et al. (2012) also observed an increase in the thigh + drumstick yield (34.65%) and reduced breast yield (31.98%) in 47-day old broilers kept at 32 °C, compared with those kept at 22 °C.

These results might be related to the fact that the broilers' breast muscle show a glycolytic metabolism (white fibers), whereas the muscles in the legs are characterized by oxidative metabolism (red fibers). Thus, due to broilers' panting during heat stress, higher breast muscle activity occurs, consuming part of the glycogen reserves and damaging muscle development. On the other hand, the yield of thigh + drumstick results from the formation of large amount of red fibers, and might show a considerable reserve of fat, which is used as an energy source (OBA et al., 2012).

Choi et al. (2014) reported that the breast muscle of flying birds, including quails, has a higher proportion of oxidative fibers (Type IIA) compared with the glycolytic type (Type IIB), while in broilers, this muscle predominantly includes glycolytic fibers (Type IIB). The authors explained the lower proportion of white fibers (glycolytic) in the breast of quails to the greater frequency of flight exercise, requiring high energy expenditure by glycolytic pathways, resulting in a greater proportion of red fibers (oxidative), compared with broilers.

Yunianto et al. (1997) associated the changes in muscle protein "turnover" of broilers kept at high temperatures to changes in the functions of the endocrine system, especially the rise of plasma corticosterone concentration. According to the results of Silva et al. (2005) in studies on mammals, the catabolic effects of steroid hormones (corticosterone and cortisol) are more intense on the

glycolytic fibers than on the oxidative ones.

Considering that quails have a higher surface-to-body volume ratio when compared with broilers, and consequently have a higher dissipation capacity of metabolic heat by sensitive media, it is possible that quails are less susceptible to changes in the endocrine system, allowing the plasma levels of corticosterone without major unbalances, even when kept at temperatures that are considered high for broilers.

Moreover, considering that the catabolic effects of steroid hormones are more intense on glycolytic fibers than on oxidative ones (SILVA et al., 2005), and given that quails have a higher proportion of oxidative fibers than broilers (CHOI et al., 2014), the absence of temperature effects on carcass and meat yield observed in this study might be justified.

At 28 days, no effects ($P > 0.05$) of the environment were observed on the RWDT (%), RWLIV (%), RWGIZ (%), and RWINT (%) of quails, but there was an effect on their ($P < 0.05$) RWHRT (%). At 42 days, similar results were observed ($P > 0.05$) for RWDT, RWHRT, RWLIV, and RWINT, with the exception of RWGIZ, where the effect ($P < 0.05$) of the environment/feed intake was evident (Table 4).

There was no direct effect of the environment observed on the RWHRT in 28-day old quails, since the NE group did not differ ($P > 0.05$) from the CECC group, however, it was found ($P < 0.05$) that the value of this variable was lower in birds of the NE and CECC groups when compared with those of the CE group. This superiority of CE quails in terms of RWHRT can be explained by the indirect effect of feed intake (FI), since the birds in this group had higher FI (346.37 g) than those of the NE (330.50 g) and CECC (331.40 g) groups. Generally, the higher FI might be related to an increased cardiac activity, to meet the demand for oxygen (WIDEMAN; TACKETT, 2000), mainly in birds with a higher body weight (OLIVEIRA NETO et al., 2005), as noted in quails of the CE group, in this study.

Table 4. Relative weights of the digestive tract (RWDT), heart (RWHRT), liver (RWLIV), gizzard (RWGIZ), and intestine (RWINT) of quails in different environments/feed intake at 28 and 42 days of age.

Variable	Treatments			VC ¹	P>F ²
	CE (26 °C)	NE (32 °C)	CECC (26 °C)		
28 days					
RWDT (%)	9.83 ^a	9.40 ^a	9.49 ^a	5.53	0.4617
RWHRT (%)	1.09 ^a	0.97 ^b	0.95 ^b	4.66	0.0011
RWLIV (%)	3.10 ^a	3.14 ^a	2.92 ^a	7.31	0.2503
RWGIZ (%)	2.88 ^a	2.76 ^a	2.93 ^a	10.81	0.6402
RWINT (%)	2.92 ^a	2.80 ^a	2.99 ^a	10.82	0.6193
42 days					
RWDT (%)	8.94 ^a	8.00 ^a	9.14 ^a	11.27	0.1350
RWHRT (%)	1.02 ^a	0.96 ^a	1.05 ^a	8.70	0.2752
RWLIV (%)	3.13 ^a	2.53 ^a	2.90 ^a	14.00	0.0730
RWGIZ (%)	2.40 ^a	1.99 ^b	2.49 ^a	6.34	0.0002
RWINT (%)	3.16 ^a	2.97 ^a	2.80 ^a	14.49	0.3733

Averages followed by the same letters in the same line do not differ by the SNK test ($P > 0.05$).

¹Variation Coefficient (%).

²Significance of the “F” test of the variance analysis.

At 42 days of age, there was a decrease ($P < 0.05$) in the RWGIZ (%) of NE quails compared with those from the CE and CECC groups, with no additional evidence to justify such a result.

Based on the results observed in broilers, it was expected that the quails kept at 32 °C would present a lower relative weight of the lower organs than those kept at 26 °C. However, this was not the case, which suggested that for physiological adaptation, there was no need to reduce the relative weight of these organs; the results were different from those obtained with broilers (ROSA et al., 2007; OLIVEIRA NETO et al., 2000).

Rosa et al. (2007) observed a decrease in the relative weights of the heart and liver in broiler chickens (42 days of age) kept under heat stress (32 °C), compared with those kept at thermoneutral conditions (23 °C). These authors linked the reduction in maintenance requirements to the decrease in mass of the internal organs and the lower metabolic activity of these organs in the birds kept at 32 °C.

Oliveira Neto et al. (2000) also found a reduction of the relative weights of heart, liver, gizzard, and intestine in broilers (22-42 days of age) kept at 32 °C, compared with those kept at 23 °C. According to the authors, the high ambient temperatures can cause various adaptive physiological changes resulting in modification of the size of the organs.

The metabolic heat production rate is reduced when birds are exposed to high temperatures (GERAERT et al., 1996), increasing the plasma concentration of corticosterone and reducing the serum levels of triiodothyronine hormones (T3) and thyroxine (T4) (SAHIN et al., 2002), thus, changing the relative weight of the digestive tract and organs. Based on this, the physiological and hormonal changes in the quails (at 42 days of age) kept at 32 °C might not reach a high enough value to change the relative weights (RWDT (%), RWHRT (%), RWLIV (%), and RWINT (%)), confirming a greater resistance of these birds in hot environments, compared with broilers, as a consequence of the greater efficiency of dissipation of metabolic heat to the environment.

At 28 and 42 days of age, it was observed that the CLT (°C) of the quails was not influenced ($P > 0.05$)

by the environment. The WT (°C) at 28 days was also not influenced ($P > 0.05$) by the environment (Table 5).

Table 5. Temperatures of the (°C) cloaca (CLT), head (HT), breast (BT), back (BCKT), wing (WT), leg (LGT), and average surface (AST) and average body (ABT) temperature of quails (28- and 42-day old) in different environments.

Variable	Treatments			VC ¹	P>F ²
	CE (26 °C)	NE (32 °C)	CECC (26 °C)		
28 days					
CLT	40.46 ^a	40.74 ^a	40.35 ^a	1.03	0.2726
HT	33.87 ^c	37.18 ^a	35.19 ^b	2.05	0.0001
BT	35.08 ^b	37.87 ^a	37.87 ^a	3.28	0.0075
BCKT	33.85 ^b	37.79 ^a	35.19 ^b	3.13	<0.0001
WT	36.86	38.12	37.62	3.08	0.1986
LGT	33.33 ^b	37.45 ^a	33.16 ^b	2.27	<0.0001
AST ³	33.92 ^c	37.76 ^a	35.20 ^b	2.32	<0.0001
ABT ⁴	38.41 ^c	39.85 ^a	38.88 ^b	0.85	<0.0001
42 days					
CLT	40.20 ^a	41.01 ^a	40.10 ^a	2.16	0.1756
HT	35.37 ^b	37.68 ^a	34.60 ^b	4.00	0.0056
BT	36.31 ^{ab}	37.58 ^a	35.38 ^b	3.09	0.0141
BCKT	36.46 ^{ab}	37.79 ^a	35.85 ^b	3.43	0.0484
WT	36.43 ^b	38.46 ^a	35.60 ^b	3.50	0.0048
LGT	34.03 ^b	36.41 ^a	32.57 ^c	2.99	<0.0001
AST ³	36.06 ^b	37.66 ^a	35.29 ^b	2.78	0.0032
ABT ⁴	38.96 ^b	40.00 ^a	38.66 ^b	2.10	0.0306

Averages followed by the same letters in the same line do not differ by the SNK test ($P > 0.05$).

¹Variation Coefficient (%).

²Significance of the “F” test of the variance analysis.

³AST = (0.12*WT) + (0.03*HT) + (0.15*LGT) + (0.70*BCKT); ⁴ABT = (0.3*AST) + (0.7*CLT) (RICHARDS, 1971).

The fact that a difference ($P > 0.05$) among the CLT of quails (28 and 42 days of age) of the experimental groups (CE, NE, and CECC) was not seen might be an indication that even when kept at high temperature, the NE quails efficiently dissipated the body heat, thus, maintaining their homeothermy. Exposure to high ambient temperatures was the reason for quails of the NE group (28 and 42 days of age) to show higher average temperatures (°C) of the head (CA), breast (BT), back (BCKT), and surface (AST) ($P < 0.05$) than those of the groups kept at 26 °C.

These results are in agreement with those found by Dahlke et al. (2005), who observed an increase in the surface temperatures of the head, breast, back, and the average surface temperature of 42-day old broilers kept at 32 °C, compared with those kept at 22 °C.

Dahlke et al. (2005) argue that the temperature variations on the outer surface of broilers (head, chest, and back) are mechanisms available to the birds to keep their internal temperature constant, suggesting an increase in the heat flow from the body core towards the external surface of the

body. Based on these factors, the higher surface temperatures of quails kept at 32 °C than those kept at 26 °C is justified.

The cloacal temperature of the birds generally varies between 40-42 °C (BROWN-BRANDT et al., 2003), and when this limit is exceeded, there is an observed difficulty in dissipating body heat by the sensitive means. The results of this study showed that ambient temperatures of 26 °C or 32 °C allowed the maintenance of a constant body-core temperature, suggesting that the heat loss mechanisms (sensitive and latent) were sufficient to prevent its variation.

ABT of the NE quails was higher ($P < 0.05$) when compared with the quails kept at 26 °C, at 28 and 42 days of age. Considering the absence of effect ($P > 0.05$) of the ambient temperature on the CLT of quails, most ABT observed in the NE can be attributed to the increased AST, keeping in mind that it is responsible for 30% of the ABT (RICHARDS, 1971).

The sensitive exchange (conduction, convection, and radiation) of heat is of major importance for thermal balance; however, such a heat-flow mechanism depends directly on a temperature difference (temperature gradient) between the average surface temperature of the bird and the ambient temperature. According to Brown-Brandt et al. (1997), the higher is the temperature differential, more efficient would be the exchange of sensitive heat.

Considering this, the calculated thermal gradients (°C) were 7.82 (CE); 5.56 (NE), and 9.10 (CECC) for 28-day old quails, while for the 42-day old ones, the thermal gradients (°C) were 9.96 (CE); 5.46 (NE), and 9.19 (CECC). These results indicate that quails of the NE group dissipated the metabolic heat by sensitive media at a lower proportion than the birds kept at 26 °C (28 and 42 days), using evaporative cooling (latent exchange mechanism) as a resource. This mechanism causes greater energy expenditure, resulting in a worse performance (YAHAV et al., 2004), as noted in the quails of the NE group (Table

3) during the accumulated period (14-42 days).

The results of this study showed that, despite the fact that quails raised at 32 °C showed a lower FI, WG, W42, and CW, compared with those kept at 26 °C, the FC, EEWG, and carcasses and meat yields were not influenced, showing that this is a technically viable activity, especially in regions where temperatures are high during most of the year, such as the Brazilian Northeast. In this context, there is an alternative for producers in these regions, considering that small areas are needed for the development of this activity, in addition to low need for labor, low initial investment, and quick return of capital, as well as the favorable demand for this meat in the consumer market.

The breeding of meat quails from 14 to 42 days of age is technically feasible in acclimatized (26 °C) and natural (32 °C) environments. Compared with quails from the acclimatized ambient environment, those in the natural environment showed higher performance up to 28 days of age, but the performance decreased from that age onward. The carcass and organ yields and the internal temperature of quails were not influenced by the environment.

Acknowledgments

We thank Professor Dr. Celso Yoji Kawabata (*In Memoriam*) for his valuable cooperation in designing, conducting, and writing the essay. We also thank the Foundation for Support to Research and Scientific and Technological Development of the State of Maranhão (FAPEMA) for funding the research. We express our gratitude to the Higher Education Personnel Improvement Coordination (CAPES) for the scholarships granted.

References

ALVARES, C. A.; STAPE, J. L.; SENTELHAS, P. C.; MORAES, G. J. L. de; SPAROVEK, G. Köppen's climate classification map for Brazil. *Meteorologische Zeitschrift*, Stuttgart, v. 22, n. 6, p. 711-728, 2013.

- BONNET, S.; GERAERT, P. A.; LESSIRE, M.; CARRE, B.; GUILLAUMIN, S. Effect of high ambient temperature on feed digestibility in broilers. *Poultry Science*, Savoy, v. 76, n. 6, p. 857-863, 1997.
- BROWN-BRANDT, T. M.; BECK, M. M.; SCHULTE, D. D.; PARKHURST, A. M.; DeSHAZER, J. A. Physiological responses of tom turkeys to temperature and humidity change with age. *Journal of Thermal Biology*, Oxford, v. 22, n. 1, p. 43-52, 1997.
- BROWN-BRANDT, T. M.; TADAYUKI, Y. J.; HONGWEI, X.; GATES, R. S.; BUCKLIN, R. A. A new telemetry system for measuring core body temperature in livestock and poultry. *Applied Engineering in Agriculture*, St. Joseph, v. 19, n. 5, p. 583-589, 2003.
- BUFFINGTON, D. E.; AROCHO, A. C.; CANTON, G. H.; PITT, D.; THATCHER, W. W.; COLLIE, R. J. Black globe humidity index (BGHI) as a comfort equation for dairy cows. *Transaction of the ASAE*, St. Joseph, v. 24, n. 3, p. 711-714, 1981.
- CHOI, Y. M.; SUH, Y.; SHIN, S.; LEE, K. Skeletal muscle characterization of Japanese quail line selectively bred for lower body weight as an avian model of delayed muscle growth with hypoplasia. *PLoS One*, San Francisco, v. 9, n. 4, p. 1-14, 2014.
- DAHLKE, F.; GONZALES, E.; GADELHA, A. C.; MAIORKA, A.; BORGES, S. A.; ROSA, P. S.; FARIA FILHO, D. E.; FURLAN, R. L. Empenamento, níveis hormonais de triiodotironina e tiroxina e temperatura corporal de frangos de corte de diferentes genótipos criados em diferentes condições de temperatura. *Ciência Rural*, Santa Maria, v. 35, n. 3, p. 664-670, 2005.
- EI-DEEP, M. H.; MAHMOUD, H.; DAICHI, I.; EID, Y. Z.; HISAYA, Y.; AKIRA, O. Effects of dietary supplementation with *Aspergillus awamori* on growth performance and antioxidative status of broiler chickens exposed to high ambient temperature. *Journal of Poultry Science*, Savoy, v. 51, n. 3, p. 281-288, 2014.
- GERAERT, P. A.; PADILHA, J. C.; GUILLAUMIN, S. Metabolic and endocrine changes induced by chronic heat exposure in broiler chickens: Growth performance, body composition and energy retention. *British Journal of Nutrition*, Cambridge, v. 75, n. 2, p. 195-204, 1996.
- INSTITUTO NACIONAL DE PESQUISAS ESPACIAIS - INPE. Manual do software SPRING (Sistema de processamento de informações georreferenciadas). Rio de Janeiro: Divisão de Processamento de Imagens, 2010. v.4.3.3.
- JORDÃO FILHO, J.; SILVA, J. H. V.; SILVA, C. T.; COSTA, F. G. P.; SOUSA, J. M. B.; GIVISIEZ, P. E. N. Energy requirement for maintenance and gain for two genotypes of quails housed in different breeding rearing systems. *Revista Brasileira de Zootecnia*, Viçosa, MG, v. 40, n. 11, p. 2415-2422, 2011.
- OBA, A.; LOPES, P. C. F.; BOIAGO, M. M.; SILVA, A. M. S.; MONTASSIER, H. J.; SOUZA, P. A. Características produtivas e imunológicas de frangos de corte submetidos a dietas suplementadas com cromo, criados sob diferentes condições de ambiente. *Revista Brasileira de Zootecnia*, Viçosa, MG, v. 41, n. 5, p. 1186-1192, 2012.
- OLIVEIRA NETO, A. R.; OLIVEIRA, R. F. M.; DONZELE, J. L.; ROSTAGNO, H. S.; FERREIRA, R. A.; MAXIMIANO, H. C.; GASPARINO, E. Efeito da temperatura ambiente sobre o desempenho e características de carcaça de frangos de corte alimentados com dieta controlada e dois níveis de energia metabolizável. *Revista Brasileira de Zootecnia*, Viçosa, MG, v. 29, n. 1, p. 183-190, 2000.
- OLIVEIRA NETO, A. R.; OLIVEIRA, R. F. M.; DONZELE, J. L.; ROSTAGNO, H. S.; FERREIRA, R. A.; MAXIMIANO, H. C.; GASPARINO, E. Níveis de metionina + cistina para pintos de corte mantidos em ambiente termoneutro. *Revista Brasileira de Zootecnia*, Viçosa, MG, v. 34, n. 6, p. 1956-1962, 2005.
- OLIVEIRA, D. L.; NASCIMENTO, J. W. B.; CAMERINI, N. L.; SILVA, R. C.; FURTADO, D. A.; ARAUJO, T. G. P. Desempenho e qualidade de ovos de galinhas poedeiras criadas em gaiolas enriquecidas e ambiente controlado. *Revista Brasileira de Engenharia Agrícola e Ambiental*, Campina Grande, v. 18, n. 11, p. 1186-1191, 2014.
- OZBEY, O.; OZCELIK, M. The effect of high environmental temperature on growth performance of Japanese quails with different body weights. *Journal of Poultry Science*, Savoy, v. 3, n. 7, p. 468-470, 2004.
- PASTORE, S. M.; OLIVEIRA, W. P.; MUNIZ, J. C. L. Panorama da coturnicultura no Brasil. *Revista Eletrônica Nutritime*, Viçosa, v. 9, n. 6, p. 2041-2049, 2012.
- RICHARDS, S. A. The significance of changes in the temperature of the skin and body core of the chicken in the regulation of heat loss. *Journal of Physiology*, Cambridge, v. 216, n. 1, p. 1-10, 1971.
- ROSA, P. S.; FARIA FILHO, D. E.; DAHLKE, F.; VIEIRA, B. S.; MACARI, M.; FURLAN, R. L. Performance and carcass characteristics of broiler chickens with different growth potential and submitted to heat Stress. *Brazilian Journal of Poultry Science*, Savoy, v. 9, n. 3, p. 181-186, 2007.

- SAHIN, K.; OZBEY, O.; ONDERCI, M.; CIKIM, G.; AYSONDU, M. H. Chromium supplementation can alleviate negative effects of heat stress on egg production, egg quality and some serum metabolites of laying Japanese quail. *The Journal of Nutrition*, Bethesda, v. 132, n. 6, p. 1265-1268, 2002.
- SAKOMURA, N. K.; ROSTAGNO, H. S. *Métodos de pesquisa em nutrição de monogástricos*. Jaboticabal: Funep, 2007. 283 p.
- SILVA, E. C.; SHIMANO, A. C.; NEDER, L.; ROSSI, A. C.; MARTINEZ, J. A. B. Efeitos da administração sistêmica de corticosteroide sobre propriedades mecânicas musculares. *Revista Brasileira de Fisioterapia*, São Carlos, v. 9, n. 2, p. 203-209, 2005.
- SILVA, J. H. V.; COSTA, F. G. P. *Tabela para codornas japonesas e europeias*. 2. ed. Jaboticabal: FUNEP, 2009. 110 p.
- SOUSA, M. S.; TINÔCO, I. F. F., BARRETO, S. L. T.; AMARAL, A. G.; PIRES, L. C.; FERREIRA, A. S. Determinação de limites superiores da zona de conforto térmico para codornas de corte aclimatizadas no Brasil de 22 a 35 dias de idade. *Revista Brasileira de Saúde e Produção Animal*, Salvador, v. 15, n. 2, p. 350-360, 2014.
- STATISTICAL ANALYSIS SYSTEM INSTITUTE - SAS Institute. Statistical Analysis System for Windows. v. 9.0. Cary: SAS Institute, 2002.
- VELDKAMP, T.; TINÔCO, I. F. F.; BARRETO, S. L. T.; AMARAL, A. G.; PIRES, L. C.; FERREIRA, A. S. Growth responses to dietary energy and lysine at high and low ambient temperature in male turkeys. *Poultry Science*, Savoy, v. 84, n. 2, p. 273-282, 2005.
- WIDEMAN, R. F. J.; TACKETT, C. D. Cardio-pulmonary function in broilers reared at warm or cool temperatures: Effect of acute inhalation of 100% oxygen. *Poultry Science*, Savoy, v. 79, n. 72, p. 257-264, 2000.
- YAHAV, S.; STRASCHNOW, A.; LUGER, D.; SHINDER, D.; TANNY, J.; COHEN, S. Ventilation, sensible heat loss, broiler energy, and water balance under harsh environmental conditions. *Poultry Science*, Savoy, v. 83, n. 2, p. 253-258, 2004.
- YUNianto, V. D.; HAYASHI, K.; KANEDA, S.; OHTSUKA, A.; TOMITA, Y. D. Effect of environmental temperature on muscle protein turnover and heat production in tube-fed broiler chickens. *British Journal of Nutrition*, London, v. 77, n. 6, p. 897-909, 1997.