



Revista de Biología Tropical

ISSN: 0034-7744

rbt@cariari.ucr.ac.cr

Universidad de Costa Rica

Costa Rica

Ferré, Lilian E.; Medesani, Daniel A.; García, C. Fernando; Grodzielski, Matías; Rodríguez, Enrique M.
Vitellogenin levels in hemolymph, ovary and hepatopancreas of the freshwater crayfish *Cherax*
quadricarinatus (Decapoda: Parastacidae) during the reproductive cycle
Revista de Biología Tropical, vol. 60, núm. 1, marzo, 2012, pp. 253-261
Universidad de Costa Rica
San Pedro de Montes de Oca, Costa Rica

Available in: <http://www.redalyc.org/articulo.oa?id=44923251017>

- How to cite
- Complete issue
- More information about this article
- Journal's homepage in redalyc.org

redalyc.org

Scientific Information System
Network of Scientific Journals from Latin America, the Caribbean, Spain and Portugal
Non-profit academic project, developed under the open access initiative

Vitellogenin levels in hemolymph, ovary and hepatopancreas of the freshwater crayfish *Cherax quadricarinatus* (Decapoda: Parastacidae) during the reproductive cycle

Lilian E. Ferré¹, Daniel A. Medesani¹, C. Fernando García², Matías Grodzielski¹ & Enrique M. Rodríguez¹

1. Dept. of Biodiversity and Experimental Biology, DBBE-FCEyN, University of Buenos Aires, Argentina. Ciudad Universitaria, Pab. II, Intendente Guiraldes 2620, C1428EHA Buenos Aires, Argentina; ferredoxina@yahoo.com.ar, medesani@bg.fcen.uba.ar, mgrodzi@gmail.com, enrique@bg.fcen.uba.ar
2. Instituto de Investigaciones Bioquímicas de La Plata (INIBIOLP), CONICET-UNLP, 60 and 120 St., 1900 La Plata, Argentina; cfgarcia1123@yahoo.com.ar

Received 18-II-2011. Corrected 10-V-2011. Accepted 13-VI-2011.

Abstract: The freshwater crayfish *Cherax quadricarinatus* is a tropical species of great interest for aquaculture. Vitellogenin (Vg), a lipoprotein precursor of the vitellum accumulated in spawned eggs, can be synthesized in the ovary and/or hepatopancreas of most crustaceans, being the hemolymph the way for transporting Vg throughout the reproductive cycle. Concentration of Vg in hemolymph, ovary and hepatopancreas of *Cherax quadricarinatus* adult females was measured by means of ELISA, specifically developed after purifying the native Vg. Measurements were made at four periods of the reproductive cycle: pre-reproductive, mid-reproductive, late reproductive and post-reproductive. Besides, both hepatosomatic (HSI) and gonadosomatic (GSI) indexes were determined in each period. Significant variations in Vg levels were detected in both hemolymph and hepatopancreas, being the highest values observed during the mid-reproductive period. Besides, such variations were positively correlated to the HSI. A positive correlation between Vg levels in hepatopancreas and ovary was also seen. These results support previous evidences about the central role of the hepatopancreas as a site of Vg synthesis in the studied species, together with the relevancy of hemolymph for transporting Vg from the hepatopancreas to the ovary. For aquaculture purposes, Vg monitoring in hemolymph could be used as a non-injurious method, to check the reproductive activity of *C. quadricarinatus* females. Rev. Biol. Trop. 60 (1): 253-261. Epub 2012 March 01.

Keywords: crustaceans, reproduction, energy allocation, vitellogenin, reproductive cycle.

Vitellogenesis is a key process in the reproduction of decapod crustaceans. Vitellogenin (Vg) is a complex lipoprotein acting as a precursor of the vitellins used by embryos to build their tissues throughout the egg incubation period (Harrison 1990, Soroka *et al.* 2000, Tsukimura 2001, Abdu *et al.* 2002). During the primary or endogenous vitellogenesis, Vg is only synthesized by the oocytes, while during the secondary or exogenous vitellogenesis, although endogenous synthesis by oocytes continues, Vg synthesis mainly occurs in cells

other than oocytes, to be later up taken by them (Meusy 1980, Charmantier *et al.* 1997). Vg synthesized in extraovarian sites is transported to oocytes by hemolymph (Yehezkel *et al.* 2000, Tahara *et al.* 2005).

The freshwater crayfish *Cherax quadricarinatus* (von Martens 1868) is a tropical and subtropical species widely used in aquaculture. In tropical regions, a continuous spawning throughout the year has been reported, but in subtropical climates a reproductive cycle can be recognized (Barki *et al.* 1997), comprising

three basic periods: pre-reproductive (late June-late September), reproductive (late September-late March) and post-reproductive (late March-late June). Spawning mainly occurs during the reproductive period, usually being the post-reproductive a quiescent period, returning to the ovarian maturation during the pre-reproductive one (Jones & Ruscoe 1996, Barki *et al.* 1997).

Determination of the relative ovarian growth (gonadosomatic index), is a methodology commonly used for evaluating the ovarian growth as a final point of an experimental assay. However, since vitellogenin circulating levels have shown to be correlated to the reproductive condition of several crustaceans (Tahara *et al.* 2005, Ibarra *et al.* 2009), their detection and quantification could be an useful tool for monitoring the reproductive state of *C. quadricarinatus* females, during their entire reproductive cycle. The advantage of using this kind of non-injurious technique in aquaculture programs is evident.

The ELISA (enzyme-linked immunosorbent assay) technique is considered as a sensitive and precise method to quantify lipoprotein compounds, such as vitellins or vitellogenin (Specker & Anderson 1994). Vitellogenin levels have been previously measured in several crustacean species, either in hemolymph or other tissues (Lee & Chang 1997, Pateraki & Stratakis 2000, Tsukimura 2001, Vazquez Boucard *et al.* 2002, Chen *et al.* 2004, Tahara *et al.* 2005, García *et al.* 2006, Santhoshi *et al.* 2009, Ibarra *et al.* 2009). In mature females of *C. quadricarinatus*, both the ovary and hepatopancreas have been reported as the main sites for vitellogenin synthesis (Serrano Pinto *et al.* 2003, 2004, 2005). Besides, the presence of vitellogenin in hemolymph has been associated to the secondary vitellogenesis that takes place in the ovary (Yehezkel *et al.* 2000, Abdu *et al.* 2002). Vitellogenin levels have been previously quantified in *C. quadricarinatus* by ELISA, at the onset of secondary vitellogenesis (Sagi *et al.* 1999). However, no monitoring of Vg hemolymphatic levels throughout the reproductive cycle of this species has been done.

The current study was aimed at determining the vitellogenin levels in hemolymph, ovary and hepatopancreas of *C. quadricarinatus* at the different periods of its reproductive cycle, by means of the ELISA technique.

MATERIALS AND METHODS

Adult females of *C. quadricarinatus* (overall mean weight=43.67±11.8g, N=38) were obtained during 2009 from a local dealer (Pinzas Rojas S.R.L, Tucumán, Argentina). Once in the laboratory, the animals were maintained for one week in a glass aquarium (40x60cm glass bottom, 15L capacity), at a density of five females per aquarium, each one filled with dechlorinated tap water (hardness: 80mg/L as CaCO₃ equivalents), under constant aeration. A temperature corresponding to the mean value for each considered period of the reproductive cycle, as well as a natural photoperiod, was maintained. Animals were daily fed *ad libitum* with a commercial pellet (Tetra^R) having 32% protein, and leaves of *Elodea canadensis*.

After this acclimation period to laboratory conditions, females were weighed and a sample of hemolymph (100µL) was withdrawn from the base of the fifth pair of pereopods, by means a 1mL syringe, provided with a 25G needle. Hemolymph samples were then transferred to Eppendorf tubes containing 15µL of potassium oxalate 10% and protease inhibitors (PMSF 0.01M) in a 3:1 proportion (v/v), to be finally freeze at -70°C until analysis by ELISA. Females were then cold-anaesthetized and sacrificed, both ovary and hepatopancreas were carefully dissected, and both gonadosomatic index (GSI) and hepatosomatic index (HSI) were finally calculated as (weight of gonad or hepatopancreas/body weight)x100. Both tissues were subsequently cut in small fragments (0.1 to 0.2g), which were homogenized in sodium phosphate buffer (50mM, pH=7.4, with 2µL/mL of protease inhibitor), in a 1:3 (w/v) ratio. Each homogenate was then centrifuged at 10 000g for 20min, at 4°C. Supernatants were further ultra-centrifuged (100 000g for 50min, at 4°C); the resulting supernatants were

separated in Eppendorf tubes and freeze until -70°C until analysis by ELISA.

Biological samples were taken at the following periods of the reproductive cycle:

Pre-reproductive: July 16 (N=10)

Mid-reproductive: December 9 (N=8)

Late-reproductive: February 10 (N=10)

Post-reproductive: June 2 (N=10)

Vitellogenin (Vg) purification process:

Isolation and purification of Vg was made from three females of *C. quadricarinatus* having GSI values higher than two, i.e., with completely mature ovaries, according to Abdu *et al.* (2002). The procedure followed that described by García *et al.* (2008) for *Macrobrachium borellii*. Briefly, ovaries from three females were pooled and homogenized in PBS 20mM, pH=7.4, containing protease inhibitor (PMSF) at 2µL/mL. Homogenate was then centrifuged at 10 000g for 20min, and the supernatant was further centrifuged at 100 000g for 60min. Aliquots of cytosol were overlayed on NaBr (density 1.26g/mL) containing 0.01% sodium azide, to be finally centrifuged in a vertical gradient, by means of a Beckman L8 70M centrifuge provided with a SW60 Ti rotor, at 178 000g and 10°C for 24h. A saline solution of the same density to that of samples was centrifuged in parallel to determine relative densities, and check the correct gradient formation. The total volume of the tubes was fractionated from top into 0.2mL aliquots, and the protein content of each fraction was monitored spectrophotometrically at 280nm. The protein peak containing the lipoproteins was separated as a whole fraction, measuring the total protein content by the method of Lowry *et al.* (1951). All samples were frozen at -70°C until analysis.

Gel electrophoresis: Lipoprotein was analyzed by native PAGE using a gradient of 4-23% acrylamide (Laemmli 1970), and

stained with Coomassie Brilliant Blue R-250 (Sigma Chemical Co., St. Louis, MO).

Procedure to obtain antibody against native Vg and Enzymelinked immunosorbent assay (ELISA) development: A primary antibody against Vg was obtained by inoculating rabbits with purified Vg, according to previous studies (Dreon *et al.* 2003, García *et al.* 2008). Anti IgG from rabbit, conjugated with Bio-Rad Lab. Peroxidase, was used as the secondary antibody. Purified Vg in a 1/500 dilution was used to prepare the standard (0 to 300ng). 15µL of either the standard or sample were placed, in triplicate, in a 96-wells plate (Nunc-Immunoplate Polisorp). Samples were previously diluted in coating buffer (15mM sodium carbonate, 35mM sodium bicarbonate, pH=9.6). Both primary and secondary antibodies were diluted (1/500) in PBS-0.05% Tween-6% powder milk. Absorbance was measured in all wells at 415nm, by using an ELISA-plates reader (Bio-Rad Lab., Model 680). ABTS [2-2'-azino-di-(3-ethylbenzthiazoline sulfonic acid)] was used as chromogen.

GSI, HSI and Vg concentration (µg/g), both in ovary and hepatopancreas, were analyzed by means of a one way ANOVA (considering period as factor), followed by the Tuckey test to compare mean values by pairs (Sokal & Rohlf 1981). Correlation between variables was also estimated, testing the significance of each correlation made (Sokal & Rohlf 1981). A 5% confidence level was considered.

RESULTS

Figure 1 shows the result obtained by electrophoretic analysis of lipoprotein purified from mature ovaries of *C. quadricarinatus* under native conditions. Electrophoretic mobility revealed a protein band of around 500kDa.

Table 1 shows the mean values of both GSI and HSI, together with the Vg concentration in hemolymph, ovary and hepatopancreas, for each period of the reproductive cycle of *C. quadricarinatus* adult females. No significant differences ($p>0.05$) were seen among periods,




Fig. 1. Results of native PAGE for lipovitellins of *C. quadricarinatus* purified from ovaries of mature females (Sample 1 indicates 6mg/L). St: biomarkers of known molecular weight (kDa).

for either GSI or ovarian Vg level. On the contrary, HSI values showed significant differences ($p<0.05$) comparing the pre-reproductive period to either the mid or late reproductive period; significant differences ($p<0.01$) in HSI

were also noted between mid and late reproductive periods, and between this later period and the post-reproductive one. Hepatopancreatic Vg showed a significant ($p<0.05$) lower level at the late reproductive period, with respect to the remaining periods. The lowest hemolymphatic Vg levels were also observed during the late reproductive period, compared to any other period ($p<0.01$).

The result of the correlation made by pairs of variables throughout the entire reproductive cycle is shown in Figures 2 and 3. As Vg level increases in hepatopancreas, an increasing Vg level was also observed in ovary ($p<0.05$, Fig. 2A). On the other hand, a positive correlation ($p<0.01$) between hepatopancreatic Vg and HSI was observed (Fig. 2B). Figures 3A and 3B show how the circulating levels of Vg positively correlate ($p<0.01$) to either Vg hepatopancreatic level or HSI.

DISCUSSION

Purification of lipoproteins from *C. quadricarinatus* mature ovaries showed to be successful, i.e., a discrete band with a molecular weight near 500kDa was obtained. This band closely corresponds to one of the main vitellin forms identified in mature ovaries of *C. quadricarinatus* by Serrano Pinto *et al.* (2003). A vitellogenin with a molecular weight by the 440kDa was observed for the vitellogenin purified from *M. borellii* (García *et al.* 2006). Crustacean vitellin molecular mass has been reported as ranging from near 320 to 630kDa, for a wide list of species (Tsukimura 2001).

TABLE 1
Gonadosomatic (GSI) and hepatosomatic (HSI) indexes, as well as mean levels of vitellogenin (Vg) in hemolymph (HL), ovary (OV) and hepatopancreas (HP) of *C. quadricarinatus* adult females, measured at different periods of the reproductive cycle

Periods	GSI	HSI	VgHL(ng/ μ L)	VgOV(μ g/g)	VgHP(μ g/g)	N
Pre-reproductive	1.66 \pm 0.14 (a)	6.39 \pm 0.19 (a)	313.49 \pm 10.31 (a)	1015.28 \pm 15.93 (a)	839.44 \pm 90.49 (a)	9
Mid-reproductive	1.57 \pm 0.40 (a)	8.20 \pm 0.56 (b)	457.33 \pm 41.27 (b)	983.04 \pm 309.56 (a)	962.24 \pm 209.46 (a)	8
Late-reproductive	2.00 \pm 0.52 (a)	4.93 \pm 0.31 (c)	221.32 \pm 4.24 (c)	1005.59 \pm 161.82 (a)	315.013 \pm 38.92 (b)	10
Post- reproductive	1.33 \pm 0.16 (a)	7.26 \pm 0.26 (ab)	302.69 \pm 5.13 (a)	875.85 \pm 133.86 (a)	804.76 \pm 91.41 (a)	10

Mean value \pm standard error is indicated. Different letters indicate significant ($p<0.05$) differences among periods.




Fig. 2. Significant ($p < 0.05$) correlation of vitellogenin (Vg) level in hepatopancreas (HP) with (A) Vg level in ovary (OV) or (B) hepatosomatic index (HSI), during the entire reproductive cycle of *C. quadricarinatus*.

Lipoproteins associated to the vitellogenesis process (i.e., vitellogenin and vitellins) have been identified and characterized in several crustacean species, and they were used as a tool for studying the reproductive biology of such species (Quackenbush 1994, Oberdörster *et al.* 2000, Pateraki & Stratakis 2000, Okumura 2004, Chen *et al.* 2004, García *et al.* 2006, 2008). In several decapod crustaceans used in aquaculture, Vg circulating levels have

been quantified. In some species of decapod crustaceans no correlation between hemolymphatic Vg levels and ovarian development was observed (Lee & Chang 1997). However, in most cases a good correlation between Vg circulating levels and ovarian growth was seen in females of reproductive age (Tsukimura 2001, Tahara *et al.* 2005, Ibarra *et al.* 2009, Santhoshi *et al.* 2009), although Vg circulating levels decreases in some extent towards




Fig. 3. Significant ($p < 0.05$) correlation of vitellogenin (Vg) level in hemolymph (HL) with (A) Vg level in hepatopancreas (HP) or (B) hepatosomatic index (HSI), during the entire reproductive cycle of *C. quadricarinatus*.

the final ovarian maturation, i.e., just before the oviposition takes place (Tsukimura 2001, Tahara *et al.* 2005).

Vitellins of *C. quadricarinatus* have been previously purified and characterized in ovary and eggs of mature females (Serrano Pinto *et al.* 2003), while both the hepatopancreas and the ovary have been characterized as the main sites of vitellogenin synthesis for the same

species (Serrano Pinto *et al.* 2004, 2005). Abdu *et al.* (2002) have cloned the complete vitellogenin cDNA of *C. quadricarinatus*, while Yehezkel *et al.* (2000) have identified specific lipoproteins in the hemolymph of this species, associated to the onset of secondary vitellogenesis. An ELISA for detecting both vitellogenin and vitellins has been previously developed by Sagi *et al.* (1999), specifically

for *C. quadricarinatus*. However, no previous information about Vg fluctuations during the entire reproductive cycle is available for this species.

Results of the current study represent a first report, in the studied species, of the annual variation of Vg levels in hemolymph, ovary and hepatopancreas, and their correlation to both GSI and HSI. Significant variations were detected in both hemolymphatic and hepatopancreatic Vg levels, which were highest during the mid-reproductive period. Besides, such variations were positively correlated to the HSI, during the entire cycle. The increment of Vg concentration in hepatopancreas would be responsible for an enhanced HSI. On the other hand, a higher synthesis of Vg in hepatopancreas would be causing an increase in the Vg circulating level, stressing the relevance of hemolymph as a transporting way for extra-ovarian Vg, from its site of synthesis to the ovary. Although differences in both Vg ovarian levels (in terms of $\mu\text{g/g}$) and GSI could not be detected throughout the reproductive cycle, the significant positive correlation found in Vg levels between ovary and hepatopancreas was in accordance with the expected higher uptake of Vg by the former tissue, as increases the production by the later.

As mentioned before, the role of the hepatopancreas as the extraovarian site of vitellogenin has been previously verified for *C. quadricarinatus* (Serrano Pinto *et al.* 2005). This hepatopancreatic function has been also reported for other crustaceans (Quackenbush 1994, Lee & Chang 1997), although in the case of the shrimp *Fenneropenaeus indicus*, a reduced contribution of the hepatopancreas to the ovarian growth was suggested (Vazquez Boucard *et al.* 2002). According to Serrano Pinto *et al.* (2003, 2005), the ovary of *C. quadricarinatus* could be playing a relevant role in Vg production only in first-maturation females, but not in previously spawners, whose hepatopancreas would be the main site of Vg production. Since the minimum size at maturity for *C. quadricarinatus* was reported as 15g (Vazquez & López Greco 2007), females used

in the current study (averaging 44g) were probably multiparous, i.e., they have had previous spawning. Therefore, the changes observed in their Vg hepatopancreatic and hemolymphatic levels would be in accordance with the central role of the hepatopancreas as a site of Vg synthesis, suggested by Serrano Pinto *et al.* (2003, 2005) for this kind of females.

Finally, the ELISA technique developed during the current study has shown to be a sensitive and useful method to detect, through a practical and non-injurious method, the degree of reproductive development at which females of the studied species can be found at different period of the reproductive cycle. In fact, hemolymphatic level of Vg has been previously used as biomarker for estimating the reproductive activity of crustaceans (Vazquez Boucard *et al.* 2002, Tahara *et al.* 2005). Vg circulating level could be also taken as biomarker to evaluate the effect of different diets, hormones and neuroregulators, among other factors, to the reproductive success, in order to optimize the culture of this species.

ACKNOWLEDGMENTS

Authors wish to help the financial support given by grants from both the University of Buenos Aires UBACYT program (X241) and the Argentine National Council of Scientific and Technical Research (CONICET, PIP 2010-2012, code 100884).

RESUMEN

La langosta de agua dulce *Cherax quadricarinatus* es una especie tropical de gran interés para la acuicultura. Se midió la concentración de vitelogenina (Vg) en hemolinfa, ovario y hepatopáncreas de hembras adultas de esta especie, por medio de ELISA. Las mediciones fueron hechas en los cuatro períodos del ciclo reproductivo: pre-reproductivo, reproductivo medio, reproductivo tardío y post-reproductivo. Se detectaron variaciones significativas en los niveles de Vg tanto en hemolinfa como en hepatopáncreas, se observó el mayor valor durante el período reproductivo medio. Además, tales variaciones se correlacionaron positivamente con el índice hepatosomático. Se observó además una correlación positiva de los niveles de Vg entre hepatopáncreas y ovario. Estos

resultados apoyan evidencias previas sobre el papel central del hepatopáncreas como sitio de síntesis de Vg, en esta especie, y también enfatizan la importancia de la hemolinfa para el transporte de la Vg del hepatopáncreas al ovario. Para propósitos de acuicultura, la medición de Vg en hemolinfa podría ser utilizada como un método no lesivo, con el fin de constatar la actividad reproductiva de hembras de *C. quadricarinatus*.

Palabras clave: crustáceos, reproducción, inversión energética, vitelogenina, ciclo reproductivo.

REFERENCES

- Abdu, V., C. Davis, I. Khalaila & A. Sagi. 2002. The vitellogenin cDNA of *Cherax quadricarinatus* encodes a lipoprotein with calcium binding ability and its expression is induced following the removal of the androgenic gland in a sexually plastic system. *Gen. Comp. Endocrinol.* 127: 263-272.
- Barki, A., T. Levi, G. Hulata & I. Karplus. 1997. Annual cycle of spawning and molting in the red-claw crayfish, *Cherax quadricarinatus*, under laboratory conditions. *Aquaculture* 157: 239-249.
- Charmantier, G., M. Charmantier-Daures & F. Van Herp. 1997. Hormonal regulation of growth and reproduction in crustaceans, p. 109-161. In M. Fingerman, R. Nagabhushanam & M.F. Thompson (eds.). *Recent advances in marine biotechnology*, vol. I. Science, Enfield, New Hampshire, USA.
- Chen, L., H. Jiang, Z. Zhou, K. Li & G.Y. Deng. 2004. Purification of vitellin from the ovary of chinese mitten-handed crab (*Eriocheir sinensis*) and development of an antivitelin ELISA. *J. Comp. Physiol. B. Biochem. Syst. Environ. Physiol.* 138: 305-311.
- Dreon, M.S., H. Heras & R.J. Pollero. 2003. Metabolism of ovorubin, the major egg lipoprotein from the apple snail. *Mol. Cell. Biochem.* 243: 9-14.
- García, C.F., M. Cunningham, J.L. Soluages, H.A. Garda & R. Pollero. 2006. Structural characterization of the lipovitellin from de shrimp *Macrobrachium borellii*. *J. Comp. Physiol. B. Biochem. Syst. Environ. Physiol.* 145: 365-370.
- García, F., M.L. Cunningham, H. Garda & H. Heras. 2008. Embryo lipoproteins and yolk lipovitellin consumption during embryogenesis in *Macrobrachium borellii* (Crustacea: Palaemonidae). *J. Comp. Physiol. B. Biochem. Syst. Environ. Physiol.* 151: 317-322.
- Harrison, K.I. 1990. The role of nutrition in maturation, reproduction and embryonic development of decapod crustaceans: a review. *J. Shellfish Res.* 9: 1-28.
- Jones, C.M. & I. Ruscoe. 1996. Production technology for Redclaw Crayfish (*Cherax quadricarinatus*). Final Report FRDC Project 92/119. Fisheries Research and Development Corporation, Camberra, Australia.
- Ibarra, A.M., T.R. Famula & F.G. Arcos. 2009. Heritability of vitellogenin in hemolymph, a pre-spawning selectable trait in *Penaeus (Litopenaeus) vannamei*, has a large genetic correlation with ovary maturity measured as oocytes mean diameter. *Aquaculture* 297: 64-69.
- Laemmli, U.K. 1970. Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature* 227: 680-685.
- Lee, F.Y. & C.F. Chang. 1997. The concentrations of vitellogenin (vitellin) and protein in hemolymph, ovary and hepatopancreas in different ovarian stages of the freshwater prawn *Macrobrachium rosenbergii*. *Comp. Biochem. Physiol. Mol. Integr. Physiol.* 117: 433-439.
- Lowry, O.H., N.J. Rosebrough, A.L. Farr & R.J. Randall. 1951. Protein measurement with the Folin phenol reagent. *J. Biol. Chem.* 193: 265-275.
- Meusy, J.J. 1980. Vitellogenin, the extra ovarian precursor of the protein yolk in Crustacea: a review. *Reprod. Nutr. Dev.* 20: 1-21.
- Oberdörster, E., C.D. Rice & L.K. Irwin. 2000. Purification of vitellin from grass shrimp *Palaemonetes pugio* generation of monoclonal antibodies, and validation for the detection of lipovitellin in crustacean. *Comp. Biochem. Physiol. C. Pharmacol. Toxicol. Endocrinol.* 127: 199-207.
- Okumura, T. 2004. Perspectives on hormonal manipulation of shrimp reproduction. *JARQ* 38: 49-54.
- Pateraki, L. & E. Stratakis. 2000. Synthesis and organization of vitellogenin and vitellin molecules from the land crab *Potamon potamios*. *J. Comp. Physiol. B. Biochem. Syst. Environ. Physiol.* 125: 53-61.
- Quackenbush, L.S. 1994. Lobster reproduction: a review. *Crustaceana* 67: 82-94.
- Sagi, A., I. Khalaila, U. Abdu, R. Shoukrum & S. Weil. 1999. A new established ELISA showing the effect of the androgenic gland on secondary -vitellogenin-specific protein in the hemolymph of the crayfish *Cherax quadricarinatus*. *Gen. Comp. Endocrinol.* 115: 37-45.

- Santoshi, S., V. Sugumar & N. Munuswamy. 2009. Serotonergic stimulation of ovarian maturation and hemolymph vitellogenin in the Indian white shrimp, *Fenneropenaeus indicus*. *Aquaculture* 291: 192-199.
- Serrano-Pinto, V., C. Vazquez-Boucard & H. Villareal-Colmenares. 2003. Yolk proteins during ovary and egg development of mature female freshwater crayfish (*Cherax quadricarinatus*). *Comp. Biochem. Physiol. Mol. Integr. Physiol.* 134: 33-43.
- Serrano-Pinto, V., I. Landais, M.H. Ogliastro, M. Gutierrez Ayala, H. Mejía Ruíz, H. Villareal Colmenares, A. García Gasca & C. Vazquez Boucard. 2004. Vitellogenin mRNA during secondary vitellogenic at first maturation females. *Mol. Reprod. Dev.* 69: 17-21.
- Serrano-Pinto, V., M.G. Carrisoza-Valenzuela & M. Ramírez-Orozco. 2005. Determination site of vitellogenin synthesis in freshwater crayfish *Cherax quadricarinatus* at different maturation stages females. *Investigaciones Marinas* 33: 195-200.
- Sokal, R.R. & F.J. Rohlf. 1981. *Biometry*. Freeman, New York, USA.
- Soroka, Y., A. Sagi, I. Khalaila, U. Abdu & Y. Milner. 2000. Changes in protein kinase C during vitellogenesis in the crayfish *Cherax quadricarinatus*, possible activation by methyl farnesoate. *Gen. Comp. Endocrinol.* 118: 200-208.
- Specker, J.L. & T.R. Anderson. 1994. Developing and ELISA for a model protein-vitellogenin, p. 567-578. *In* P. Hochachka & T. Mommsen (eds.). *Biochemistry and molecular biology of fishes*, vol. 3. The Netherlands: Elsevier Science B.V. Amsterdam, Netherlands.
- Tahara, D., K. Suitoh & H. Hattori. 2005. Hemolymph vitellogenin levels during final maturation and post-spawning in the female kuruma prawn *Marsupenaeus japonicus*. *Aquaculture* 245: 311-319.
- Tsukimura, B. 2001. Crustacean Vitellogenesis: its role in oocyte development. *Am. Zool.* 41: 465-475.
- Vazquez, F.J. & L.S. López Greco. 2007. Diferenciación sexual en la langosta de agua dulce *Cherax quadricarinatus* (Decapoda, Parastacidae). *Rev. Biol. Trop.* 55: 33-38.
- Vazquez Boucard, C.G., P. Levy, H.J. Ceccaldi & C.H. Brogren. 2002. Development changes in concentrations of vitellin, vitellogenin and lipids in hemolymph, hepatopancreas and ovaries from different ovarian stages of indian white prawn *Fenneropenaeus indicus*. *J. Exp. Mar. Biol. Ecol.* 281: 63-75.
- Yehezkel, G., R. Chayoth, U. Abdu, I. Khalaila & A. Sagi. 2000. High-density lipoprotein associated with secondary vitellogenesis in the hemolymph of the crayfish *Cherax quadricarinatus*. *J. Comp. Physiol. B. Biochem. Syst. Environ. Physiol.* 127: 411-421.