



Revista de Matemática: Teoría y Aplicaciones

ISSN: 1409-2433

mta.cimpa@ucr.ac.cr

Universidad de Costa Rica

Costa Rica

López-Bonilla, José; Cruz-Beltrán, L.

An approach to ladder operators for the two-dimensional harmonic oscillator

Revista de Matemática: Teoría y Aplicaciones, vol. 15, núm. 1, enero-junio, 2008, pp. 97-99

Universidad de Costa Rica

San José, Costa Rica

Available in: <http://www.redalyc.org/articulo.oa?id=45326937008>

- How to cite
- Complete issue
- More information about this article
- Journal's homepage in redalyc.org

redalyc.org

Scientific Information System

Network of Scientific Journals from Latin America, the Caribbean, Spain and Portugal

Non-profit academic project, developed under the open access initiative

AN APPROACH TO LADDER OPERATORS FOR THE TWO-DIMENSIONAL HARMONIC OSCILLATOR

JOSÉ LÓPEZ-BONILLA* L. CRUZ-BELTRÁN†

Recibido/Received: 14 May 2007 — Aceptado/Accepted: 31 Oct 2007

Abstract

We show that elementary properties of associated Laguerre polynomials generate the ladder operators for the harmonic oscillator radial wave function in two dimensions.

Keywords: Harmonic oscillator, ladder operators, Laguerre polynomials.

Resumen

Demostramos que propiedades elementales de polinomios de Laguerre generan operadores escalera para la función de onda del oscilador radial armónico en dos dimensiones.

Palabras clave: Oscilador armónico, operadores escalera, polinomios de Laguerre.

Mathematics Subject Classification: 33C45; 34B30.

1 Introduction

Let $\psi_{MN}(\varrho, \varphi)$ be the wave function of the two-dimensional harmonic oscillator (*2DHO*) in polar coordinates with natural units ($\hbar = 1$ and Mass=1), which can be expressed in its radial and angular parts as [1]:

$$\psi_{MN} = R_{|M|N}(\varrho)e^{iM\varphi}, \quad N = 0, 1, 2, \dots \quad (1)$$

*Sección de Estudios de Posgrado e Investigación Escuela Superior de Ingeniería Mecánica y Eléctrica
Instituto Politécnico Nacional Edif. Z-4, 3er Piso, Col. Lindavista, C.P. 07738 México, D.F., México.
E-Mail: jlopezb@ipn.mx

†Misma dirección de J. López. E-Mail: beltranb2000@yahoo.com

with $M = -N, -N + 2, \dots, N - 2, N$ where

$$R_{MN} = C_{mN} \varrho^m e^{-\frac{\varrho^2}{2}} L_{\frac{N-m}{2}}^m(\varrho^2), \quad C_{mN} = \left[\frac{(\frac{N-m}{2})!}{\pi (\frac{N+m}{2})!} \right]^{\frac{1}{2}}, \quad m \geq 0 \quad (2)$$

and L_q^p denoting the associated Laguerre polynomials [2].

Here we shall determine the ladder operators \hat{O}_{mN}^\pm , for the radial wave function(2), such that:

$$R_{m\pm 2, N} = \hat{O}_{mN}^\pm R_{mN}, \quad (3)$$

employing only elementary properties of L_q^p , that is, it is possible to construct \hat{O}_{mN}^\pm without the use of specific techniques as the factorization method [3-6].

2 Ladder operators for the 2DHO

From (2) we have that:

$$R_{m+2, N} = \frac{C_{mN} \varrho^{m+2}}{\left[\left(\frac{N-m}{2} \right) \left(\frac{N+m}{2} + 1 \right) \right]^{\frac{1}{2}}} e^{-\frac{\varrho^2}{2}} L_{\frac{N-m}{2}-1}^{m+2}(\varrho^2), \quad (4)$$

but if we remember the known relations [2]:

$$\frac{d}{d\mu} L_\beta^\alpha(\mu) = -L_{\beta-1}^{\alpha+1}(\mu), \quad L_\gamma^{\beta+1}(\mu) = \left(1 - \frac{d}{d\mu} \right) L_\gamma^\beta(\mu), \quad (5)$$

then (4) may be written as:

$$R_{m+2, N} = -\frac{C_{mN} \varrho^{m+1}}{\sqrt{(N-m)(N+m+2)}} e^{-\frac{\varrho^2}{2}} \frac{d}{d\varrho} \left(1 - \frac{1}{2\varrho} \frac{d}{d\varrho} \right) L_{\frac{N-m}{2}}^m(\varrho^2), \quad (6)$$

From the associated Laguerre equation[2] it is immediate the expression:

$$\frac{d^2}{d\varrho^2} L_{\frac{N-m}{2}}^m(\varrho^2) = \left(2\varrho - \frac{1+2m}{\varrho} \right) \frac{d}{d\varrho} L_{\frac{N-m}{2}}^m(\varrho^2) - 2(N-m) L_{\frac{N-m}{2}}^m(\varrho^2), \quad (7)$$

and thus (6) adopts the form:

$$R_{m+2, N} = -\frac{C_{mN} \varrho^m e^{-\frac{\varrho^2}{2}}}{\sqrt{(N-m)(N+m+2)}} \left[\frac{(1+m)}{\varrho} \frac{d}{d\varrho} L_{\frac{N-m}{2}}^m(\varrho^2) + (N-m) L_{\frac{N-m}{2}}^m(\varrho^2) \right], \quad (8)$$

but (2) implies that $L_{\frac{N-m}{2}}^m(\varrho^2) = \frac{\varrho^{-m}}{C_{mN}} e^{\frac{\varrho^2}{2}} R_{mN}$, then (8) leads to (3) with:

$$\hat{O}_{mN}^+ = \frac{m+1}{\sqrt{(N-m)(N+m+2)}} \left(\frac{m}{\varrho^2} - \frac{N+1}{m+1} - \frac{1}{\varrho} \frac{d}{d\varrho} \right) R_{mN}. \quad (9)$$

A similar process permits to deduce the other ladder operator:

$$\hat{O}_{mN}^- = \frac{m-1}{\sqrt{(N+m)(N-m+2)}} \left(\frac{m}{\varrho^2} - \frac{N+1}{m-1} + \frac{1}{\varrho} \frac{d}{d\varrho} \right) R_m N. \quad (10)$$

The radial wave function for the Coulomb potential is in terms of L_q^p , to see [7,8], the method here showed is applicable to obtain its ladder operators, which represents an alternative procedure to another techniques [9,10].

References

- [1] Wallace, P.R. (1984) *Mathematical Analysis of Physical Problems*. Dover, New York.
- [2] Abramowitz, M.; Stegun, I.A. (1972) *Handbook of Mathematical Functions*. John Wiley and Sons, New York.
- [3] Infeld, L. (1942) “A generalization of the factorization method of solving eigenvalue problems”, *Trans. Canad. Roy. Soc. Ser. III* **36**: 7–18.
- [4] Infeld, L.; Hull, T.E. (1948) “The factorization method, hydrogen intensities and related problems”, *Phys. Rev.* **74**: 905–909.
- [5] Infeld, L. (1949) “The factorization method and its application to differential equations in Theoretical Physics”, *Proc. Symp. Appl. Math.* **28**: 58–65.
- [6] Infeld, L.; Hull, T.E. (1951) “The factorization method”, *Rev. Mod. Phys.* **23**: 21–68.
- [7] López-Bonilla, J. ; Morales, J.; Rosales, M. (1995) “Hypervirial theorem and matrix elements for the Coulomb potential”, *Int. J. Quantum Chem.* **53**: 3–7.
- [8] Bautista-Moedano, M.; López Bonilla, J.; Morales, J. (2006) “Matrix elements $\langle n_2 \ell_2 | r^k | n_1 \ell_1 \rangle$ for the Coulomb interaction”, *Apeiron* **13**(1): 34–42.
- [9] Gaftoi, V.; López Bonilla, J.; Peña, J.J. (2000) “Ladder operators for the Coulomb potential”, *Indian J. Phys.* **B74**: 171–172.
- [10] Martínez, R.P.; Romero, Y.; Núñez-Yépez, H.N.; Salas-Brito, A.L. (2007) “Algebraic approach to radial ladder operators in the hydrogen atom”, *Int. J. Quantum Chem.* **107**: 1606–1613.