



Revista de Matemática: Teoría y Aplicaciones

ISSN: 1409-2433

mta.cimpa@ucr.ac.cr

Universidad de Costa Rica

Costa Rica

Caro-Lopera, Francisco J.; Díaz-García, José A.; González-Farías, Graciela

A formula for complex zonal polynomials of second order

Revista de Matemática: Teoría y Aplicaciones, vol. 13, núm. 1, enero, 2006, pp. 35-39

Universidad de Costa Rica

San José, Costa Rica

Available in: <http://www.redalyc.org/articulo.oa?id=45326957004>

- How to cite
- Complete issue
- More information about this article
- Journal's homepage in redalyc.org

redalyc.org

Scientific Information System

Network of Scientific Journals from Latin America, the Caribbean, Spain and Portugal

Non-profit academic project, developed under the open access initiative

A FORMULA FOR COMPLEX ZONAL POLYNOMIALS OF SECOND ORDER

FRANCISCO J. CARO-LOPERA ^{*} JOSÉ A. DÍAZ-GARCÍA[†]
GRACIELA GONZÁLEZ-FARÍAS[‡]

Recibido/Received: 22 Nov 2005 — Aceptado/Accepted: 30 Ago 2006

Abstract

A formula for complex zonal polynomials of second order is derived by solving a particular partial differential equation.

Keywords: Laplace-Beltrami operator, zonal polynomials, Hermitian matrix, Legendre's differential equation.

Resumen

En el presente trabajo se encuentra una fórmula para polinomios zonales complejos de segundo orden, a partir de una ecuación diferencial parcial.

Palabras clave: Operador de Laplace-Beltrami, polinomios zonales, matriz hermitiana, ecuación diferencial de Legendre.

Mathematics Subject Classification: 62H05, 33A70.

1 Introduction

Recently Díaz-García and Caro-Lopera (2006b) computed the zonal polynomials of positive definite hermitian matrix by the use of the Laplace-Beltrami operator. In the same

^{*}Department of Basic Mathematics, Centro de Investigación en Matemáticas, Callejón de Jalisco s/n, Mineral de Valenciana, 36240 Guanajuato, Guanajuato, México. E-mail: fjcaro@cimat.mx

[†]Department of Statistics and Computation, Universidad Autónoma Agraria Antonio Narro, 25315 Buenavista, Saltillo, Coahuila, México, E-Mail: jadiaz@uaaan.mx

[‡]Department of Probability and Statistics, Centro de Investigación en Matemáticas, Callejón de Jalisco s/n, Mineral de Valenciana, 36240 Guanajuato, Guanajuato, México, E-Mail: farias@cimat.mx

way for the real case, the zonal polynomials of positive definite and a semi-definite positive symmetric matrices were calculated by James (1968) and Díaz-García and Caro-Lopera (2006a), respectively. It is known that general formulae for complex and real zonal polynomials are not available, and the general partial differential equations for both polynomials are also unsolved. However, the differential equation for the zonal polynomials of definite positive symmetric matrix argument of the second order was solved by James (1968).

In this paper we reduce the partial differential equation for zonal polynomials of positive definite hermitian matrix of second order to a Hypergeometric differential equation type, which is analogous to the results obtained by James (1968) in the real case. By solving the ordinary differential equation, we get an explicit formula for the corresponding zonal polynomials, see Section 2.

2 Complex Zonal Polynomials of Second Order.

Let the partition $\kappa = (k_1, \dots, k_m)$ of k a decreasing sequence of nonnegative integers. In Díaz-García and Caro-Lopera (2006b) was proved that the zonal polynomials $\tilde{C}_\kappa(Y)$ of an $m \times m$ positive definite hermitian matrix Y satisfy the partial differential equation

$$\sum_{i=1}^m y_i^2 \frac{\partial^2}{\partial y_i^2} \tilde{C}_\kappa(Y) + 2 \sum_{i=1}^m \sum_{j=1(j \neq i)}^m y_i^2 (y_i - y_j)^{-1} \frac{\partial}{\partial y_i} \tilde{C}_\kappa(Y) = [\tilde{\rho}_\kappa + k(2m - 1)] \tilde{C}_\kappa(Y), \quad (1)$$

where

$$\tilde{\rho}_\kappa = \sum_{i=1}^m k_i(k_i - 2i),$$

y_1, \dots, y_m are the eigenvalues of the matrix Y and $\kappa = (k_1, \dots, k_m)$ is a partition of k .

When $m = 2$ in (1) we get the partial differential equation

$$y_1^2 \frac{\partial^2 \tilde{C}}{\partial y_1^2} + y_2^2 \frac{\partial^2 \tilde{C}}{\partial y_2^2} + 2y_1^2 (y_1 - y_2)^{-1} \frac{\partial \tilde{C}}{\partial y_1} - 2y_2^2 (y_1 - y_2)^{-1} \frac{\partial \tilde{C}}{\partial y_2} - [k_1(k_1 + 1) + k_2(k_2 - 1)] \tilde{C} = 0, \quad (2)$$

where we denote $\tilde{C}_\kappa(Y)$ as \tilde{C} .

Let us replace $u = y_1 + y_2$ and $v = y_1 y_2$ in (2), then we find

$$(u^2 - 2v) \frac{\partial^2 \tilde{C}}{\partial u^2} + 2v^2 \frac{\partial^2 \tilde{C}}{\partial v^2} + 2uv \frac{\partial^2 \tilde{C}}{\partial u \partial v} + 2u \frac{\partial \tilde{C}}{\partial u} + 2v \frac{\partial \tilde{C}}{\partial v} - [k_1(k_1 + 1) + k_2(k_2 - 1)] \tilde{C} = 0.$$

Substituting $z = \frac{u}{2\sqrt{v}}$ and $t = \sqrt{v}$ we obtain

$$(1 - z^2) \frac{\partial^2 \tilde{C}}{\partial z^2} - t^2 \frac{\partial^2 \tilde{C}}{\partial t^2} - 3z \frac{\partial \tilde{C}}{\partial z} - t \frac{\partial \tilde{C}}{\partial t} + 2[k_1(k_1 + 1) + k_2(k_2 - 1)] \tilde{C} = 0.$$

It is easy to see that the last equation is homogeneous in t . Thus by taking

$$\tilde{C} = t^k f(z),$$

the next ordinary differential equation results

$$(1 - z^2) \frac{d^2 f}{dz^2} - 3z \frac{df}{dz} + [(k_1 - k_2)(k_1 - k_2 + 2)]f = 0.$$

Taking $w = (1 - z)/2$ as the independent variable, the differential equation becomes

$$w(1 - w) \frac{d^2 f}{dw^2} + \frac{3}{2}(1 - 2w) \frac{df}{dw} + \rho(\rho + 2)f = 0, \quad (3)$$

with $\rho = k_1 - k_2$, a non negative integer, according to the definition of the partition κ .

Comparing with the general hypergeometric equation

$$w(1 - w) \frac{d^2 f}{dw^2} + [c - (a + b + 1)w] \frac{df}{dw} - abf = 0, \quad (4)$$

we see that the complex zonal polynomials are involved in the solution of an hypergeometric differential equation of parameters $a = -\rho$, $b = \rho + 2$ and $c = 3/2$.

Following Erdélyi *et al.* (1981), we know that a solution of (4) which is regular at $w = 0$ is given by

$$f(w) = \sum_{n=0}^{\infty} \frac{(a)_n (b)_n}{(c)_n n!} w^n = {}_2F_1(a, b; c; w),$$

where ${}_2F_1(a, b; c; w)$ is the classical hypergeometric function, which we will denote in front by $F(a, b; c; w)$.

Thus a solution of (3) is

$$f(z) = F\left(-\rho, \rho + 2; \frac{3}{2}; \frac{1 - z}{2}\right),$$

Let us refined the above solution by applying properties of the hypergeometric functions. From Erdélyi *et al.* (1981, Section 2.11, p.111), equation (2), we see that

$$F\left(2d, 2e; d + e + \frac{1}{2}; t\right) = F\left(d, e; d + e + \frac{1}{2}; 4t(1 - t)\right),$$

then

$$\begin{aligned} f(z) &= F\left(-\rho, \rho + 2; \frac{3}{2}; \frac{1 - z}{2}\right) \\ &= F\left(-\frac{\rho}{2}, \frac{\rho}{2} + 1; \frac{3}{2}; 1 - z^2\right). \end{aligned} \quad (5)$$

By Erdélyi *et al.* (1981, Section 2.10, p.108), equation (1),

$$\begin{aligned} F(a, b; c; t) &= A_1 F(a, b; a + b - c + 1; 1 - t) \\ &\quad + A_2 (1 - t)^{c-a-b} F(c - a, c - b; c - a - b + 1; 1 - t), \end{aligned}$$

where

$$A_1 = \frac{\Gamma(c)\Gamma(c-a-b)}{\Gamma(c-a)\Gamma(c-b)} \quad \text{and} \quad A_2 = \frac{\Gamma(c)\Gamma(a+b-c)}{\Gamma(a)\Gamma(b)}.$$

Then (5) can be written as follows

$$\begin{aligned} F\left(-\frac{\rho}{2}, \frac{\rho}{2} + 1; \frac{3}{2}; 1 - z^2\right) &= A_1 F\left(-\frac{\rho}{2}, \frac{\rho}{2} + 1; \frac{1}{2}; z^2\right) \\ &\quad + A_2 z F\left(\frac{3}{2} + \frac{\rho}{2}, \frac{1}{2} - \frac{\rho}{2}; \frac{3}{2}; z^2\right), \end{aligned}$$

where

$$A_1 = \frac{\Gamma(\frac{3}{2})\Gamma(\frac{1}{2})}{\Gamma(\frac{3}{2} + \frac{\rho}{2})\Gamma(\frac{1}{2} - \frac{\rho}{2})} \quad \text{and} \quad A_2 = \frac{\Gamma(\frac{3}{2})\Gamma(-\frac{1}{2})}{\Gamma(-\frac{\rho}{2})\Gamma(\frac{\rho}{2} + 1)}.$$

If $\rho = k_1 - k_2 = 2n$, $n = 0, 1, 2, \dots$, and using the fact that $\Gamma(\frac{1}{2} + z)\Gamma(\frac{1}{2} - z) = \pi \sec(z\pi)$ and $\Gamma(z)\Gamma(1 - z) = \pi \csc(z\pi)$ which implies $A_1 = \frac{(-1)^n}{2n+1}$ and $A_2 = 0$, respectively, then the even complex zonal polynomials of second order are given by

$$\tilde{C}_{(k_1, k_2)}(Y)/\tilde{C}_{(k_1, k_2)}(I_2) = (y_1 y_2)^{\frac{k}{2}} \frac{(-1)^n}{2n+1} F\left(-n, n+1; \frac{1}{2}; \frac{(y_1 + y_2)^2}{4y_1 y_2}\right).$$

If $\rho = k_1 - k_2 = 2n + 1$, $n = 0, 1, 2, \dots$ and using the same properties for the simplification of A's then we find the odd complex zonal polynomials of second order

$$\tilde{C}_{(k_1, k_2)}(Y)/\tilde{C}_{(k_1, k_2)}(I_2) = (y_1 y_2)^{\frac{k}{2}} (-1)^n \frac{y_1 + y_2}{2\sqrt{y_1 y_2}} F\left(n+2, -n; \frac{3}{2}; \frac{(y_1 + y_2)^2}{4y_1 y_2}\right),$$

where $z^2 < 1$ and

$$\tilde{C}_{(k_1, k_2)}(I_2) = k! \frac{(k_1 - k_2 + 1)^2}{(k_1 + 1)!(k_2)!},$$

see Khatri (1970).

These formulae agree with the expressions found for the complex zonal polynomials in Farrell (1980) until the 5th degree.

Acknowledgments

This research work was partially supported by CONACYT-México, research grant no. 45974-F.

References

Díaz-García, J.A.; Caro, F.J. (2006a) “An alternative approach for deriving the Laplace-Beltrami operator for the zonal polynomials of positive semidefinite and definite matrix argument”, *Far East Journal of Mathematical Sciences (FJMS)* **22**(3): 273–290.

- Díaz-García, J.A.; Caro, F.J. (2006b) “Derivation of the Laplace-Beltrami operator for the zonal polynomials of positive definite Hermitian matrix argument”. Submitted.
- Erdélyi, A.; Magnus, W.; Oberhettinger, F.; Tricomi, F.; Bertin, D.; Fulks, W.; Harvey, A.; Thomsen, D.; Weber, M.; Whitney E. (1981) *Higher Transcendental Functions 1*. Robert E. Krieger Publishing Company, Malabar.
- Farrell R.H. (1980) “Calculation of complex zonal polynomials”, in: P.R. Kishnaiah (Ed.) *Multivariate Analysis V*, North Holland: 301–320.
- James, A. T. (1968) “Calculation of zonal polynomial coefficients by use of the Laplace-Beltrami operator”, *Ann. Math. Statist.* **39**: 1711–1718.
- Khatri, C. G. (1907) “On the moments of traces of two matrices in three situations for complex multivariate normal populations”, *Sankhyā, A* **32**: 65–80.