



SHILAP Revista de Lepidopterología

ISSN: 0300-5267

avives@orange.es

Sociedad Hispano-Luso-Americana de
Lepidopterología
España

Ilyinykh, A.; Dubovskiy, I.; Polenogova, O.; Ponomarev, V.; Glupov, V.
Embryonic death as a probable reason for the collapse of population densities in
Lymantria dispar (Linnaeus, 1758) (Lepidoptera: Erebidae, Lymantriinae)
SHILAP Revista de Lepidopterología, vol. 45, núm. 179, septiembre, 2017, pp. 457-465
Sociedad Hispano-Luso-Americana de Lepidopterología
Madrid, España

Available in: <http://www.redalyc.org/articulo.oa?id=45552790011>

- How to cite
- Complete issue
- More information about this article
- Journal's homepage in redalyc.org

redalyc.org

Scientific Information System

Network of Scientific Journals from Latin America, the Caribbean, Spain and Portugal

Non-profit academic project, developed under the open access initiative

Embryonic death as a probable reason for the collapse of population densities in *Lymantria dispar* (Linnaeus, 1758) (Lepidoptera: Erebidae, Lymantriinae)

A. Ilyinykh, I. Dubovskiy, O. Polenogova, V. Ponomarev
& V. Glupov

Abstract

For the first time the causes of mass death of embryos in a population of *Lymantria dispar* (Linnaeus, 1758) with biochemical and molecular methods are investigated. In this study egg masses were collected in the territory of Russia from two populations, with low (the West Siberian population) and high (Trans-Ural) density and analyzed. Spring hatching of larvae from the eggs stored under constant laboratory conditions at 2° C was two times lower for insects of the Trans-Ural (TU) population compared to the West Siberian (WS) population ($43 \pm 6\%$ and $86 \pm 7\%$ accordingly). The level of virus defined by PCR for TU and WS populations was $76 \pm 9\%$ and $36 \pm 6\%$, respectively. The major components of eggs were the same between populations. However, we found decreased concentrations of proteins important for embryogenesis in TU population, such as an arilforin-like molecule and no vitellin 2-like component. Field data were comparable to those of the laboratory, and in TU populations there was a decrease of density. The obtained results suggest that alteration in hormonal balance of the infected with nucleopolyhedrovirus (NPV) insects may lead to a decrease or absence of some key proteins of embryogenesis in eggs. It can influence hatching of larvae from eggs and, accordingly, population dynamics of the lepidopteran.

KEY WORDS: Lepidoptera, Erebidae, Lymantriinae, *Lymantria dispar*, population dynamics, embryonic death, nucleopolyhedrovirus.

Muerte embrionaria como probable razón para el colapso de la densidad de población en *Lymantria dispar* (Linnaeus, 1758) (Lepidoptera: Erebidae, Lymantriinae)

Resumen

Se investiga por primera vez con métodos bioquímicos y moleculares, las causas de la muerte masiva de embriones en una población de *Lymantria dispar* (Linnaeus 1758). En este estudio se recogieron y analizaron masas de huevos de dos poblaciones en el territorio de Rusia: con baja (la población siberiana occidental) y alta densidad (Trans-Ural), respectivamente. La eclosión en primavera de las larvas de los huevos, guardados bajo condiciones constantes del laboratorio a 2° C, fue dos veces menor (la mitad) en las poblaciones de insectos de la Trans-Ural (TU) que en la del oeste siberiano (WS) ($43 \pm 6\%$ y $86 \pm 7\%$ respectivamente). El nivel de virus definido por PCR para las poblaciones de TU y WS era de $76 \pm 9\%$ y $36 \pm 6\%$, respectivamente. La mayoría de los componentes importantes de los huevos eran prácticamente los mismos entre las poblaciones. Sin embargo, encontramos una presencia reducida de proteínas importantes para la embriogénesis en la población de TU, como por ejemplo, una molécula parecida al pro-arilforin y una falta del componente pro-vitelín 2. Los datos de campo fueron comparables a éstos del laboratorio, y en las poblaciones de TU, había un decrecimiento de la densidad.

Los resultados obtenidos indican que la alteración en el balance hormonal de los insectos infectados con nucleopolihedrovirus (NPV) podría llevar a un disminución o falta de algunas proteínas de la embriogénesis en huevos. Puede influir en el desarrollo de las larvas desde los huevos y, en consecuencia, en la dinámica de población del lepidóptero.

PALABRAS CLAVE: Lepidoptera, Erebidae, Lymantriinae, *Lymantria dispar*, dinámica de poblaciones, muerte embrionaria, nucleopolihedrovirus.

Introduction

The gypsy moth is one of the most biologically and economically significant defoliator that periodically forms outbreaks in the territories of Eurasia, North America and North Africa (GIESE & SCHNEIDER, 1979; JOHNSON *et al.*, 2005). Population dynamics of this insect can influence both abiotic and biotic factors, causing death of insects at various stages, including an egg phase. Insect eggs represent a self-sustaining system which provides the raw materials for building the larval body and the energy reserves for embryogenesis (SANDER *et al.*, 1985). The development of the embryo is dependent upon the appropriate physiological and environmental conditions. The most important environmental condition for development of the embryo is favorable temperature and humidity (HAMILTON, 1950).

There are several studies of mass embryonic mortality of gypsy moth in natural populations (KONDAKOV, 1963; ILYINSKY & TROPIN, 1965; KOLTUNOV *et al.*, 1998). This research demonstrates which abiotic or biotic factors cause embryonic death. Moreover, there are several observations about the unknown etiology of embryonic death. EGOROV (1958) demonstrated that in 1953 in the Altay territory there were up to 15 egg masses per tree. However, in the spring of 1954 a large portion of the embryos in the eggs were dead. That has led to the collapse of population density. Research of KONDAKOV (1963) in the Krasnoyarsk region in 1954 and 1955 has revealed a mass death of eggs without clear etiology. Laboratory cultivation of gypsy moth larvae from the Trans-Ural population in 1991, where the eggs were kept under 0° C and 60-70 % humidity, hatched in May at a rate of 5-10 % (KOLTUNOV *et al.*, 1998). The same hatching rate authors observed in nature. Moreover, there is a study that in the Novosibirsk region decrease of gypsy moth density in 1997-1998 basically has been connected with mass death of embryos for unstated reasons (ILYINYKH, 2002).

It was shown that nucleopolyhedrovirus (NPV) could be one of the main factors of mortality in gypsy moth populations (ELKINTON, 1990; DWYER & ELKINTON, 1995; HOCH *et al.*, 2001). Probably, NPV infections can be a factor resulting in the mortality of embryos (straight or indirectly). However, there are cases when the death rate from NPV at an embryo stage has still not been identified in our practice and in the literature. Moreover, nutrition content of the eggs could be one of the main reasons for successful development of embryos. It is known that lipid and carbohydrate reserves decrease as embryogenesis progresses (QUICKENDEN, 1970). Yolk proteins also take part in both energy storage and embryogenesis and they are critically important for development of the embryo (IZUMI *et al.*, 1994).


Thus, in the known literature there are at least four studies about high mortality of embryos without clear etiology (EGOROV, 1958; KONDAKOV, 1963; KOLTUNOV *et al.*, 1998; ILYINYKH, 2002). Therefore, in the present work the causes of this phenomenon on two populations of gypsy moth with low (WS population) and high (TU population) egg mortality are investigated using both biochemical and molecular methods.

Material and Methods

(a) EXPERIMENTAL DESIGN

The field and laboratory experiments were conducted during an outbreak of the gypsy moth in birch forests (*Betula pendula* Roth.) of Sverdlovsk (Trans-Ural population, Kamensk district) and

Novosibirsk (West Siberian population, Karasuksky district) regions of Russia in 2011-2012 (see figure 1). The investigations were performed at two stages of the outbreak: density increase in the West Siberian population (WS) and decrease in the Trans-Ural population (TU) (data from the Novosibirsk and Sverdlovsk Centers of Forest Protection).



For detection NPV-caused mortality, the number of studied plots per population varied from five to seven, and each 100 m² plot contained three to five model trees. The number of alive and dead insects was established at the larva and pupa stages by cutting branches from model trees under study, as described previously (ILYINYKH *et al.*, 2004). To count the number of dead insects and to detect NPV in laboratory, three branches were cut per tree (one from low down, one from the middle and one from top of the crown). Cut branches fell on a parachute spread under the crown. The numbers of dead insects were counted on each branch and then this result was multiplied by the total number of branches in a tree (ILYINYKH *et al.*, 2004). The cause of insect death (45-207 individuals per population) was determined by light microscopy (see next paragraph).

In 2011, more than 100 egg masses were collected in the third decade of September in both populations in studied plots. Eggs were stored at a temperature of 2° C prior to testing of hatching, fertility, mortality, weight, and virus with PCR. In January 2012 part of the eggs was stored at or below -20° C to test protein, fat, and carbohydrate content. Eggs from both populations were tested on fertility and weight (presented as mass per 100 eggs). Hatching and number of egg masses in field conditions for both populations were provided by the Centers of Forest Protection in 2011-2012. Moreover, the above- mentioned organizations provided data about the dynamics of the total areas of major defoliation (above 60 %) of birch forests caused by the gypsy moth for both populations in 2003-2014.

(b) DETECTION OF NUCLEOPOLYHEDROVIRUS

For detection of mortality from entomopathogens, 200 larvae from both populations were cultivated in May until reaching imago under conditions preventing exogenous virus on the artificial medium (AM) using the technique described by ILYINYKH (1997). Dead insects were examined on infection with light microscope (Biolam - R15; LOMO, Russia) using phase contrast.

Twenty cleared eggs from each of the studied egg masses were selected for PCR detection of NPV. Eggs were sterilized within 10 min in 0.25 % NaOH solution on a magnetic mixer, washed with sterile water and dried. Embryos were collected from eggs under sterile conditions (20 individuals per sample). Samples were stored at - 70° C until DNA was extracted.

Total DNA from samples of insects was extracted using the DNA Extraction kit ("MEDIGEN Laboratory", Russia) according to the protocol of the manufacturer. Detection of viral conjugation protein gene Ld130 was carried out in 20 l of buffer contained 10 l of PyroStart™ Fast PCR Master Mix (2X) ("Fermentas", USA); 0.1 l of forward (5' CGGGCATCATCCGCGGCC 3' (127651 - 127668)) and reverse (5' CGCCCTCCAGCTCCGCGC 3' (127944 - 127927)) primers and 27.5 % of DNA on volume. Specific primers were designed using the full-genomic sequence (GenBank database at number NC_001973) of virus of gypsy moth. PCR carried out on a thermocycler "DNA Engine Dyad® Peltier Thelmar Cycler" ("BIO-RAD", USA) using the following protocol (denaturation 30s with 94° C, annealing 30s with 68° C, synthesis 30 with 72° C (37 cycles)); synthesis of 7 min. - 72° C. The size of the detected gene fragment was 294 bp.

(c) ANALYSIS

For protein, fatty acid and carbohydrate detection, 100 mg per sample of eggs from both populations were collected and homogenized using the FastPrep MP biomedical (ICN) homogenizer and freeze dried. For testing of carbohydrate and protein concentrations frozen-dried samples were dissolved in distilled water. For determination of fatty acid concentration, the freeze-dried samples were dissolved in concentrated sulfuric acid.

Detection of fatty acid, carbohydrate and protein concentrations were carried out in triplicate using a spectrophotometer. Fat content was checked according to CHABROL & CHARONNAT (1937) with some modifications. The freeze-dried sample was incubated with 1 ml of sulfuric acid for 20 min at 100° C. The mix was cooled for 5 min and added to 500 l of vanillin (Sigma) and then dissolved in distilled water (13 mM). Optical density of the solution registered at 530 nm in 30 min. Concentration of fatty acids was determined with a calibration curve, using vegetable fatty acid (olive oil, 99 %) standards.

Detection of carbohydrates was carried out according to HANSEN & MOLLER (1975) with some modifications. We mixed 100 l of sample with 500 l of 0.5 % Antron (Sigma) solution in 72 % sulfuric acid and incubated for 11 min at 100° C. Then the mixture was quickly cooled to 0° C on ice. Optical density was detected at 630 nm in 60 min at 22° C. Concentration of carbohydrates were determined with a calibration curve, using starch (99 %, Sigma) standards. The protein concentration of samples was estimated by using the Bradford method (BRADFORD, 1976), using bovine serum albumin standards. Qualitative composition of proteins was detected with capillary electrophoresis (protein kit) (Agilent Bioanalyzer).


Data are presented as mean \pm the standard error. To check the data for normal distribution, the Wilk Shapiro W criterion was used. All results were assessed using one-way ANOVA, followed by Tukey's post-hoc tests to identify specific differences between means.

Results**(a) WEIGHT, FERTILITY AND MORTALITY OF EGGS**

We found that weight and fertility of eggs from both the WS (population increase) and TU

(population decline) populations did not differ (table 1). Moreover, the hatching of larvae in January was the same in both populations. However, in May (post-diapause) the hatching of larvae from eggs in the WS population was higher compared with the TU population ($86 \pm 7\%$ and $43 \pm 6\%$ respectively, $p < 0.01$) (table 1).

The same data was found in natural conditions during testing in the forest. Spring hatching in the TU population was about 30 % while in the WS population the hatching was about 85 %. The majority of larvae (approximately 90 %) that hatched from eggs in the TU population died of unclear reason at first and second instar. The study from the forest showed that the area around the TU population had low egg mass density: 21 ± 10 egg masses/ha (previous year 3650 ± 457 egg masses/ha). While the density of the egg masses in the WS population territory increased up 1240 ± 265 egg masses/ha (previous year 64 ± 23 egg masses/ha). Moreover, the total area of major defoliation (above 60 %) in the region increased from 28 hectares to 2500 while in the TU population this index decreased from 39945 hectares to 3. The data for both populations in 2003-2014 are shown on figure 2. These data demonstrate that the TU gypsy moth population (Sverdlovsk region) decreased, whilst the WS gypsy moth population (Novosibirsk region) increased.





(b) NUTRITION CONTENT OF EGGS

The concentrations of proteins, lipids and carbohydrates in eggs were similar between both the TU and WS populations (figure 3). Concentration of lipids in eggs was on average 104 ± 13 mg/g, proteins averaged 16 ± 0.65 mg/g, and 1.93 ± 0.14 mg/g was the average concentration of carbohydrates. However, a qualitative protein assay showed that in eggs from the TU population, concentration of an arylphorin-like molecule (~ 75 kDa) was 1.5 times lower ($p < 0.001$) than in eggs from the WS population. Additionally, in eggs from the TU population, the vitellin 2-like component (~ 45 kDa) was absent.

(c) INFECTIONS OF INSECTS

We did not find any infections, including NPV, in the embryos and hatched larvae with the light microscope. However, at cultivation on the artificial media, mortality of the TU population larvae from

NPV was $5 \pm 1\%$ while larvae from the WS population did not die from virus ($p < 0.01$). The mortality of the larvae due to unstated reasons for the TU and WS populations was $92 \pm 8\%$ and $12 \pm 3\%$ ($p < 0.01$) respectively. The highest mortality ($82 \pm 7\%$) for the TU population was observed in first instar larvae.



In nature, the NPV carrying level (detected by PCR) among insects (embryos) from the TU population in 2011 was $76 \pm 9 \%$, but mortality at the larva and pupa stages from NPV was $12 \pm 5 \%$. The larvae from the WS population did not die from NPV in 2011; however, virus carrying quantity among embryos was $36 \pm 6 \%$ in the autumn. PCR detection of viral gene (LD 130) in embryos from the TU population showed 6 positive cases from 10 samples (figure 4). Data on the WS population aren't provided.

Discussion

The TU population of gypsy moth was found to have increased mortality of embryos. This is probably the main factor of decline of population density in the area of the TU population. Several possible biotic reasons for embryonic death were analyzed in the study. Nutrition in eggs, an important factor of embryonic feeding, was similar for both the TU and WS populations. However, some key proteins of embryogenesis (arylphorin-like molecule and the vitellin 2-like component) were decreased or absent in eggs from the TU population. It is possible that the lowered survival of eggs from the TU population can be connected with the shortage of these substances.

It was demonstrated that NPV could be one of the main factors of mortality in population dynamics of the gypsy moth (DWYER & ELKINTON, 1995; HOCH *et al.*, 2001; FULLER *et al.*, 2012). At the same time, it is shown that the impact of this factor in various parts of gypsy moth areas can be different. In particular it is well known that in populations in the territory of Trans Ural and Western Siberia, NPV was detected locally but at an insignificant level (about 10%) of larvae (ILYINYKH *et al.*, 2004).

Baculoviruses infect over 600 species of insects (ROHRMANN, 2008), and in some cases, they were successfully used to control different insect pests (reviewed by INCEOGLU *et al.*, 2006). Although horizontal route is thought to be the major pathway for baculovirus transmission (CORY & MYERS, 2003), some studies also reveal vertical transfer in field populations (reviewed by KUKAN, 1999; ZHOU *et al.*, 2005; KOUASSI *et al.*, 2009). Moreover, the vertical transmission of gypsy moth NPV was described earlier in our own investigations (ILYINYKH *et al.*, 2004; ILYINYKH & POLENOGOVA, 2013). Individuals exposed to low doses of virus may acquire a non-fatal sublethal infection, but transmit the virus vertically to the next generation of insects (BURDEN *et al.*, 2002; CABODEVILLA *et al.*, 2011; MURILLLO *et al.*, 2011). This may affect insect health, weight and fecundity (MYERS *et al.*, 2000; VILAPLANA *et al.*, 2008). Probably, NPV could be one of the reasons for embryo mortality in the TU population. The NPV is capable of affecting hormonal balance of the infected insects. In some baculoviruses the *egt*-gene was found, which is capable of coding the UDP-glucosyltransferase catalyzing binding of sugars by ecdysteroids (O'REILLY & MILLER, 1989; SLAVICEK *et al.*, 1999). In particular infection of gypsy moth by NPV containing the *egt*-gene, led to abnormality of molting and growth of insects. Virus with a deletion of the *egt*-gene did not change the growth of insects (SLAVICEK *et al.*, 1999).

Probably, alteration in hormonal balance of the infected insects (O'REILLY & MILLER, 1989; SLAVICEK *et al.*, 1999) may lead to a decrease or absence of some key proteins of embryogenesis (arylphorin-like molecule and the vitellin 2-like component) in eggs from the TU population.

Perhaps, insects infected with NPV can demonstrate delay of development and lay eggs later in comparison with non-infected individuals or insects with smaller quantity of virus carriers. It can lead to a decrease in the sum of the effective temperatures necessary for normal embryogenesis of insects (especially in the conditions of a continental climate) and/or can change embryonic diapause. It can influence hatching of larvae from eggs and, accordingly, population dynamics of the gypsy moth.

Possibly, in further research, the biochemical methods can be employed to diagnostics of a phase of depression in population dynamics of the gypsy moth.

Acknowledgements

The authors would like to thank Colette Pietzsch for reading the manuscript. This research was supported by the Federal Fundamental Scientific Research Programme for 2013-2020 (VI.51.1.5) and in part the Russian Foundation for Basic Research, project no. 14-04-00615a.

BIBLIOGRAPHY

- BRADFORD, M., 1976.– A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding.– *Analytical Biochemistry*, **72**: 248-254.
- BURDEN, J. P., GRIFFITS, C. M., CORY, J. S., SMITH, P. & SAIT, S. M., 2002.– Vertical transmission of sublethal granulovirus infection in the Indian meal moth, *Plodia interpunctella*.– *Molecular Ecology*, **11**: 547-555.
- CABODEVILLA, O., VILLAR, E., VITRO, C., MURILLO, R., WILLIAMS, T. & CABALLERO, P., 2011.– Intra- and intergenerational persistence of an insect nucleopolyhedrovirus: adverse effects of sublethal disease on host development, reproduction, and susceptibility to superinfection.– *Applied and Environmental Microbiology*, **77**: 2954-2960.
- CHABROL, E. & CHARONNAT, R., 1937.– Une nouvelle reaction pour l'étude des lipids. L'eleidemie.– *La presse médicale*, **45**: 1713-1714.
- CORY, J. S. & MYERS, J. H., 2003.– The ecology and evolution of insect baculoviruses.– *Annual Review of Ecology Evolution and Systematics*, **34**: 239-272.
- DWYER, G. & ELKINTON, J. S., 1995.– Host dispersal and the spatial spread of insect pathogens.– *Ecology*, **76**: 1262-1275.
- EGOROV, N. N., 1958.– Pest insects of pine forests of West Siberia.– *Zoologicheskii zhurnal*, **37**: 1488-1499 [in Russian with English summary].
- ELKINTON, J. S., 1990.– Population dynamics of gypsy moth in North America.– *Annual Review of Entomology*, **35**: 517-596.
- FULLER, E., ELDERD, B. D. & DWYER, G., 2012.– Pathogen Persistence in the Environment and Insect-Baculovirus Interactions: Disease-Density Thresholds, Epidemic Burnout and Insect Outbreaks.– *American Naturalist*, **179**: 70-96.
- GIESE, R. L. & SCHNEIDER, M. L., 1979.– Cartographic comparisons of Eurasian gypsy moth distribution (*Lymantria dispar* L., Lepidoptera: Lymantriidae).– *Entomological News*, **90**: 1-16.
- HAMILTON, A. G., 1950.– Further studies on the relation of humidity and temperature to the development of two species of African locusts - *Locusta migratoria migratorioides* (R. & F.) and *Schistocerca gregaria* (Forsk.).– *Transactions of the Royal Entomological Society of London*, **101**: 1-58.
- HANSEN, J. & MØLLER, I., 1975.– Percolation of starch and soluble carbohydrates from plant tissue for quantitative determination with anthrone.– *Analytical Biochemistry*, **68**: 87-94.
- HOCH, G., ZUBRIC, M., NOVOTNY, J. & SCHOPF, A., 2001.– The natural enemy complex of the gypsy moth, *Lymantria dispar* (Lepidoptera, Lymantriidae) in different phases of its population dynamics in eastern Austria and Slovakia, a comparative study.– *Journal of Applied Entomology*, **125**: 217-227.
- ILYINSKY, A. I. & TROPIN, I. V. (eds.), 1965.– *Supervision, registration and prognosis of outbreaks of needle- and leaf-eating insects in forests of the USSR*. Moscow: Lesnaya Promishlennost' [in Russian].
- ILYINYKH, A. V., 2002.– Analysis of causes of waning of mass reproduction of gypsy moth (*Lymantria dispar* L.) in the territory of the Novosibirsk region.– *Contemporary problems of ecology*, **6**: 697-702.
- ILYINYKH, A. V., 1997.– The method for the laboratory cultivation of *Ocneria dispar* L.– *Biotechnologiya*, **9-10**: 27-29. [in Russian with English summary].
- ILYINYKH, A. V. & POLENOGOVA, O. V., 2013.– Demonstration of remote effect for vertical transmission of baculovirus based on example of gypsy moth, *Lymantria dispar* L. (Lepidoptera, Lymantriidae).– *Biology Bulletin Reviews*, **3**: 214-218.
- ILYINYKH, A. V., SHTERNISH, M. V. & KUZMINOV, S. V., 2004.– Exploration into a mechanism of transgenerational transmission of nucleopolyhedrovirus in *Lymantria dispar* L. in Western Siberia.– *BioControl*, **49**: 441-454.
- INCEOGLU, A. B., KAMITA, S. G. & HAMMOCK, B. D., 2006.– Genetically modified baculoviruses: a historical overview and future outlook.– *Advances in Virus Research*, **68**: 323-360.

- IZUMI, S., YANO, K., YAMAMOTO, Y. & TAKAHSHI, S. Y., 1994.– Yolk proteins from insect eggs: Structure, biosynthesis and programmed degradation during embryogenesis.– *Journal of Insect Physiology*, 40: 735-746.
- JOHNSON, D. M., LIEBHOLD, A. M., BJØRNSTAD, O. N. & MCMANUS, M. L., 2005.– Circumpolar variation in periodicity and synchrony among gypsy moth populations.– *Journal of Animal Ecology*, 74: 882-892.
- KOLTUNOV, E. V., PONOMAREV, V. I. & FEDORENKO, S. I., 1998.– *The gypsy moth Lymantria dispar ecology in conditions of anthropogenic influences*. Ekaterinburg: UB RAS [in Russian with English summary].
- KONDAKOV, Yu. P., 1963.– The gypsy moth (*Ocneria dispar* L.) in Krasnoyarsk region forests.– In *Protection of Siberian forests from pest insects*: 30-77. Lesnaya Promishlennost', Moscow. [in Russian].
- KOUASSI, L. N'G., TSUDO, K., GOTO, C., MUKARAVA, S., SAKAMAKI, S., KUSIGEMATI, K. & NAKAMURA, M., 2009.– Prevalence of latent virus in *Spodoptera litura* (Fabricius) (Lepidoptera: Noctuidae) and its activation by heterologous virus.– *Applied Entomology and Zoology*, 44: 95-102.
- KUKAN, B., 1999.– Vertical transmission of nucleopolyhedrovirus in insects.– *Journal of Invertebrate Pathology*, 74: 103-111.
- MURILLO, R., HUSSEY, M. S. & POSSEE, R. D., 2011.– Evidence for covert baculovirus infections in a *Spodoptera exigua* laboratory culture.– *Journal of General Virology*, 92: 1061-1070.
- MYERS, J., MALAKAR, H. R. & CORY, J. S., 2000.– Sublethal nucleopolyhedrovirus infection effects on female pupal weight, egg mass size, and vertical transmission in gypsy moth (Lepidoptera: Lymantriidae).– *Environmental Entomology*, 29: 1268-1272.
- O'REILLY, D. R. & MILLER, L. K., 1989. - A baculovirus blocks insect molting by producing ecdysteroid UDP-glucosyl transferase.– *Science*, 245: 1110-1112.
- QUICKENDEN, K. L., 1970.– Carbohydrates in eggs of the grasshopper, *Aulocara elliotti*, during development.– *Journal of Insect Physiology*, 16: 171-183.
- ROHRMANN, G. R., 2008.– *Baculovirus molecular biology*. Available from <http://www.ncbi.nlm.nih.gov/books/NBK1736/>.
- SANDER, K., GUTZEIT, H. O. & JACKIE, H., 1985.– Insect Embryogenesis: Morphology, Physiology, Genetical and Molecular Aspects.– *Comprehensive insect physiology, biochemistry and pharmacology*, 1: 319-385.
- SLAVICEK, J. M., POPHAM, H. J. R. & RIEGEL, C. I., 1999.– Deletion of the *Lymantria dispar* multicapsid nucleopolyhedrovirus ecdysteroid UDP-glucosyltransferase gene enhances viral killing speed in the last instar of the gypsy moth.– *Biological Control*, 16: 91-103.
- VILAPLANA, L., REDMAN, E. M., WILSON, K. & CORY, J. S., 2008.– Density- related variation in vertical transmission of a virus in the African armyworm.– *Oecologia*, 155: 237-246.
- ZHOU, M., SUN, X., SUN, X., VLAK, J. M., HU, Z. & VAN DER WERF, W., 2005.– Horizontal and vertical transmission of wild-type and recombinant *Helicoverpa armigera* single-nucleocapsid nucleopolyhedrovirus.– *Journal of Invertebrate Pathology*, 89: 165-175.

*A. I.; I. D.; O. P.; V. G

Institute of Systematics and Ecology of Animals
Siberian Branch of Russian Academy of Sciences
Frunze, 11
630091 Novosibirsk
RUSSIA / RUSSIA
E-mail: avilyinykh@mail.ru

V. P.

Institute Botanic Garden
Ural Branch of Russian Academy of Sciences
8-Marta, 202a
620144, Ekaterinburg
RUSSIA / RUSSIA
E-mail: v_i_ponomarev@mail.ru

*Autor para la correspondencia / *Corresponding author*

(Recibido para publicación / *Received for publication* 11-XI-2016)

(Revisado y aceptado / *Revised and accepted* 20-II-2017)

(Publicado / *Published* 30-IX-2017)