



Brazilian Journal of Physics

ISSN: 0103-9733

luizno.bjp@gmail.com

Sociedade Brasileira de Física  
Brasil

DuBois, R. D.; de Lucio, O. G.; Gavin, J.  
Differential Ionization Studies for Positron Impact  
Brazilian Journal of Physics, vol. 36, núm. 2B, june, 2006, pp. 522-523  
Sociedade Brasileira de Física  
São Paulo, Brasil

Available in: <http://www.redalyc.org/articulo.oa?id=46403607>

- How to cite
- Complete issue
- More information about this article
- Journal's homepage in redalyc.org

redalyc.org

Scientific Information System  
Network of Scientific Journals from Latin America, the Caribbean, Spain and Portugal  
Non-profit academic project, developed under the open access initiative

## Differential Ionization Studies for Positron Impact\*

R. D. DuBois, O. G. de Lucio, and J. Gavin  
University of Missouri-Rolla, Rolla, MO 65409 USA

Received on 29 July, 2005

Recent studies of differential ionization by positron impact being performed at the University of Missouri-Rolla are described. Recoil ion-scattered positron and recoil ion-scattered positron-ejected electron coincidences are measured to provide doubly and triply differential ionization data (DDCS and TDCS) for  $e^+$  - Ar collisions. The DDCS and TDCS studies are for 750 and 200 eV impact respectively. For the TDCS studies the positron scattering angles are between approximately  $\pm 5^\circ$  and electrons ejected with energies less than 15 eV and angles between  $90^\circ \pm 50^\circ$  with respect to the beam direction are detected. Examples of TDCS data for the scattered positron and the ejected electron are presented.

Keywords: Positrons; Ionizations; Differential

### I. INTRODUCTION

Data for ionizing interactions between particles and antiparticles can be compared to provide information about known atomic processes or about unexplored regimes of physics. For example, charge and mass effects on the interaction dynamics and the influence of electron exchange can be studied by comparing results obtained for electron and positron impact, or positron impact ionization can be used to study ultrainelastic collisions where the projectile transfers all of its energy to the target. For these purposes, highly differential experimental data such as that provided by  $e,2e$  type experiments are needed since they provide extremely sensitive tests of theoretical models.

Numerous theoretical predictions are available for positron impact, e.g., either quantal calculations such as those provided by Sharma and Srivastava [1] and Brauner, Briggs and Klar [2]. For antiproton impact, the classical trajectory Monte Carlo calculations of Wood et al. [3] are available. But, on the experimental side only a limited number of differential studies exist for positron impact. This is because the beam intensities that can be obtained are extremely weak as compared to those for electron beams. In spite of this, a few ground breaking studies have been performed. For example, doubly differential electron emission resulting from positron impact has been measured by Schmitt et al. [4] and Kövér et al. [5], [6]. In the late 90's, more detailed information became available when Kövér and Laricchia measured the first triply differential cross sections for positron impact [7], [8]. They studied zero degree electron emission. To date, no experimental differential data exists for antiproton impact.

### II. PRESENT WORK

This paper describes work in progress at the University of Missouri-Rolla (UMR) where several years ago we started an experimental program to measure differential ionization cross sections for positron and electron impact. The apparatus consists of a  $^{22}\text{Na}$  source and moderator for producing positron and electron beams, an electrostatic beam transport system,

and a scattering chamber containing a gas jet target and a projectile energy analyzer. The electrostatic energy analyzer is equipped with a position sensitive detector which samples a range of energies and angles for forward scattered projectiles. In our first studies [9]-[11], differential energy loss for single and multiple ionization of Ar and Kr was measured for 750 eV positron impact. This was done by measuring coincidences between extracted target ions and forward scattered projectiles. These data represent the only differential information for multiple target ionization by antiparticle impact. For both targets it was found that the percentage of double ionization systematically increased for energy losses of 0 to  $\sim 50$  eV above threshold and that the degree of double ionization was smaller for positron impact than for electron impact. For larger energy losses, the percentage of double ionization remained rather constant and was similar in magnitude for both positron and electron impact.

Recently we have added another position sensitive detector that views electrons emitted between approximately  $40^\circ$  and  $140^\circ$  along and transverse to the beam direction. The x,y positions recorded for each detected electron provide information about its  $\theta, \phi$  emission angles while for single ionization the electron energy can be determined from knowledge of the initial and final beam energies and the ionization potential. Since we also know the scattered projectile energy and angles, recoil ion-scattered projectile-ionized electron coincidences provide triply differential cross section information, i.e., so-called  $(e,2e)$  measurements. Double ionization information is also obtained but in this case only the sum of the energies of the two ejected electrons is known. This corresponds to  $(e,3e-1)$  type measurements.

An example of our preliminary TDCS data is shown in the Figure. The initial positron energy is 200 eV while the final energy and scattering angle is approximately  $183 \pm 2$  eV and  $3 \pm 2^\circ$ . As seen, the limited statistics acquired to date indicate approximately equal magnitudes for the binary and recoil peaks. Slightly higher electron energies (not shown) indicate an increasing binary/recoil peak ratio and possible structure in the peaks. Additional data are currently being collected in order to clarify the peak magnitudes and shapes and to improve the statistics. With regard to the data shown, the reader should keep in mind that the influence of the recoil ion extraction field




FIG. 1: Triply differential electron emission data for single ionization of argon by 200 eV positrons. The positron scattering angle is approximately  $-1$  to  $-5^\circ$  and the electron energies are between 0 and 4 eV. The error bars are statistical uncertainties.

on these low-energy electrons still needs to be deconvoluted. Also, the calibration of the projectile energies and angles are still tentative. However, these data demonstrate that TDCS data for positron impact, such as that which has been avail-

able for electron impact since the 1960's, are finally feasible.

\*This work supported by the National Science Foundation.

- 
- [1] S. Sharma and M.K. Srivastava, *Phys. Rev. A* **38**, 1083 (1988).
  - [2] M. Brauner, J.S. Briggs, and H. Klar, *J. Phys. B* **22**, 2265 (1989).
  - [3] C.J. Wood, C. R. Feeler, and R.E. Olson, *Phys. Rev. A* **56**, 3701 (1977).
  - [4] A. Schmitt, H. Cerny, H. Möller, R. Raith, and M. Weber, *Phys. Rev. A* **49**, R5 (1994).
  - [5] Á. Kövér, G. Laricchia, and M. Charlton, *J. Phys. B* **27**, 2409 (1994).
  - [6] Á. Kövér, R.M. Finch, M. Charlton, and G. Laricchia, *J. Phys. B* **30**, L507 (1997).
  - [7] Á. Kövér, K. Paludan, and G. Laricchia, *J. Phys. B* **34**, L219 (2001).
  - [8] Á. Kövér and G. Laricchia, *Phys. Rev. Lett.* **80**, 5309 (1998).
  - [9] R.D. DuBois, C. Doudna, C. Lloyd, M. Kahveci, Kh. Khayyat, Y. Zhou, and D.H. Madison, *J. Phys. B* **34**, L783 (2001).
  - [10] A.C.F. Santos, A. Hasan, T. Yates, and R.D. DuBois, *Phys. Rev. A* **67**, 052708 (2003).
  - [11] A.C.F. Santos, A. Hasan, and R.D. DuBois, *Phys. Rev. A* **69**, 032706 (2004).