



Brazilian Journal of Physics

ISSN: 0103-9733

luizno.bjp@gmail.com

Sociedade Brasileira de Física

Brasil

Jiménez-Cruz, Camilo Andrés; Rodríguez López, J.-Alexis; Martinez, Roberto

Light Higgs Boson Production in Two Higgs Doublets Models Type III

Brazilian Journal of Physics, vol. 38, núm. 3B, septiembre, 2008, pp. 455-458

Sociedade Brasileira de Física

São Paulo, Brasil

Available in: <http://www.redalyc.org/articulo.oa?id=46415795013>

- How to cite
- Complete issue
- More information about this article
- Journal's homepage in redalyc.org

redalyc.org

Scientific Information System

Network of Scientific Journals from Latin America, the Caribbean, Spain and Portugal

Non-profit academic project, developed under the open access initiative

Light Higgs Boson Production in Two Higgs Doublets Models Type III

Camilo Andrés Jiménez-Cruz,^{*} J.-Alexis Rodríguez López,[†] and Roberto Martínez[‡]

Departamento de Física - Universidad Nacional de Colombia - Sede Bogotá, Colombia

(Received on 16 April, 2008)

By using the Cheng, Sher and Yuan's ansatzs, we study the light Higgs Boson production associated with b quark production at TEVATRON using the 2HDM type III. We compare the simulations with experimental results coming from TEVATRON, finding valid ranges for the bb coupling. By using these results, we calculate the cross section for the process $pp \rightarrow b\bar{b}h(b\bar{b})$ for the LHC collider.

Keywords: Light Higgs Boson; b Production; LHC

I. INTRODUCTION

Although a relatively light Higgs boson coming from the SM can support the idea of the SM being true all the way to the Plank scale, most theorists consider such a possibility unlikely [1]. A light Higgs boson is preferred by precision fits of the Standard Model (SM) [2] and also theoretically required by many frameworks that can be effectively SM-like at low energies. The production of a Higgs boson in association with a heavy quark and antiquark pair (both $t\bar{t}$ and $b\bar{b}$) at hadron colliders will be sensitive to the Higgs-fermion couplings and can help discriminate between models. One of such models is the so-called 2HDM, which includes a second scalar doublet with the same properties of the first one.

In this paper, by using the framework of the 2HDM-III and the CSY ansatzs (section II), we calculate the associated b-quark/Higgs boson production at the TEVATRON collider. By comparing the results obtained in the simulation with experimental data released by the D0 collaboration we find bounds over the λ_{bb} parameter of the coupling (section III). The bounds found let us estimate the associated higgs/bottom production in the Large Hadron Collider (section IV). Finally, in section V we state our concluding remarks.

II. TWO HIGGS DOUBLET MODEL (2HDM)

The 2HDM includes a second scalar doublet, with each doublet acquiring a vacuum expectation value different from zero.

The scalar content of the model becomes:

$$\Phi_i = \begin{pmatrix} \phi_i^+ \\ \phi_i^0 \end{pmatrix}, \quad \langle \Phi_i \rangle = \begin{pmatrix} 0 \\ \frac{v_i}{\sqrt{2}} \end{pmatrix}, \quad i = 1, 2 \quad (1)$$

In this way, the scalar spectrum of mass eigenvalues contains two CP-even neutral Higgs bosons (h_0, H_0) coming from the mixing of the real part of the neutral components of both doublets with a mixing angle α ; two charged Higgs bosons (H^\pm), which mix with the would-be Goldstone bosons (G_W^\pm) through

the mixing angle $\tan\beta = v_2/v_1$ and one CP-odd Higgs (A_0), which mixes with the neutral would-be Goldstone.

A. The Yukawa Lagrangian

The most general Lagrangian that can be written in this kind of models includes interactions between all the fermions and both doublets:

$$\begin{aligned} -\mathcal{L}_Y = & \eta_{ij}^{U,0} \bar{Q}_{iL}^0 \tilde{\Phi}_1 U_{jR}^0 + \eta_{ij}^{D,0} \bar{Q}_{iL}^0 \tilde{\Phi}_1 D_{jR}^0 \\ & + \xi_{ij}^{U,0} \bar{Q}_{iL}^0 \tilde{\Phi}_2 U_{jR}^0 + \xi_{ij}^{D,0} \bar{Q}_{iL}^0 \tilde{\Phi}_2 D_{jR}^0 \\ & + \text{l.s.} + \text{h.c} \end{aligned} \quad (2)$$

The strong suppression of FCNC at tree level makes customary to impose discrete symmetries over the doublets. Those symmetries end in one of three types of models [3].

• Type I

Φ_1 is the responsible of giving mass to the matter, while Φ_2 decouples totally from them.

• Type II

Φ_1 couples to the up sector while Φ_2 couples to the down sector. This is the case of the MSSM.

• Type III

Both doublets couple to both sectors.

B. 2HDM type III

As there are two non-diagonal 3×3 matrices in the Yukawa lagrangian, and the suffix 0 means that these fermion states are not mass eigenstates. It is clear that the mass terms for the


Parameter	Range
$\xi_{\mu\tau}^2$	$[7.62 \times 10^{-4} : 4.44 \times 10^{-2}]$
$\xi_{\tau\tau}$	$[-1.8 \times 10^{-2} : 2.2 \times 10^{-2}]$
$\xi_{\mu\mu}$	$[-0.12 : 0.12]$
$\xi_{\mu e}$	$[-0.39 : 0.39]$
λ_{bb}	$[-100 : 100]$
λ_{tt}	$[-\sqrt{8} : \sqrt{8}]$

TABLE I: Experimental constraints over the ξ and λ matrices

^{*}Electronic address: cajimenezc@unal.edu.co

[†]Electronic address: jarodriguezl@unal.edu.co

[‡]Electronic address: remartinez@unal.edu.co

FIG. 1: Processes involved in the associated ϕ/b production.

matter will depend on two Yukawa coupling matrices. The rotation of the quarks and leptons allow us to diagonalize one of the matrices but in general not both simultaneously, then one Yukawa coupling remains non-diagonal, leading to the FCNC at tree level.

C. Cheng, Sher and Yuan's Ansatz (CSY)

In this work, we use the CSY parameterization of the Yukawa's couplings [4]. This ansatz is based on the SM $\phi f \bar{f}$ couplings and states that

$$\xi^{ij} \equiv \frac{\sqrt{m_i m_j}}{v} \lambda_{ij}$$

This is an ansatz for the Yukawa texture matrices looking for a phenomenological similarity with SM couplings. Here, v has been defined as $v \equiv \sqrt{v_1^2 + v_2^2}$.


Some restrictions over the λ_{ij} and the ξ_{ij} parameter sets have been found [5] and can be seen in Table I. The λ_{bb} parameter can be constrained by the coupling of the b quark to the scalar sector.

III. ASSOCIATED $h_0 b \bar{b}$ PRODUCTION

In the SM, the scalar sector couples to the fermions with strength proportional to the mass of the fermion. In this way, the Yukawa coupling to bottom quarks ($m_b \sim 5\text{GeV}$) leads to small cross sections in hadron colliders. However, processes involving b quarks and light Higgs bosons (Fig. 1) can be strongly enhanced in some 2HDM scenarios. This associated production has been extensively studied for 2HDM type II [6-9], being the Minimal Supersymmetric Standard Model a favorite framework for calculation. However, 2HDM type III models' cross sections can be enhanced in a similar way.

The predominant modes are

$$qq, gg \rightarrow b \bar{b} h_0 (b \bar{b}), \text{ and } gb \rightarrow bh(b \bar{b}) \quad (3)$$

FIG. 2: Light Higgs boson production associated with two high p_T b -quarks and decaying into a $b \bar{b}$ pair, for different values of λ_{bb} and M_h , $\alpha = \sqrt{2}/2$.

so the interesting events would have at least four jets in the final state where two of these jets should come from a Higgs resonance. The signal-to-background ratio can be enhanced by requiring that the four jets (or at least three) of the highest transverse energies are tagged as b -jets. This mode could be of great interest when the mass of the scalar boson is greater than 150 MeV, where the sensitivity to the $h \rightarrow \tau \tau$ channel disappears [10].

Specifically, in the frame of the 2HDM-III and by using the CSY ansatz, the coupling between the lightest Higgs boson and the b quarks is given by

$$A(h_0 b \bar{b}) = h_0 \bar{b} \left(\frac{\cos \alpha + \sin \alpha \tan \beta}{\sqrt{2}} \xi^D - \frac{\sin \alpha}{v} m_b \right) b \quad (4)$$

where α and β are the angles defined before.

In terms of the *fundamental parameterization* [3], the coupling becomes

$$A(h_0 b \bar{b}) = h_0 \bar{b} \left(-\frac{\sin \alpha}{v} m_b + \frac{\cos \alpha}{\sqrt{2}} \frac{m_b}{v} \lambda_{bb} \right) b. \quad (5)$$

where we see explicitly the high dependence on the λ_{bb} parameter defined in the CSY ansatz.

IV. RESULTS

By using the CalcHEP package [12] as the Monte Carlo event generator, we calculated the associated bottom-quark/Higgs production in the TEVATRON collider for different values of λ_{bb} and m_ϕ . The results can be seen in Figure 2. We used the CTEQ6M Parton Distribution Function for the proton and the antiproton, a momentum of 1980 GeV at center of mass and a representative value of $\alpha = \pi/4$.

As in this parameter region the h_0 boson is nearly degenerate in mass with either the h_0 or the A_0 bosons, they cannot be




FIG. 3: Plot matching experimental data [11] coming from the D0 experiment (TEVATRON) and simulated data running m_h and λ_{bb} .




FIG. 4: Estimated cross section for the associated bottom quark production for the LHC collider ($pp \rightarrow b\bar{b}h(b\bar{b})$), here we used $\lambda_{bb} = 6$.

distinguished experimentally [11] and the total cross section for the signal is assumed to be twice of the light boson [13].

The total cross section for the process is then [13]:

$$\sigma(p\bar{p} \rightarrow b\bar{b}h_0 \rightarrow b\bar{b}b\bar{b}) \sim 2 \times \sigma(p\bar{p} \rightarrow b\bar{b}\phi) \times BR(h_0 \rightarrow b\bar{b}). \quad (6)$$

The results were matched against experimental results coming from the D0 experiment [11] (Fig. 3), showing a great agreement for small values of λ_{bb} . In this way, we got the following bounds over the λ_{bb} parameter, given $\alpha \sim \frac{\sqrt{2}}{2}$:

$$-6 \leq \lambda_{bb} \leq 6 \quad (7)$$

By using the bounds obtained, we can estimate the associated b-quark-higgs boson production rates for the LHC collider. The results are depicted in the Fig. 4.

V. CONCLUDING REMARKS

The coupling between the light neutral Higgs boson and the fermions is quite sensitive to the existence of a non-minimal scalar sector.

By using the Cheng, Sher and Yuang parameterization of the coupling between the b quark and the scalar sector we calculated the total cross section for the process $p\bar{p} \rightarrow b\bar{b}h(b\bar{b})$ at a center of mass energy of 1.98TeV. We compared this result with the reported data coming from the D0 experiment at TEVATRON [14]. In this way, we obtained that the λ_{bb} parameter of the anzats should be in the range $[-6, 6]$ improving the actual bound provided by unitarity and included in the table I.

With this results, we calculated the associated b quark production of the light Higgs boson for the LHC.

Acknowledgments

R.M. acknowledges the financial support of the Banco de la Republica de Colombia.

- [1] M. Carena, J. S. Conway, H. E. Haber, and J. D. Hobbs, *Report of the Higgs Working Group of the Tevatron Run 2 SUSY/Higgs Workshop*, (2000), URL <http://www.citebase.org/abstract?id=oai:arXiv.org:hep-ph/0010338>.
- [2] T. L. C. A. Collaboration, D. Collaboration, L. Collaboration, O. Collaboration, and the LEP Electroweak Working Group, *Precision Electroweak Measurements and Constraints on the Standard Model*, 2007, URL <http://www.citebase.org/abstract?id=oai:arXiv.org:0712.0929>.
- [3] R. A. Dias, *Phenomenological analysis of the Two Higgs Doublet Model*, 2002, URL <http://www.citebase.org/abstract?id=oai:arXiv.org:hep-ph/0212237>.

- [4] M. Sher, *Scalar-Mediated Flavor-Changing Neutral Currents*, (1998), URL <http://www.citebase.org/abstract?id=oai:arXiv.org:hep-ph/9809590>,
- [5] R. Martinez, D. A. Milanes, and J. -Alexis Rodriguez, *Phys. Rev. D*, **72**, 035017 (2005), URL <http://www.citebase.org/abstract?id=oai:arXiv.org:hep-ph/0502087>.
- [6] J. Campbell, R. K. Ellis, F. Maltoni, and S. Willenbrock, 2003, URL, arXiv:hep-ph/0204093v2.
- [7] S. Dawson, C. B. Jackson, L. H. O. L. Reina, and D. Wackeroth, 2003, URL, arXiv:hep-ph/0311216v1.
- [8] J. Campbell, S. Dawson, S. Dittmaier, C. Jackson, M. Kramer, F. Maltoni, L. Reina, a M. Spira, D. Wackeroth, and S. Willen-

- brock 2004, URL [arXiv:hep-ph/0405302v1](http://arXiv.org/abs/hep-ph/0405302v1).
- [9] M. Kramer 2004, URL, [arXiv:hep-ph/0407080v1](http://arXiv.org/abs/hep-ph/0407080v1).
- [10] A. Collaboration, Tech. Rep. CERN (2000).
- [11] D0 Collaboration and V. Abazov, Phys. Rev. Lett. **95**, 151801 (2005),
URL <http://www.citebase.org/abstract?id=oai:arXiv.org:hep-ex/0504018>.
- [12] A. Pukhov et al, [arXiv:hep-ph/9908288](http://arXiv.org/abs/hep-ph/9908288), 2007.
- [13] M. Carena, S. Heinemeyer, C. E. M. Wagner, and G. Weiglein, European Physical Journal C, **45**, 797 (2006),
URL <http://www.citebase.org/abstract?id=oai:arXiv.org:hep-ph/0511023>.
- [14] P. K. Mal (2006), URL, [arXiv:hep-ex/0610083v1](http://arXiv.org/abs/hep-ex/0610083v1),