



Brazilian Journal of Physics

ISSN: 0103-9733

luizno.bjp@gmail.com

Sociedade Brasileira de Física
Brasil

Almeida, Eduardo de; Sharma, S. Shelly
SU(3) Model Description of Be Isotopes
Brazilian Journal of Physics, vol. 34, núm. 3A, september, 2004, pp. 962-965
Sociedade Brasileira de Física
São Paulo, Brasil

Available in: <http://www.redalyc.org/articulo.oa?id=46434568>

- How to cite
- Complete issue
- More information about this article
- Journal's homepage in redalyc.org

redalyc.org

Scientific Information System
Network of Scientific Journals from Latin America, the Caribbean, Spain and Portugal
Non-profit academic project, developed under the open access initiative

SU(3) Model Description of Be Isotopes

Eduardo de Almeida and S. Shelly Sharma

Departamento de Física, Universidade Estadual de Londrina, 86051-990, Londrina, PR, Brazil

Received on 20 October, 2003

Experimental data on light nuclei close to dripline suggests that as the nucleon number asymmetry increases, the shell structure from stability line is not preserved. In contrast with spherical shell model, Elliott's SU(3) model, uses a deformed multi-nucleon basis to describe nuclear states. The SU(3) symmetry is a strong feature of 1p shell nuclei, where symmetry breaking spin-orbit force is rather weak. We have calculated the binding energies and low-lying energy spectra of Be isotopes ($A=6$ to $A=14$), within the framework of Elliott's SU(3) model, with special emphasis on effects due to the presence of intruder orbit $1S_{1/2}$ in the region. The model space includes SU(3) basis states that maximize the quadrupole-quadrupole interaction ($-\chi Q \cdot Q$). An extended model space includes, in addition, a set of 2p-2h excitations, with excited nucleons occupying lowest energy SU(3) states in harmonic oscillator shell $N=2$. Group theoretical methods have been used to classify the states and calculate the nuclear interaction matrix elements. Good angular momentum states are projected out from intrinsic deformed SU(3) states by using standard angular momentum projection techniques. The interaction used contains the monopole-monopole, quadrupole-quadrupole and isospin dependent terms. Interaction parameters are fixed so as to reproduce the binding of 4 nucleons in $N=1$ orbit for the $N=Z$ isotope- ^8Be in ground state and first excited 2^+ state, along with the known systematics of single neutron separation energies for Be isotopes. The calculated energy spectra are compared with available experimental data. The calculated excitation energies of intruder states in ^8Be and ^{10}Be , are discussed in the light of available experimental evidence and theoretical results from shell model as well as deformed oscillator model calculations of Fayache et al..

1 Introduction

Shell model predictions for ground state spins and parities of Be isotopes with $6 \leq A \leq 10$ and $A = 12, 14$, are verified experimentally. For ^{11}Be and ^{13}Be nuclei, however, the model predicts ground state spin and parity values of $1/2^-$ and $5/2^+$, to be compared with the experimental values of $1/2^+$ and $1/2^-$ [1], respectively. Low lying spectra of ^9Be , ^{11}N , ^{11}Be and ^{13}Be , also point to a change in nuclear structure close to the drip line. The presence of positive parity intruder orbit $1S_{1/2}$, in the region of $N=1$ single particle orbits, offers a possible explanation for experimentally observed spin and parity values. As a consequence, intruder states are expected to be found in low lying spectra of Be isotopes [2, 3, 4].

The object of the present work is to understand the nuclear structure of Be isotopes in the low-energy region, within the framework of SU(3) model. The SU(3) model was proposed and used by Elliott [5] to describe rotational bands in light nuclei. Appropriate representations ($\lambda\mu$) of symmetry group SU(3) are used, in this model, to generate multi-nucleon basis states for a given nucleus. Using a four parameter residual interaction with monopole-monopole, quadrupole-quadrupole and isospin dependent interaction terms with strength parameters (P_0, P_1), χ , and β respectively, we have calculated the binding energies and low-lying spectra of Be isotopes. The ground state band is described by the representation ($\lambda\mu$) that maximizes the ground state binding energy for a given strength χ of quadrupole-quadrupole interaction. An extended model space including (2p-2h) excitations is then used to obtain

the likely position of intruder state in $^{8,10,12}\text{Be}$. Interaction parameters χ and P_0 are fitted to experimental ground state binding energy and excitation energy of 2_1^+ state in $N=Z$ isotope ^8Be . Interaction parameters P_1 and β are adjusted to give a best fit to experimental ground state binding energies of Be isotopes with $6 \leq A \leq 14$. The calculated energy spectra are compared with available experimental data. The New Mass Formula (NMF) of reference [6], that made possible the identification and prediction of new magic numbers, is also used to calculate the residual interaction. A comparison of results from NMF, our calculation and experimental data shows that the interaction used by us reproduces reasonably well the experimental ground state binding energies and neutron separation energies for Be isotopes.

2 SU(3) Model

Symmetry group SU(3) [7] has found many applications in nuclear physics as well as particle physics. We outline the main features of SU(3) model of Elliott [5], writing down the generators, subgroups and relevant Casimir operators. An SU(3) representation is characterized by quantum numbers ($\lambda\mu$), that determine the eigenvalue of Casimir operator for symmetry group SU(3). The states belonging to a given representation can be further labeled by quantum numbers that determine the eigenvalues of Casimir operators for subgroups of SU(3). Multi-nucleon deformed states, having good orbital angular momentum, can be projected out from a state belonging to a given representation of SU(3) group by using angular momentum projection techniques.

TABLE I. Casimir operators, Casimir operator eigenvalues and quantum numbers of SU(3) group and subgroups.

	SU(3)	SU(2)	U ₁ (1)	U ₂ (1)
Casimir op.	$(Q, Q+3L, L)$	$(3L_0^2+Q_2Q_{-2}+Q_{-2}Q_2)$	$L_0/2$	Q_0
eigenvalue	$\frac{2}{3}(\lambda^2 + \mu^2 + \lambda\mu + 3\lambda + 3\mu)$	$\Lambda(\Lambda + 1)$	$K/2$	ϵ
quantum number	$(\lambda\mu)$	Λ	K	ϵ

In spherical basis, three components of orbital angular momentum operator L , and five components of quadrupole moment operator Q constitute the eight generators of symmetry group SU(3). A component of quadrupole moment, is defined as

$$Q_q = \sqrt{\frac{4\pi}{5}} \alpha^2 \left(r^2 Y_{2,q}(\hat{r}) + \frac{p^2}{\alpha^2 \hbar} Y_{2,q}(\hat{p}) \right), \quad q = 0, \pm 1, \pm 2$$

where $\alpha^2 = m\omega/\hbar$ and $Y_{2,q}$ are the spherical harmonics. The eight SU(3) generators form a Lie algebra and satisfy the following commutation relations,

$$\begin{aligned} [L_q, L_{q'}] &= -\sqrt{2}(11qq'|1q+q') L_{q+q'}, \\ [Q_q, L_{q'}] &= -\sqrt{6}(21qq'|2q+q') Q_{q+q'}, \\ [Q_q, Q_{q'}] &= 3\sqrt{10}(22qq'|1q+q') L_{q+q'}. \end{aligned}$$

A convenient labeling scheme can be achieved by noting that the generators of symmetry group SU(2) are proportional to Q_2 , Q_{-2} and L_0 . Next L_0 and Q_0 are identified as the generators of subgroups U₁(1) and U₂(1), respectively. We show in Table I the relevant Casimir operators, Casimir operator eigenvalues, and the corresponding quantum numbers for symmetry group SU(3) and its subgroups. The quantum numbers $(\lambda\mu)$ characterize the permutation symmetry between harmonic oscillator quanta and Λ is proportional to the number of quanta in xy plane for the case $\mu = 0$ [8, 9].

Elliott [5] has shown that we can expand a state $\chi((\lambda\mu)_{\epsilon_{\max}} \Lambda K)$ belonging to SU(3) representation $(\lambda\mu)$ in terms of states $\Psi((\lambda\mu) K L K)$, having good angular momentum quantum number L that is

$$\chi((\lambda\mu)_{\epsilon_{\max}} \Lambda K) = \sum_L c(\lambda\mu K, L) \Psi((\lambda\mu) K L K),$$

where the coefficients c are real and ϵ_{\max} is the highest eigenvalue of operator Q_0 . We use the method of angular momentum projection and obtain the wave functions $\Psi((\lambda\mu) K L K)$, which are in turn used to obtain analytically the matrix elements of operator $Q \cdot Q$. A fortran routine has been written to calculate c 's and eventually matrix elements of $Q \cdot Q$ operator between relevant good L states.

3 Interaction Parameters

The nuclei studied are the isotopes of Be with $6 \leq A \leq 14$. We define the binding energy of isotope A in residual interaction zero state, $I(A)$ as

$$I(A) = B(A, Z) - B(4, 2) - 2S_p(5, 3) - (A-6)S_n(5, 2), \quad (1)$$

where $B(A, Z)$ is the binding energy of the nucleus with Z protons and $A-Z$ neutrons, $S_p(5, 3)$ is one proton separation energy of ${}^5\text{Li}$ and $S_n(5, 2)$ is one neutron separation energy of ${}^5\text{He}$. All terms on the right hand side in Eq. 1 can be obtained from available experimental data [1]. The experimental ground state residual interaction between active nucleons, $I(A)$, serves as a constraint on interaction parameters used in theoretical calculations.

We use an interaction Hamiltonian containing monopole-monopole, quadrupole-quadrupole and isospin dependent interaction terms that is

$$H_{res} = -F_0(\underline{n}) - \chi Q \cdot Q + \beta T(T+1), \quad (2)$$

where \underline{n} is the number of active nucleons ($\underline{n} = A-4$). The isospin dependent interaction is repulsive while the monopole-monopole and quadrupole-quadrupole interactions are attractive. The strength of monopole-monopole interaction for nucleons in the same shell is P_0 and for nucleons in different shells is P_1 . We write the monopole-monopole interaction for n_1 and n_2 ($\underline{n} = n_1 + n_2$) nucleons in oscillator shells $N=1$ and $N=2$, respectively, as

$$F_0(\underline{n}) = P_0 \frac{n_1(n_1-1)}{2} + P_0 \frac{n_2(n_2-1)}{2} + P_1 n_1 n_2.$$

From Table I, the quadrupole-quadrupole interaction operator may be expressed as

$$-\chi Q \cdot Q = -\chi 3(2C_{SU(3)} - L \cdot L),$$

where $C_{SU(3)}$ is the Casimir operator of the SU(3) group. The ground state isospin is defined as $T = \left| \frac{(n-p)}{2} \right|$, where n and p are the number of neutrons and protons respectively.

3.1 New Mass Formula (NMF)

Comparing the mass formula of Bethe-Weizsacker [10, 11] with the New Mass Formula (NMF) [6], the NMF has a new parameter $\Delta(N, Z)$ and has a redefinition of the pairing term δ_{new} . The NMF is written as

$$\begin{aligned} B(A, Z)_{new} &= a_v A - a_s A^{2/3} - a_c \frac{Z(Z-1)}{A^{1/3}} \\ &\quad - a_{sym} \frac{(A-2Z)^2}{A} + \delta_{new} + \Delta(N, Z), \end{aligned}$$

TABLE II. Values of $(\lambda_1\mu_1 L_1)$ and $(\lambda_2\mu_2 L_2)$ for active nucleons in N=1 and N=2 oscillator shells.

	g. s. band				2p-2h excitations			
	$\lambda_1\mu_1$	L_1	$\lambda_2\mu_2$	L_2	$\lambda_1\mu_1$	L_1	$\lambda_2\mu_2$	L_2
${}^6\text{Be}$	20	0,2						
${}^7\text{Be}$	30	1,3						
${}^8\text{Be}$	40	0,2,4			20	0,2	40	0,2,4
${}^9\text{Be}$	31	1,2,3,4			30	1,3	40	0,2,4
${}^{10}\text{Be}$	22	0,2 ² ,3,4			40	0,2,4	40	0,2,4
${}^{11}\text{Be}$	22	0,2 ² ,3,4	20	0,2	40	0,2,4	41	1,2,3,4,5
${}^{12}\text{Be}$	22	0,2 ² ,3,4	40	0,2,4	20	0,2		
${}^{13}\text{Be}$	21	1,2,3	40	0,2,4	31	1,2,3,4	42	0,2 ² ,4 ² ,3,5,6
${}^{14}\text{Be}$	20	0,2	40	0,2,4	22	0,2 ² ,3,4	42	0,2 ² ,4 ² ,3,5,6

where $a_v = 15,85$ MeV, $a_s = 18,34$ MeV, $a_c = 0,71$ MeV, $a_{\text{sym}} = 23,21$ MeV, $\delta_{\text{new}} = (1 - \exp(-\frac{Z}{c}))\delta$, $c = 6/\ln 2$, δ is equal to $12A^{-1/2}$ for even-even nuclei, $-12A^{-1/2}$ for odd-odd nuclei, 0 for odd A nuclei and $\Delta(N, Z) = |N - \frac{4}{3}Z|N^k Z \exp(-\frac{Z}{3})$ with $k = 0,45$.

Using the NMF to calculate the binding energy we obtain the residual interaction $I_{\text{new}}(A)$ for Be isotopes as

$$I_{\text{new}}(A) = B_{\text{new}}(A, 4) - B(4, 2) - 2S_p(5, 3) - (A-6)S_n(5, 2). \quad (3)$$

4 Model Space

Table II lists the representations $(\lambda\mu)$ that maximize the quadrupole-quadrupole interaction, and the possible L values for the ground state band. The representations $(\lambda\mu)$ and possible angular momenta for the case when two nucleons are excited out of ground state (2p-2h excitations) are also presented in Table II.

We predict correct spin and parity for ${}^{11}\text{Be}$ ground state if we consider the last neutron occupying N=2 oscillator orbit. The isotopes, ${}^{11}\text{Be}$, ${}^{12}\text{Be}$, ${}^{13}\text{Be}$ and ${}^{14}\text{Be}$ have at least one neutron occupying the N=2 oscillator shell, in the ground state. Therefore we must subtract $\hbar\omega$ from residual interaction for ${}^{11}\text{Be}$ and $2\hbar\omega$ for ${}^{12}\text{Be}$, ${}^{13}\text{Be}$ and ${}^{14}\text{Be}$. We use the formula $\hbar\omega = 45A^{-1/3} - 40A^{-2/3}$ to get the value of oscillator constant. In comparison Fayache et al [4] have used $\hbar\omega = 45A^{-1/3} - 25A^{-2/3}$, with a milder surface dependence.

5 Results and Conclusions

In Fig. 1 we plot the ground state binding energies of Be isotopes, due to residual interaction between active nucleons, (i) calculated by using SU(3) model, (ii) calculated from New Mass Formula (NMF Eq. 3) and (iii) obtained from experimental data (Eq. 1). The parameter $\chi = 0.17$ MeV reproduces the excitation energy of 2_1^+ state in N=Z isotope ${}^8\text{Be}$. For this choice of χ , a parameter value of $P_0 = 2.3$ MeV is needed to obtain experimental ground state binding

energy of ${}^8\text{Be}$. Interaction parameters P_1 and β are next adjusted to give a best fit to experimental ground state binding energies of Be isotopes with $6 \leq A \leq 14$. The fit shown in Fig. 1 is obtained with $P_1 = 2.2$ MeV and $\beta = 4.1$ MeV. We can verify that the calculated binding energies for Be isotopes show a good agreement with experimental values. Correct ground state spins and parities are reproduced including the inversion of spectra in ${}^{11}\text{Be}$. We notice that the results of New Mass Formula, also reproduce the trend of experimental binding energies of nuclei obtained by adding neutrons to ${}^8\text{Be}$.




Figure 1. Experimental and calculated spectra for ${}^8\text{Be}$, ${}^{10}\text{Be}$ and ${}^{12}\text{Be}$. Calculated (2p-2h) excitation states are shown in square brackets.

In Fig. 2, we compare with experimental and calculated spectra for ${}^8, {}^{10}, {}^{12}\text{Be}$ obtained by using the interaction of Eq. 2 with the set of parameter values chosen earlier. In this calculation we have used the states defined by SU(3) representations and allowed angular momenta listed in Table II. The states obtained by using representations with (2p-2h) excitations are shown in square brackets in Fig. 2. We observe that the low lying states of ${}^8\text{Be}$ are not likely to mix strongly with states from (2p-2h) excitations. Table III lists the calculated and experimental excitation energies of $0_1^+, 2_1^+, 0_2^+$, and 2_2^+ states for comparison. The energy difference between the

TABLE III. Excitation energies of 0_1^+ , 2_1^+ , 0_2^+ , and 2_2^+ states in ^8Be , ^{10}Be and ^{12}Be nuclei in MeV. Square brackets represent states arising from (2p-2h) excited configurations.

^8Be				^{10}Be				^{12}Be			
Calc.		Expt.		Calc.		Expt.		Calc.		Expt.	
0_1^+	0.0	0_1^+	0.0	0_1^+	0.0	0_1^+	0.0	0_1^+	0.0	0_1^+	0.0
2_1^+	3.04	2_1^+	3.04	2_1^+	3.04	2_1^+	3.37	2_1^+	3.04	2_1^+	2.10
$[0_2^+]$	18.68	0_2^+	20.2	$[0_2^+]$	3.83	0_2^+	6.18	$[0_2^+]$	2.98		
$[2_2^+]$	21.70			$[2_2^+]$	6.86			$[2_2^+]$	6.01		




Figure 2. Calculated (SU(3) model and NMF) and experimental ground state binding energy due to residual interaction between active nucleons for Be Isotopes.

states 2_2^+ and 2_1^+ (2p-2h) for ^8Be is 18.66 MeV while for ^{10}Be and ^{12}Be it is just 3.82 MeV and 3.97 MeV respectively. Therefore the ground states as well as the 2_1^+ states in nuclei ^{10}Be and ^{12}Be are expected to have strong contributions from (2p-2h) configurations. We are studying the effect of such mixing on radii and transitions of Be isotopes.

The calculated excitation energy of 0^+ (2p-2h) intruder state in ^8Be is 18.6 MeV. Experimentally a 0^+ state is seen at 20.2 MeV. Using a deformed oscillator model calculation Fayache et al [4] have reported, a value of 17.23 MeV in ^8Be . As such our SU(3) model result is close to the calculated value from deformed oscillator model and not far from the experimentally observed 0^+ state energy. For the case of ^{10}Be our result of 3.83 MeV and the value of 6.36 MeV

calculated in ref. [4] may possibly be compared with the energy of a 0^+ state seen at 6.18 MeV, experimentally. We may point out, however that our results are preliminary results. As pointed out by Fayache et al.[4], it is essential to take into consideration triaxial nature of the ground state in ^{10}Be .

Acknowledgements

Financial support from CAPES, Brazil and Fundação Araucaria is acknowledged.

References

- [1] National Nuclear Data Center, <http://www.nndc.bnl.gov/nndc/ensdf/>.
- [2] F. C. Barker, Aust. J. Phys. **41**, 743 (1988).
- [3] F. C. Barker, Aust. J. Phys. **42**, 25 (1989).
- [4] M. S. Fayache, E. Moya de Guerra, P. Sarriguren, Y. Y. Sharon, and L. Zamick, Phys. Rev. C **57**, 2351 (1998).
- [5] J. P. Elliott, Proc. R. Soc. (London) A **245**, 128 (1958); Proc. R. Soc. (London) A **245**, 562 (1958).
- [6] C. Samanta and S. Adhikari, Phys. Rev. C **65**, 037301 (2002).
- [7] M. Hamermesh, Group Theory and its applications to Physical problems, Addison-Wesley, Reading MA 1962.
- [8] V. G. Gueorguiev and J. P. Draayer, Revista Mexicana de Fisica **44**, 2 (1999).
- [9] A. Bohr and B. R. Mottelson, Nuclear Structure, V. 1, W. A. Benjamin, Inc., (1969).
- [10] C. F. von Weizsäcker, Z. Physik **96**, 431 (1935).
- [11] H. Bethe, Rev. Mod. Phys. **8**, 82 (1936).