



Brazilian Journal of Physics

ISSN: 0103-9733

luizno.bjp@gmail.com

Sociedade Brasileira de Física

Brasil

C. de Souza, José C.; Saa, Alberto  
Phase Space Solutions in Scalar-Tensor Cosmological Models  
Brazilian Journal of Physics, vol. 35, núm. 4B, december, 2005, pp. 1041-1043  
Sociedade Brasileira de Física  
São Paulo, Brasil

Available in: <http://www.redalyc.org/articulo.oa?id=46435704>

- How to cite
- Complete issue
- More information about this article
- Journal's homepage in redalyc.org

redalyc.org

Scientific Information System  
Network of Scientific Journals from Latin America, the Caribbean, Spain and Portugal  
Non-profit academic project, developed under the open access initiative

## Phase Space Solutions in Scalar-Tensor Cosmological Models

José C. C. de Souza

*Departamento de Física Matemática, Instituto de Física,  
Universidade de São Paulo, C.P. 66318, 05315-970 São Paulo, SP, Brazil*

and Alberto Saa

*Departamento de Matemática Aplicada, IMECC – UNICAMP, C.P. 6065, 13083-859 Campinas, SP, Brazil  
(Received on 15 October, 2005)*

An analysis of the solutions for the field equations of generalized scalar-tensor theories of gravitation is performed through the study of the geometry of the phase space and the stability of the solutions, with special interest in the Brans-Dicke model. Particularly, we believe to be possible to find suitable forms of the Brans-Dicke parameter  $\omega$  and potential  $V$  of the scalar field, using the dynamical systems approach, in such a way that they can be fitted in the present observed scenario of the Universe.

### I. SCALAR-TENSOR THEORIES OF GRAVITATION

In a homogeneous and isotropic space, described by the Friedmann-Lemaître-Robertson-Walker metric

$$ds^2 = -dt^2 + a^2(t) \left[ \frac{dr^2}{1 - Kr^2} + r^2(d\theta^2 + \sin^2\theta d\phi^2) \right], \quad (1)$$

where  $a$  is the scale factor and  $K$  is the spatial curvature index, gravitation can be described by an action of the kind

$$S = \frac{1}{16\pi} \int d^4x \sqrt{-g} \left[ \phi R - \frac{\omega(\phi)}{\phi} g^{ab} \nabla_a \phi \nabla_b \phi - V(\phi) \right] + S^m, \quad (2)$$

where  $S^m$  is the action of usual matter,  $g$  is the determinant of the metric tensor,  $\omega$  is a coupling function (which we will eventually assume to be a constant, known as Brans-Dicke parameter) and  $V(\phi)$  is the potential of the scalar field  $\phi$  [2].

From (2), we obtain for the field equations:

$$H^2 = -H \left( \frac{\dot{\phi}}{\phi} \right) + \frac{\omega}{6} \left( \frac{\dot{\phi}}{\phi} \right)^2 + \frac{V(\phi)}{6\phi} - \frac{K}{a^2} + \frac{8\pi\rho^m}{3\phi}, \quad (3)$$

$$\begin{aligned} \dot{H} = & -\frac{\omega}{2} \left( \frac{\dot{\phi}}{\phi} \right)^2 + 2H \left( \frac{\dot{\phi}}{\phi} \right) \\ & + \frac{1}{2(2\omega+3)\phi} \left[ \phi \frac{dV}{d\phi} - 2V + \frac{d\omega}{d\phi} (\dot{\phi})^2 \right] \\ & + \frac{K}{a^2} - \frac{8\pi}{(2\omega+3)\phi} [(\omega+2)\rho^m + \omega P^m], \end{aligned} \quad (4)$$

$$\ddot{\phi} + \left( 3H + \frac{1}{2\omega+3} \frac{d\omega}{d\phi} \right) \dot{\phi} = \frac{1}{2\omega+3} \left[ 2V - \phi \frac{dV}{d\phi} + 8\pi(\rho^m - 3P^m) \right], \quad (5)$$

where  $H \equiv \dot{a}/a$  is the Hubble parameter and  $\rho^m$  and  $P^m$  are the energy density and the pressure of the material fluid.

As usual, we parameterize the equation of state for the fluid as  $P^m = (\gamma - 1)\rho^m$  with  $\gamma$  a constant chosen to indicate a variety of fluids that are predominantly responsible for the energy density of the Universe. We can see that through the energy conservation equation  $\dot{\rho}^m + 3H(\rho^m + P^m) = 0$  we obtain  $\rho^m = \rho_0/a^{3\gamma}$ , with  $\rho_0$  a constant.




FIG. 1: Upper sheet of the of the phase space for a model with  $\omega = 10$  (Brans-Dicke), corresponding to the positive sign in eq. (7) [2]. The “hole” in the surface indicates the region forbidden for the orbits of the solutions.

### II. THE CASE FOR $V = \frac{1}{2}m^2\phi^2$ AND $K = 0$

In the referred paper [2], the author proceeds to show the phase space allowed for the orbits of solutions for these field equations in several cases with different potentials and parameters  $\omega$ . For example, in the case of vacuum, flat space ( $K = 0$ ) and potential  $V = \frac{1}{2}m^2\phi^2$ , equation (3) was rearranged as (making  $m \equiv 1$ )

$$\omega\dot{\phi}^2 - 6H\phi\dot{\phi} + \left( \frac{1}{2}\phi^2 - 6H^2\phi \right)\phi = 0, \quad (6)$$

which has the solutions

$$\dot{\phi}_{\pm}(H, \phi) = \frac{1}{\omega} \left[ 3H\phi \pm \sqrt{3(2\omega+3)H^2\phi^2 - \frac{1}{2}\omega\phi^2} \right]. \quad (7)$$

The assumption of flat space is required in order to reduce the dimensionality of the phase space.

We want to analyze qualitatively the geometry of the phase space  $(H, \phi, \dot{\phi})$ , expecting to infer the form of the functions  $\omega(\phi)$  and  $V(\phi)$  to fit better the available data on the structure of the Universe.

The phase space for this situation is composed of a 2-d surface with two sheets, related to the lower and upper signs in eq. (7). Figures 1-3 show the phase space for the choice  $\omega = 10$ .

The fixed points for this dynamical system, obtained making  $\dot{H} = \dot{\phi} = 0$ , are de Sitter solutions, given by  $H_0 = \pm\sqrt{\phi_0/12}$ , with constant  $H_0$  and  $\phi_0$ .




FIG. 2: Lower sheet of the phase space, now corresponding to the negative sign in eq. (7).




FIG. 3: The complete phase space composed of the upper and lower sheets linked to each other at the boundary of the forbidden region.

### III. THE CASE FOR $V = \Lambda\phi$ AND $K = 0$

Following other works ([3]-[5]) which give a complete analysis of the phase space for Brans-Dicke model with a cosmological constant  $\Lambda$  (simply making  $V(\phi) = \Lambda\phi$  in the action), we can illustrate the situation in which  $\omega$  has a very large value and  $\gamma = 0$ . Therefore, the energy density of the fluid is a constant  $\rho_0$ . It should be emphasized that recent observational and simulation results seem to favor a scenario very similar to this one ([6],[11]-[14]). The solutions in this case are written as

$$\dot{\phi}_{\pm}(H, \phi) = \frac{1}{\omega} \left[ 3H\phi \pm \sqrt{9H^2\phi^2 - \omega[\phi^2(\Lambda - 6H^2) + 16\pi\rho_0]} \right]. \quad (8)$$

With this solutions, we are able to show the phase space for a particular choice of constants  $\Lambda$ ,  $\omega$  and  $\rho_0$ .

We proceed to find the dynamical equations system for this simple model, as done before.

Naming  $\Delta$  the expression under the root in eq.(8), we can write the equation for  $\dot{H}$ :

$$\begin{aligned} \dot{H}_{\pm} = & -\frac{1}{2\omega\phi^2} \left[ 3H\phi \pm \sqrt{\Delta} \right]^2 + \frac{2H}{\omega} \left[ 3H \pm \frac{\sqrt{\Delta}}{\phi} \right] \\ & - \frac{1}{2(2\omega+3)} \left( \frac{\Lambda}{2} + \frac{16\pi\rho_0}{\phi} \right). \end{aligned} \quad (9)$$

Now, equations (8) and (9) form the system for which the fixed points are the solutions  $H_0 = \pm \sqrt{8\pi\rho_0/3\phi_0^2 + \Lambda/6}$ .

Of special interest is the search for the most adequate functions  $\omega(\phi)$  and  $V(\phi)$ , that may be more complicated than what was assumed until here.




FIG. 4: Complete phase space for a Brans-Dicke model with a cosmological constant  $\Lambda = 1$ , energy density  $\rho_0 = 2$  and constant parameter  $\omega = 50000$ , showing two sheets linked by the boundary of the forbidden region, as in the precedent case.

### IV. CONCLUSIONS

The method of analyzing the geometry of the phase space have proved to be a useful tool in the search for the solutions of the field equations of generalized gravity models. Our aim is to achieve a complete analysis of the simple model presented before (including the stability of the solutions, via Lyapunov's direct method [1], in order to investigate further its *attraction basin*) and to apply more sophisticated functions to it.

### V. ACKNOWLEDGMENTS

This work is supported by Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq). We would like to thank F. C. Carvalho for useful discussions.

- [1] J. LaSalle and S. Lefschetz, *Stability by Lyapunov's Direct Method*. Academic Press (1967)
- [2] V. Faraoni, *Annals of Physics* **317**, 366 (2005)

- [3] S. J. Kolitch, *Annals of Physics* **246**, 121 (1996)
- [4] S. J. Kolitch, *Annals of Physics* **241**, 128 (1995)
- [5] C. Santos and R. Gregory, *Annals of Physics* **258**, 111 (1997)

- [6] A. G. Sanchez et al., astro-ph/0507583
- [7] G. Esposito-Farise and D. Polarski, Phys. Rev. D **63**, 063504 (2001)
- [8] A. Saa et al., Phys. Rev. D **63** 067301 (2001); Int. J. Theor. Phys. **40**, 2295 (2001); L.R. Abramo, L. Brenig, E. Gunzig, and A. Saa, Phys. Rev. D **67**, 027301 (2003); gr-qc/0305008.
- [9] J. D. Barrow and J. P. Mimoso, Phys. Rev. D **50**(6), 3746 (1994)
- [10] F. C. Carvalho and A. Saa, Phys. Rev. D **70**, 087302 (2004)
- [11] G. Esposito-Farise, gr-qc/0409081
- [12] B. Bertotti, L. Iess, and P. Tortora, Nature **425**, 374 (2003)
- [13] V. Acquaviva, C. Baccigalupi, S. M. Leach, Andrew R. Liddle, and F. Perrotta, astro-ph/0412052
- [14] A. R. Liddle, A. Mazumdar, and J. D. Barrow, Phys. Rev. D **58**, 027302 (1998)