



Ciência e Natura

ISSN: 0100-8307

cienciaenaturarevista@gmail.com

Universidade Federal de Santa Maria  
Brasil

Poursharifi, M.; Shahi, S.  
Microwave Irradiation Assisted Synthesis 4-Hydroxy 3-Iodobenzylguanidine(HIBG)  
Ciência e Natura, vol. 37, núm. 6-1, 2015, pp. 71-74  
Universidade Federal de Santa Maria  
Santa Maria, Brasil

Available in: <http://www.redalyc.org/articulo.oa?id=467547682011>

- How to cite
- Complete issue
- More information about this article
- Journal's homepage in redalyc.org

redalyc.org

Scientific Information System  
Network of Scientific Journals from Latin America, the Caribbean, Spain and Portugal  
Non-profit academic project, developed under the open access initiative

## Microwave Irradiation Assisted Synthesis 4-Hydroxy 3-Iodobenzylguanidine(HIBG)

M.Poursharifi<sup>1\*</sup> - S.Shahi<sup>2</sup>

1-Department of Chemistry management,Tehran science and research branch Islamic Azad University

\*M.Poursharifi@srbiau.ac.ir

2- Department of Chemistry, Faculty of science, University of Guilan, Rasht, Iran

### Abstract


*Meta-iodo benzyl guanidine(MIBG) is an analogue of Noradrenalin transmitter. When radiolabelled with  $^{123}\text{I}$  and  $^{131}\text{I}$ , used for scan and diagnosis Adrenal's tumor. Labeled MIBG with  $^{131}\text{I}$  used in therapy of Neuroblastoma, Pheochromocytoma, Carcinoidsyndrome and Neuroendocrine tumors. In our continuing efforts to develop MIBG analogues with higher therapeutic efficacy, we recently evaluated the structure-activity characteristics of several analogues of MIBG. We are interested in developing HIBG analogues containing hydroxyl at the 4-postion oF HIBG.generous of the some groups to activate the aromatic ring of a drug towards electrophilic radiodiodination may have merit .The resistance of HIBG to in vivo radiodiodination and amenability to kit production make it a strong candidate for clinical evaluation as a pheochromocytoma and neuroblastoma localzing agent. Microwave assisted heating under controlled conditions has been shown to be an invaluable technology for organic chemistry and during discovery applications since it often rise yield and dramatically reduce reaction times, typically form days or hours to minutes or even seconds.*

*Microwave Irradiation Assisted Synthesis of HIBG is started to produce 4-methoxybenzylamine. 4-methoxybenzylamine and hydriodic acid are mixed and heated for this purpose. Then under MBR condition 4-Hydroxybenzyl guanidine in sulphate form is prepared with additional Cyanamide and Sulphuric acid. After that Iodination is preformed under MBR condition and 4-Hydroxy-3-iodobenzyl guanidine sulphate formed, that can labeled with  $^{131}\text{I}$  and exchange to labeled 4-Hydroxy-3-iodobenzyl guanidine sulphate.*


**Keywords:** Radiolabelling- MIBG- HIBG- Neuroblastoma- Pheochromocytoma- Carrcinoidsyndrome- Microwave Batch Reactor

## 1. Introduction


In recent years, Synthesis and labeling of MIBG analogues which are used in scan and diagnosis of adrenal tumors have become highly important. In this research, synthesis of HIBG is examined microwave irradiation assisted that resulted in higher therapeutic effect of HIBG compared to MIBG, plus a faster synthesis and higher efficiency when microwave irradiation assisted than thermal method.



Schema.1



Schema.2



Schema.3

## 2. EXPERIMENTAL METHOD

Chloramine-T , 4-methoxybenzylamine , sodium iodide , hydriodic acid and Cyanamide were provided in the laboratory microwave from laboratory resources. Proton magnetic resonance (HNMR) and infrared spectrum (IR) of them were obtained.

### 2.2 TASK PROCEDURE

#### 2.2.1 4-Hydroxybenzylamin hydroiodide

4-methoxybenzylamine(125mg, 1mmol) was MBR irradiation assisted for 20min with hydriodic acid(184mg , 3mmol)(German Merck) . methyl iodide(bp:30°C) was vacuum evaporated and extra HI was isolated with appropriate solvents (methyl acetate , acetic acid)after 3 times of recrystallization. needle crystals of 4-hydroxybenzylaminhydroiodide were appeared .(88% yield) (Shema.1)

m.p:182° C

IR(KBr)  $\nu_{\max}/\text{cm}^{-1}$ : 3400(OH), 3161.25(NH), 3006.77(NH),3000(sp<sup>2</sup> C-H, Arom)

#### 2.2.2 4-Hydroxybenzylguanidine sulphate

4-Hydroxybenzylamin hydroiodide (502mg , 2mmol) and Cyanamide (252mg , 0.6mmol) were irradiation at 150°C for 15minutes then they were cool to ambient temperature. The Remaining was solved in boiling water. Dissolved KHCO<sub>3</sub> (220mg , 2.2mmol) in 2ml of water, was added to the solution . The resulting solution was boiled for 2minutes then was left to be cool to ambient temperature. Filtrated product of bicarbonate salt made in 4ml hot water as a suspension . Added H<sub>2</sub>SO<sub>4</sub>(2N) drop by drop until pH became 2. It boiled to become dense to 1ml, then were cool. Colorless solid filtrated . it was recrystallized once more with water to achieve sulphat solid as a colorless crystal.(380mg , 90% ) (Shema.2)

m.p: 260-262 °C

HNMR(300 MHZ ,MeOH)δ: 7.11(d,2H , J=8HZ), 6.74(d,2H , J=8HZ),4.25(s,2H)

### 2.2.3 4-hydroxy-3-iodobenzylguanidine sulphate

Dissolved 4-Hydroxybenzylguanidine sulphate (225mg , 1.05mmol) in EtOH (30% , 15ml) and dense ammoniac solution and added a solution of I<sub>2</sub> (199.9mg , 0.788mmol) in EtOH(15ml) in 0 °C were mixed. Once more in ambient temperature mixed for 18hours. In vacuum, solvent evaporated and the rest was recrystallized three times with water solvent to achieve (149mg , 82%) (Shema.3)

m.p:208-210 °C

<sup>1</sup>HNMR(300MHz , MeOD)δ:6.76-7.65(m , 6H , ArH,NH<sub>2</sub>), 4.10(s,2H,CH<sub>2</sub>)




Figure3:<sup>1</sup>HNMR of 4-hydroxy-3-iodobenzyl guanidine sulphate




Figure1:<sup>1</sup>HNMR of 4-Hydroxybenzylguanidine sulphate




Figure4:<sup>13</sup>CNMR of 4-hydroxy-3-iodobenzyl guanidine sulphate




Figure2:<sup>13</sup>CNMR of 4-Hydroxybenzylguanidine sulphate

## 3. RESULT DISSCUSION

4-hydroxy-3-iodobenzyl guanidine sulphate labeled easily with carrier added and nocarrier added structures. In addition, containing hydroxyl group in para-position , radiation iodo stabled in 3-position . this function reduce large amount of releasing scatter iodine in drug. 4-hydroxybenzyl guanidine sulphate (HBG) is an intermediate that has the ability to be labeled in the clinics by provided kit production. The far hospitals from central nuclear may have merit. Microwave irradiation assisted synthesis reduce reaction time with rise yield.

#### 4. REFERENCES

- 1.H.Salehi, M.Poursharifi , M.Rezaei , M.Gourani , 12th Annal Iranian Coungress on Necleur Medicene . 25.28,2008
- 2.Helen Lee , Muthiah N.Inbasekaran , et al:Development of kit-form analogs of metaiodobenzylguanidine. I Nucl Med 27:256-267 , 1986
- 3.G.Vaidyanathan , D.J.Affeck , M.R.Zalutsky . No –carrier-added synthesis of a 4-methyl –substituted meta-iodobenzylguanidine analogue.
- 4.G.Vaidyanathan, D.J.Affeck , M.R.Zalutsky : (4-Fluoro-3-iodobenzylguanidine) , a potential MIBG analogue for positron Emission Tomography J.Med . Chem. 37,3655-3662, 1994
- 5.G.Vaidyanathan, D.J.Affeck , M.R. Zalutsky: synthesis of king-and side chain-substituted meta-iodobenzylguanidine analogue. Bioconjugate chem 2001,19,786-797
- 6.Ganesan Vaidyanathen, Sriram Shsnkar, : meta-iodobenzylguanidine donna J. Affeck, et al derivatives containing asecond guanidine moiety. Elsevier Bioorg and Med Chem , 12:1649-1656, 2004
- 7.Radiopharmaceuticls in Nuclear Pharmacy and Nuclear Medicine. Second Edition, Ed By R.J.Kowalsky and S W Falen 2004. Pg 310-314,713-716
- 8.Knickmeier, M, Matheja, P., et .al. clinical evaluation of no-carier m-iodobenzylguanidine for myocardial scintgrraphy. EJNM 2000; No.3.,302-307
- 9.T. Yassine, M. A. Bakir, S..Shanan, M. AL-Asaad: Preparation, purification and quality control of <sup>131</sup>I-m-iodobenzylguanidine. J RAD Nucl Chem, 285,3,635-637,2003
- 10.G. Prabhakar , Kiran S. Mehra , N. Ramamoorthy , et al :Evaluation of radioiodination of meta-iodobenzylguanidine (MIBG) catalyzed by in situ generated Cu(I) and directly added Cu(II). Pergamon Rad and Iso. 50,1011-1014, 1999
11. Inbasekaran M, Sherman PS , Brown LE , et al: Development of kit-form analogs of meta-iodobenzylguanidine. J Nucl Med 24:P117 , 1983 (a bstr)
12. Inbasekaran M , Lee H , Wieland DM , et al : Radiopharmaceuticls Selective for Adrenergic Nevers , 186<sup>th</sup> American Chemical Society National Meeting , Washington , DC , Aug. 28-Sept.2,1983
- 13.Sisson JC,Frager MS,Valk TW,et al: Scintirgraphic localization of pheochromocytoma .N Engl J Med 305:12-17