



Journal of the Mexican Chemical Society

ISSN: 1870-249X

editor.jmcs@gmail.com

Sociedad Química de México

México

Mendoza, Jorge A.; Jiménez Vázquez, Hugo A.; Herrera, Rafael; Liu, Jide; Tamariz, Joaquín
Synthesis, Structural, and Theoretical Study of New β -Heterosubstituted Captodative Olefins 1-Acetylvinylnyl Arenecarboxylates

Journal of the Mexican Chemical Society, vol. 47, núm. 2, abril-junio, 2003, pp. 108-116

Sociedad Química de México

Distrito Federal, México

Available in: <http://www.redalyc.org/articulo.oa?id=47547204>

- How to cite
- Complete issue
- More information about this article
- Journal's homepage in redalyc.org

redalyc.org

Scientific Information System

Network of Scientific Journals from Latin America, the Caribbean, Spain and Portugal

Non-profit academic project, developed under the open access initiative

Investigación

Synthesis, Structural, and Theoretical Study of New β -Heterosubstituted Captodative Olefins 1-Acetylvinyl Arenecarboxylates

Jorge A. Mendoza,¹ Hugo A. Jiménez-Vázquez,¹ Rafael Herrera,² Jide Liu,¹ and Joaquín Tamariz^{1*}

¹ Departamento de Química Orgánica, Escuela Nacional de Ciencias Biológicas, Instituto Politécnico Nacional, Prol. de Carpio y Plan de Ayala, 11340 México, D.F. México. Tel: (+5255) 5729-6300 / 62411; Fax: (+5255) 5396-3503; E-mail: jtamariz@woodward.enqb.ipn.mx

² Instituto de Investigaciones Químico-biológicas, Universidad Michoacana de San Nicolás de Hidalgo, Edif. B-1, Ciudad Universitaria, Francisco J. Mujica S/N, 58066 Morelia, Mich., México.

Dedicated to Dr. Alfonso Romo de Vivar on the occasion of his 50th anniversary of relevant contributions in natural products chemistry at the Instituto de Química, UNAM

Recibido el 10 de diciembre del 2002; aceptado el 6 de febrero del 2003

Abstract. A new series of β -heterosubstituted captodative olefins 1-acetylvinyl arenecarboxylates (**7a** and **10a-10h**) has been prepared, by introducing nitrogen or sulphur as heteroatoms substituted by alkyl and aryl groups. Three preparation methods were evaluated by modifying the leaving group of the starting material, as well as the nucleophilic character of the adding thiols. All of them were efficient and stereoselective, providing the desired alkenes in good yields and with the *Z* configuration of the double bond. *Ab initio* calculations (HF/6-31G*) of the FMOs, of some of the *beta* amino and bromo olefins, explained their experimental reactivity in Diels-Alder additions with respect to the unsubstituted olefin **1a**. It also appears that the HOMO and LUMO energies of the *beta* sulphur analogues are governed by the particular electronic features of the sulphur atom, and that their very low reactivity before a diene is due to steric hindrance. A comparison between bond distances obtained by X-ray crystallography of different β -substituted and unsubstituted olefins seems to correlate with the delocalization effect of the heteroatom lone electron pair for the bromo and amino β -substituted olefins.

Keywords: Captodative olefin, structure, reactivity.

Resumen. Se describe la preparación de una nueva serie de olefinas captodativas β -heterosustituidas 1-acetilvinil arenecarboxilatos (**7a** and **10a-10h**), donde el heteroátomo es nitrógeno o azufre, sustituido por grupos alquilo y arilo. Se evaluaron tres métodos para su preparación, modificando el grupo saliente en el sustrato, y el carácter nucleofílico de los tioles que se adicionaron. Los tres métodos fueron eficientes y estereoselectivos, proporcionando los alquenos deseados en buenos rendimientos y con la configuración *Z* del doble enlace. El cálculo de orbitales moleculares frontera (*ab initio*, HF / 6-31G*) de algunas de las olefinas preparadas permitió explicar sus diferencias en reactividad experimental con respecto a la olefina no sustituida **1a**. Así, se sugiere que las energías de los orbitales HOMO y LUMO de los análogos azufrados están gobernadas por las propiedades electrónicas particulares del átomo de azufre, y que su baja reactividad ante un dieno depende también del efecto estérico. Una comparación de los datos de cristalografía de rayos X entre distancias de enlace de diferentes olefinas β -sustituidas y la no sustituida parece correlacionarse con el efecto de deslocalización del par de electrones no compartidos del heteroátomo para las olefinas β -bromo y β -amino sustituidas.

Palabras clave: Olefina captodativa, estructura, reactividad.

Introduction

Captodative olefins have attracted particular attention in recent years, due to the opposite electronic demand and to the synthetic potential displayed by their geminally substituted functional groups [1]. We have shown that 1-acetylvinyl *p*-arenecarboxylates **1a-1c** were highly reactive and selective in Diels-Alder [2] and 1,3-dipolar cycloadditions [3], and they also proved to be very useful synthons in natural product synthesis [4]. More recently, the alkyl 2-aryloxy acrylates **2a-2b** were prepared, showing also high reactivity and selectivity in Diels-Alder reactions [5]. With the aim of evaluating the effect of a third substituent in the double bond on the reactivity in [4+2] additions, compound **3a** was prepared through a stereoselective synthetic route starting from **1a** [6]. A series of amines **3b** and thiols **3c** and **3d** were synthesized by treatment of **3a** with the corresponding amines and thiols. In particular,

enaminones are important organic intermediates [7], and have potential biological activity [8]. Moreover, other β -substituted captodative olefins have been prepared, showing interesting pericyclic behavior and synthetic usefulness [9].

Attempts to carry out the Diels-Alder cycloaddition of alkenes **3a-3d** with dienes such as cyclopentadiene (**4**) and isoprene (**5**) were, however, unsuccessful, except for derivative **3a**. The latter yielded adducts with diene **4** in an unexpectedly high






Table 1. Preparation of the β -sulfanyl substituted captodative olefins **10a-10h** from olefin **3a**.

Entry ^a	RSX	Base ^b	T (°C)	t (h)	10 (%) ^c
1	9a (R = Me, X = Na)	—	25	0.5	10a (85)
2	9b (R = Et, X = Na)	—	25	0.5	10b (90)
3	9c (R = <i>i</i> -Pr, X = Na)	—	25	0.5	10c (90)
4	9d (R = Et, X = H)	Et ₃ N	0-120 ^d	3.0	10b (78)
5	9e (R = <i>i</i> -Pr, X = H)	Et ₃ N	0-120 ^d	3.0	10c (81)
6	9f (R = <i>t</i> -Bu, X = H)	Et ₃ N	0-120 ^d	3.0	10d (82)
7	9g (R = Bn, X = H)	Et ₃ N	0-120 ^d	3.0	10e (90)
8	9h (R = C ₆ H ₄ <i>p</i> -Br, X = H)	Et ₃ N	0-120 ^d	3.0	10f (96)
9	9i (R = C ₆ H ₄ <i>p</i> -OMe, X = H)	Et ₃ N	0-120 ^d	3.0	10g (80)
10	9j (R = C ₆ H ₄ <i>p</i> -Me, X = H)	Et ₃ N	0-120 ^d	3.0	10h (81)

^a All under N₂ atmosphere, with 1.3 mol equiv. of RSX, in DMF as solvent.^b With 1.3 mol equiv. of Et₃N.^c After column and radial chromatography.^d The reaction was successively maintained at 0 °C for 1 h, 20 °C for 1 h, and at 120 °C for 1 h.**Fig. 1.****Fig. 2.**

exo stereoselectivity, and in comparable para/meta regioselectivity to that observed for olefin **1a** with diene **5** [6].

The high reactivity and selectivity of captodative olefins in cycloaddition reactions is rather unexpected, since the electron-releasing effect of the aryloxy group should decrease their reactivity in comparison with a dienophile or dipolarophile bearing only an electron-withdrawing group, such as methyl vinyl ketone (**6**) [10]. Structural and theoretical studies of olefin **1a** revealed that the delocalization of the oxygen lone pair of the electron-donor group toward the π -system was inhibited by conformational restrictions [11]. In addition, FMO calculations suggested a dominant effect of the acetyl electron-withdrawing group on the polarization of the olefin [11]. However, the high regioselectivity shown by olefins **1** in 1,3-dipolar additions toward nitrones and nitrile oxides was better rationalized by DFT/HSAB theory [3c], showing the relevance of the electron-donor group in controlling the interaction of the cycloaddends. Therefore, electronic and structural factors should be taken into account to explain the reactivity and regiochemistry observed in both Diels-Alder and 1,3-dipolar reactions.

It is then relevant to evaluate the perturbation of the *beta* substituent in olefins **3** on the electronic and structural properties of these molecules and, in particular, on the double bond. Accordingly, we hereby report the preparation of a large series of new captodative olefins β -substituted with a new amine, and alkyl and aryl thiols. MO calculations were also carried out to assess the effect of the third substituent on the FMO energies and coefficients.

Results and discussion

Synthesis of β -heterosubstituted captodative olefins

The β -amino substituted olefin **7a** was prepared by treating α -bromoalkene **3a** with amine **8a**, as an extension of the method used for the preparation of amino derivatives **3b** [6] (Fig. 1). Thus, addition of *N,N*-methylphenyl amine (**8a**), in methylene chloride at 10 °C for 30 min, satisfactorily led to the desired *Z*-substituted olefin **7a** in 62 % yield. The *Z* configuration of the double bond was established through NOE experiments. Irradiation of proton H-4 resulted in enhancement of the signal for the acetyl group.

In the previous report, the synthesis of β -thio (β -sulfanyl) olefins was limited to **3c** and **3d** by treatment of **3a** with the corresponding thiol in the presence of triethylamine in DMF at room temperature [6]. The use of some other alkyl mercaptanes and aryl thiophenols under these conditions was not as efficient. Therefore, three additional methodologies were investigated to improve the yields. The first method involved the addition of the sodium salt of the thiol (**9a-9c**) to **3a** (Fig. 2), leading to products **10a-10c** in high yields and under mild conditions (Table 1, entries 1-3). When thiols **9d-9g** were used, more severe conditions were applied in order to improve the preparation of olefins **10b-10e** (entries 4-7). The yields were high even for the bulky thiol **9f**. Under the same reaction conditions, substituted thiophenols **9h-9j** reacted with **3a** to give alkenes **10f-10h** also in good yields (entries 8-10).

The leaving group Y at the starting alkenes was also evaluated. Instead of the bromine atom of olefin **3a**, the dimeth-

Table 2. Preparation of the β -sulfanyl substituted captodative olefins **10b-10h** from olefin **7b**.

Entry ^a	RSH	10 (%) ^b
1	9d (R = Et)	10b (74)
2	9e (R = <i>i</i> -Pr)	10c (77)
3	9f (R = <i>t</i> -Bu)	10d (70)
4	9g (R = Bn)	10e (87)
5	9h (R = C ₆ H ₄ <i>p</i> -Br)	10f (91)
6	9i (R = C ₆ H ₄ <i>p</i> -OMe)	10g (93)
7	9j (R = C ₆ H ₄ <i>p</i> -Me)	10h (90)

^a All under N₂ atmosphere, with 1.3 mol equiv. of RSH and with 1.3 mol equiv. of Et₃N, in DMF as solvent. The reaction was successively maintained at 0 °C for 1 h, 120 °C for 1 h, and at 20 °C for 1 h.

^b After column and radial chromatography.

ylamino group was used, due to its known aptitude as a leaving group [12]. In these cases the optimum reaction conditions were found to be similar to those employed for olefin **3a**, that is, reacting olefin **7b** in the presence of thiols **9d-9j** and triethylamine as base, and heating from 0 °C up to 120 °C for 3 h (Table 2). The aliphatic thiols **9d-9g** provided compounds **10b-10e** in quite lower yields than those obtained when olefin **3a** was used (Table 2, entries 1-4 vs. Table 1, entries 4-7); however, the yields increased for the preparation of **10g-10h** when **7b** was treated with thiophenols **9i-9j** (Table 2, entries 6-7 vs. Table 1, entries 9-10).

For both olefins, **3a** and **7b**, there were no significant differences in reactivity or yields to furnish the β -substituted alkenes **10** with the *para* substituted thiophenols **9h-9j** (Table 1, entries 8-10 and Table 2, entries 5-7), considering that the temperature and the reaction time were comparable, regardless of the substituent.

It is interesting that sodium thiolates **9a-9c** reacted with olefin **3a** to give the 1,4-addition products **10a-10c** in good yields, instead of providing the hydrolysis products by addition to the *p*-nitrobenzoyloxy group, as observed when the corresponding alcohols were used. As expected, this behavior illustrates again the known greater softness of the sulphur atom with respect to the oxygen atom.

The *Z* configuration of the double bond is maintained in all of the new olefins **10a-10h**, as observed for the β -amino alkene **7a**, indicating that it is largely more stable than the *E*

configuration [6]. It is likely that the higher stability of the *Z* configuration is associated to destabilizing steric interactions found between the acetyl group and the *beta* substituent in the opposite *E* configuration. Such interactions may inhibit the effective conjugation of the enone moiety, and the delocalization of the heteroatom lone electron pair through the π system [6, 13].

The configuration of the double bond and the planar conformation of the enone moiety was confirmed by single crystal X-ray crystallography of olefin **10f** (Fig. 3). It shows that both conjugated moieties, the enone and *p*-nitrobenzoyl groups, are in a quasi-orthogonal conformation. This agrees with those structures obtained for other analogues [5, 6, 11], as well as in the *s-trans* conformation of the enone. Interestingly, the *p*-bromophenyl ring lies out of the plane formed by the enone π -system.

FMO calculations of β -heterosubstituted captodative olefins

Frontier molecular orbital (FMO) theory has proven to be a reliable model to predict reactivity and regioselectivity in Diels-Alder [14] and 1,3-dipolar cycloadditions [15]. It has been able to explain the behavior of substituted olefins as dienophiles in Diels-Alder additions. For instance, the rate increases when the dienophile bears electron-withdrawing groups, whereas it decreases with dienophiles bearing electron-donating groups [16].

FMO theory has also been useful in explaining the reactivity and regioselectivity of captodative olefins with dienes such as isoprene (**5**) [2a, 11], showing that the interaction was controlled by normal electron demand (NED), *i.e.* the HOMO-diene/LUMO-dienophile interaction was the energetically most favorable. Furthermore, these results were supported by experimental measurement of ionization energies (IEs) and vertical attachment energies (VAEs) of **1a**, whose relative values fit well with the calculated HOMO and LUMO energies, respectively [11].

Table 3. *Ab initio* HF/6-31G* Frontier Molecular Orbitals energies, and energy gaps (eV) of olefins **3a**, **7a**, **7b**, **10a**, **10f**, and **10g**, and isoprene (**5**).

Entry	Compound ^a	HOMO	LUMO	HOMO-LUMO ^b
1	1ac	-11.0460	2.4588	11.0781
2	3a	-10.4288	2.1015	10.7208
3	7a	-8.4516	2.9375	11.5568
4	7b	-8.8119	3.1767	11.7960
5	10a	-9.1412	2.4681	11.0874
6	10f	-9.2764	2.3162	10.9355
7	10g	-10.2014	2.1900	10.8093
8	5c	-8.6193	3.5337	

^a Of the non-planar *s-trans* conformation for olefins **1a**, **3a**, **7a**, **7b**, **10f**, and **10g**, and of the *s-cis* conformation for olefin **10a** and for the diene. ^b Energy gaps for the energetically more favorable HOMO-diene/LUMO-dienophile interaction. ^c Ref. [11].







Table 4. Comparison of X-ray selected bond distances (Å) (estimated standard deviations) of the crystal structure of **10f** with those of olefins **1a**, **3a**, and **7b**, and with the average lengths of bonds of the enone, enamine, vinyl ester and bromo vinyl systems.


Bond	10f	1a^a	3a^b	7b^b	Average Length ^c
O(4)-C(3)	1.208 (4)	1.215 (5)	1.20 (3)	1.231 (7)	1.222
C(2)-C(3)	1.470 (5)	1.490 (5)	1.40 (3)	1.41 (1)	1.462
C(1)-C(2)	1.320 (4)	1.306 (6)	1.42 (3)	1.366 (9)	1.340
C(1)-Y(5)	1.725 (4)		1.79 (3)	1.32 (1)	1.712 (S); 1.881 (Br); 1.358 (Nsp ²); 1.418 (Nsp ³)
C(2)-O(6)	1.412 (3)	1.398 (4)	1.41 (3)	1.423 (7)	1.353
O(6)-C(7)	1.355 (4)	1.353 (4)	1.38 (2)	1.354 (6)	1.359
C(7)-O(8)	1.195 (4)	1.200 (4)	1.21 (3)	1.188 (6)	1.201
C(7)-C(9)	1.486 (4)	1.488 (5)	1.50 (3)	1.481 (7)	1.481

^a Ref. [11]. ^b Ref. [6]. ^c Ref. [19]**Table 5.** *Ab initio* HF/6-31G* calculations of coefficients (C_i) of the frontier molecular orbitals for olefins **1a**, **3a**, **7a**, **7b**, **10a**, **10f**, and **10g**, and diene **5**.^a

1a, Y = H

3a, Y = Br

7a, Y = N(Me)Ph

7b, Y = NMe₂

Ar = C₆H₄*p*-NO₂

10a, Y = SMe

10f, Y = SC₆H₄*p*-Br

10g, Y = SC₆H₄*p*-OMe

Compd ^b	HOMO					ΔC_i^c	LUMO				
	<i>C</i> ₁	<i>C</i> ₂	<i>C</i> ₃	<i>C</i> ₄	<i>C</i> ₁		<i>C</i> ₂	<i>C</i> ₃	<i>C</i> ₄	ΔC_i^c	
3a	−0.2456	−0.2839	0.0095	0.1277	−0.0383	0.3188	−0.2499	−0.2783	0.2759	0.0689	
7a	−0.1192	−0.3110	−0.0215	0.1414	−0.1918	0.2701	−0.1547	−0.2314	0.2143	0.1154	
7b	−0.1711	−0.3702	−0.0190	0.1718	−0.1991	0.2799	−0.1433	−0.2564	0.2316	0.1366	
10a	−0.1848	−0.3135	−0.0038	0.1475	−0.1287	0.3189	−0.2200	−0.2807	0.2501	0.0989	
10f	−0.1625	−0.2694	−0.0019	0.1277	−0.1069	0.2872	−0.2073	−0.2506	0.2263	0.0799	
10g	−0.2082	−0.2400	0.0042	0.1174	−0.0318	0.2843	−0.2404	−0.2579	0.2574	0.0439	
1a^d	−0.3593	−0.3565	0.0236	0.1676	0.0028	0.2940	−0.2386	−0.2889	0.2800	0.0554	
5^d	0.3247	0.2523	−0.2180	−0.2857	0.0390	0.2591	−0.2236	−0.2306	0.2793	−0.0202	

^a These are the values of the $2p_z$ coefficients, the relative $2p_z'$ contributions and their ΔC_i are analogous.^b The FMOs of the non-planar *s-trans* conformation for olefins **1a**, **3a**, **7a**, **7b**, and **10g**, and *s-cis* for **5**, **10a**, and **10f**.^c Carbon 1 - carbon 2 for the olefins; carbon 1 - carbon 4 for the diene.^d Ref. [11]

Therefore, the electronic effect produced by the substituent in the *beta* position of the captodative olefins **3**, **7**, and **10** on their behavior in Diels-Alder reactions could be evaluated by calculating the FMO energies of these molecules, and correlating them with those of the corresponding FMOs of a diene such as **5**.

Table 3 summarizes the calculated (HF/6-31G*) FMO energies of bromo and amino alkenes **3a**, **7a**, and **7b**, as well as the thio alkenes **10a**, **10f**, and **10g**. The geometries were optimized with the same basis set, showing that the most stable geometry for **3a**, **7a**, and **7b**, corresponded to the *s-trans*

conformation of the enone moiety, and the non-planar conformation of the *p*-nitrobenzoyloxy group. However, only for thioether **10g**, the enone *s-trans* conformation was more stable, while olefins **10a** and **10f** were the exception, since the *s-cis* conformer for the enone conjugate system was slightly more stable (0.72 kcal/mol for **10a**, and 0.24 kcal/mol for **10f**), which is not in agreement with the X-ray structure of **10f**. Even though these energy differences are negligible, and, practically, both conformations are isoenergetic, the FMO energy values listed in Table 3 corresponded to those obtained for the most stable conformation in every molecule. The co-

efficient differences follow a similar trend in the other conformers.

As expected for the amino substituted olefins **7a** and **7b**, when an electron-releasing group is introduced into the double bond of the captodative olefin, both HOMO and LUMO should increase in energy (Table 3, entry 1 vs. entries 3 and 4). However, for the rest of the olefins, only the HOMO was energetically destabilized, since the LUMO energy of olefins **3a**, **10f**, and **10g** was lower than that of the unsubstituted olefin **1a**. It is likely that the effect of the heteroatom on the energy of the FMOs be the result of an interplay of different factors. The delocalization of the lone electron pair of bromine and sulphur toward the π -conjugated enone system will be less efficient than that of the nitrogen atom, due to the differences in electronic configuration. The inductive effect of these electronegative heteroatoms may increase the electron-withdrawing effect of the *beta* substituent on the electron density of the double bond [17].

The delocalization of the lone electron pair in enaminone **7b** was reflected in a shortening of the C(1)-N and C(2)-C(3) bonds, and by an increase of the C(3)=O(4) and C(1)=C(2) bond lengths [6], as observed for other analogues by X-ray diffraction [18]. Similarly, significant delocalization appears in the bromo olefin **3a**, since the C(2)-C(3) bond length is shorter and the C(1)=C(2) bond is longer than the corresponding bond lengths of **1a** and of the average values taken from X-ray data of similar functional groups [19] (Table 4). In contrast, the X-ray structure of **10f** (Fig. 2) provides bond distances similar to those for the β -unsubstituted olefin and for the non-delocalized enone [19] (Table 4). It is noteworthy that the bond distances between the atoms of the aryloxy group are not really perturbed by changes in the heteroatom at the *beta* position. This supports the idea of a non significant interaction between the lone pair of the oxygen atom of the electron donor group and the double bond, at least, in the crystalline state.

According to data in Table 3, one could anticipate a higher reactivity of olefins **3a**, **10f**, and **10g** with respect to **1a** in Diels-Alder cycloadditions with diene **5**, since the HOMO-diene / LUMO-dienophile energy gaps for the former dienophiles are smaller than that found for the latter. This expectation is only partially in agreement with the observed reactivity, since olefin **3a** added to **5** under thermal and Lewis acid catalytic conditions similar to those used for **1a** in shorter reaction times [6]. However, thioalkene **3d**, which is analogous to olefins **10**, failed to react even under catalysis. As predicted, the *beta* amino substituted alkene **7b** was less reactive than **1a**, being unable to react with cyclopentadiene (**12**) [6], which is considered a very reactive diene.

The unreliable prediction of the reactivity of olefins **10** by FMO theory may be attributed to steric hindrance [20], which would counterbalance the electronic effect, as suggested previously by establishing a correlation between the reaction rate and LUMO energies of olefins **1a** and **6** [11]. In addition, it has been found that the FMO model fails to account for the regioselectivity in Diels-Alder reactions with trisubstituted dienophiles or with phenylthiosubstituted dienes [21], or in

1,3-dipolar cycloadditions of nitrile oxides and nitrones with olefins such as **1a** [3c].

It is noteworthy that, under thermal and catalyzed conditions, the regioselectivity found in the Diels-Alder addition of dienophile **3a** with isoprene (**5**) was similar to that observed with the β -unsubstituted olefin **1a** [2a, 6]. From the FMO viewpoint, regioselectivity can be predicted on the basis of the atomic coefficient differences for the appropriate frontier orbital interaction: HOMO-diene/LUMO-dienophile, under NED control [14a, 14b]. It can be observed from Table 5 that, for olefin **3a**, the relative magnitude of the LUMO coefficient in the monosubstituted terminus of the double bond is larger than that of the geminally disubstituted carbon. In comparison with the LUMO of olefin **1a**, the difference in coefficients C_1 and C_2 for this olefin and bromo olefin **3a** are analogous, hence a comparable regioselectivity should be observed, which is, indeed, experimentally found. Thus, for diene **5**, which has the HOMO largest coefficient in carbon C-1, a preferred interaction with the largest coefficient on the double bond of the dienophile agrees with the *para* orientation as the major regioisomer.

Polarization of the π -system in the HOMO for the amino olefins **7a** and **7b**, and for the thio olefins **10** is towards the substituted captodative carbon of the double bond (Table 5). The larger difference in coefficients (ΔC_i) of these alkenes with respect to olefin **1a** reflects the electron-donor effect of the heteroatom in *beta* position. In contrast, the opposite polarization is found for the LUMO, where the larger coefficient is located in the monosubstituted carbon atom (C_1) of the double bond.

Conclusions

The stereoselective synthesis of new β -heteroatom substituted captodative olefins, including amino compound **7a**, and sulphur derivatives **10a-10h**, was feasible through three analogous routes. The common feature among them involved the replacement of a leaving group at the *beta* position of the olefin by the corresponding amino or thio compound. Both bromo and dimethylamino were efficient as the leaving groups in the starting activated substrates **3a** and **7b**, respectively. The former underwent nucleophilic attack of alkyl and aryl thiols or the sodium salt of some of them in good yields. The *Z* configuration of the double bond was established by NMR and X-ray crystallography. A comparison between bond distances by X-ray crystallography of different β -substituted and unsubstituted olefins seems to correlate with the delocalization effect of the heteroatom lone electron pair for the bromo and amino β -substituted olefins.

Ab initio calculations of FMOs of trisubstituted amino olefins **7a** and **7b** showed an increase of HOMO and LUMO energies with respect to the unsubstituted **1a**, as expected for the perturbation of the π -orbital by an electron-donating group. The presence of a bromine atom and alkyl and aryl thio groups in olefins **3a**, and **10a**, **10f**, and **10g**, however, pro-

duced an increase of the HOMO energy and a stabilization of the LUMO. Based on these results, the predicted reactivity of the Diels-Alder additions with diene **5** agrees with experiment only for alkenes **3a**, **7a** and **7b**. The regioselectivity observed for the cycloaddition of olefin **3a** is also explained by FMO theory. Therefore, these calculations clearly indicate a significant perturbation of the double bond of the captodative olefin by the third substituent in the *beta* position.

Experimental section

General. Melting points (uncorrected) were determined with an Electrothermal capillary melting point apparatus. IR spectra were recorded on a Perkin-Elmer 1600 spectrophotometer. ^1H and ^{13}C NMR spectra were obtained on a Varian Gemini-300 (300 MHz and 75.4 MHz), and Bruker DMX-500 (500 MHz and 125 MHz) instruments, with CDCl_3 as solvent and TMS as internal standard. The mass spectra (MS) were taken on a Hewlett-Packard 5971A spectrometer. X-Ray analyses were collected using Mo $\text{K}\alpha$ radiation (graphite crystal monochromator, $\lambda = 0.71073 \text{ \AA}$). Microanalyses were performed by M-H-W Laboratories (Phoenix, AZ). Analytical thin-layer chromatography was carried out using E. Merck silica gel 60 F_{254} coated 0.25 plates, visualizing by long- and short-wavelength UV lamp. All air moisture sensitive reactions were carried out under nitrogen using oven-dried glassware. DMF was freshly distilled and received on molecular sieves (4 \AA), and methylene chloride from calcium hydride, prior to use. Triethylamine was freshly distilled from NaOH. All other reagents were used without further purification. Compounds **3a** and **7b** were prepared as described previously [6].

(Z)-4-(N,N-Methylphenylamino)-3-(4-nitrobenzoyloxy)-3-buten-2-one (7a). To a solution of 1.0 g (3.18 mmol) of **3a** in CH_2Cl_2 (25 mL), at 10 $^\circ\text{C}$, 0.443 g (4.14 mmol) of **8a** were added, and the mixture was stirred for 30 min. The reaction mixture was diluted with CH_2Cl_2 (50 mL), and was washed with a cold 5 % aqueous solution of HCl ($2 \times 25 \text{ mL}$), and a cold saturated solution of NaCl ($2 \times 30 \text{ mL}$). The organic layer was dried (MgSO_4), and the solvent was evaporated under vacuum. The residue was successively purified by flash column chromatography on silica gel treated with 10 % of triethylamine (20 g, hexane/EtOAc, 90:10), and by radial chromatography (hexane/ CH_2Cl_2 , 90:10). The solid was recrystallized (hexane/ CH_2Cl_2 , 20:80) to give 0.67 g (62 %) of **7a** as pale brown crystals: R_f 0.14 (hexane/EtOAc, 8:2); mp 141–143 $^\circ\text{C}$; IR (CH_2Cl_2) 1741, 1621, 1591, 1529, 1495, 1350, 1316, 1244, 1096, 896 cm^{-1} ; ^1H NMR (300 MHz, CDCl_3) δ 2.24 (br s, 3H, CH_3CO), 3.45 (s, 3H, CH_3N), 6.99–7.04 (m, 1H, Ph-H), 7.13–7.16 (m, 2H, Ph-H), 7.22–7.27 (m, 2H, Ph-H), 7.48 (br s, 1H, HC=), 8.03–8.06 (m, 2H, Ar-H), 8.21–8.24 (m, 2H, Ar-H); ^{13}C NMR (75.4 MHz, CDCl_3) δ 24.3 (CH_3CO), 42.1 (CH_3N), 122.7, 123.2, 125.8, 126.8 (C-3), 129.2, 130.9 (C-4), 131.1, 134.3, 145.7, 150.6, 162.7 (ArCO_2), 187.0 (CH_3CO). Anal.

Calcd for $\text{C}_{18}\text{H}_{16}\text{N}_2\text{O}_5$: C, 63.52; H, 4.74; N, 8.23. Found: C, 62.75; H, 4.58; N, 7.71.

General Procedures for the Preparation of Olefins 10a–10h. Method A. To a mixture of 1.0 g (3.18 mmol) of **3a** and 1.31 mol equiv. of the sodium salt of the corresponding thiol **9a–9c**, at 0 $^\circ\text{C}$, anhydrous DMF (10 mL) was added, and the mixture was stirred for 30 min. The solution was concentrated under vacuum, the reaction crude was diluted with CH_2Cl_2 (50 mL), and was washed with a cold 5 % aqueous solution of HCl ($2 \times 25 \text{ mL}$), and a cold saturated solution of NaCl ($2 \times 30 \text{ mL}$). The organic layer was dried (MgSO_4), and the solvent was evaporated under vacuum. The residue was successively purified by column chromatography on silica gel treated with 10 % of triethylamine (30 g/1 g of crude, hexane/EtOAc, 90:10), and by radial chromatography (hexane/ CH_2Cl_2 , 80:20).

Method B. To a solution of 1.0 g (3.18 mmol) of **3a** in anhydrous DMF (20 mL) 1.31 mol equiv. of the corresponding thiol **9d–9j** were added at 20 $^\circ\text{C}$. The mixture was stirred and cooled down to 0 $^\circ\text{C}$, and a solution of 0.42 g (4.16 mmol) of triethylamine in anhydrous DMF (3 mL) was added dropwise. The mixture was maintained at the same temperature for 1 h, then warmed up to 120 $^\circ\text{C}$ for 1 h, and cooled down to 20 $^\circ\text{C}$ for 1 h. The solution was concentrated under vacuum, the reaction crude was diluted with CH_2Cl_2 (50 mL), and washed with a cold 5 % aqueous solution of HCl ($2 \times 25 \text{ mL}$), and a cold saturated solution of NaCl ($2 \times 30 \text{ mL}$). The organic layer was dried (MgSO_4), and the solvent was evaporated under vacuum. The residue was successively purified by column chromatography on silica gel treated with 10 % of triethylamine (30 g/1 g of crude, hexane/EtOAc, 90:10), and by radial chromatography (hexane/ CH_2Cl_2 , 90:10). For the solid products, the recrystallization was carried out from hexane/ CH_2Cl_2 , 10:90.

Method C. To a solution of 1.0 g (3.59 mmol) of **7b** in anhydrous DMF (25 mL) 1.3 mol equiv. of the corresponding thiol **9d–9j** were added at 20 $^\circ\text{C}$. The mixture was stirred and cooled down to 0 $^\circ\text{C}$, and a solution of 0.472 g (4.67 mmol) of triethylamine in anhydrous DMF (3 mL) was added dropwise. The mixture was maintained at the same temperature for 1 h, then warmed up to 120 $^\circ\text{C}$ for 1 h, and cooled down to 20 $^\circ\text{C}$ for 1 h. The solution was concentrated under vacuum, the reaction crude was diluted with CH_2Cl_2 (50 mL), and was washed with a cold 5 % aqueous solution of HCl ($2 \times 25 \text{ mL}$), and a cold saturated solution of NaCl ($2 \times 30 \text{ mL}$). The organic layer was dried (MgSO_4), and the solvent was evaporated under vacuum. The residue was successively purified by column chromatography on silica gel treated with 10 % of triethylamine (30 g/1 g of crude, hexane/EtOAc, 80:20), and by radial chromatography (hexane/ CH_2Cl_2 , 80:20). For the solid products, the recrystallization was carried out from hexane/ CH_2Cl_2 , 10:90.

(Z)-4-Methylsulfanyl-3-(p-nitrobenzoyloxy)-3-buten-2-one (10a). According to method A with 0.291 g (4.16 mmol) of **9a**, afforded 0.76 g (85 %) of **10a** as a pale yellow oil: R_f 0.21 (hexane/EtOAc, 8:2); IR (CH_2Cl_2) 1746, 1675, 1589, 1530, 1347, 1248, 1093, 895 cm^{-1} ; ^1H NMR (300 MHz, CDCl_3) δ 2.34 (s, 3H, CH_3CO), 2.50 (s, 3H, CH_3S), 7.39 (s, 1H, HC=), 8.32-8.34 (m, 4H, Ar-H); ^{13}C NMR (75.4 MHz, CDCl_3) δ 17.3 (CH_3S), 24.7 (CH_3CO), 123.7, 131.4, 134.1, 136.9 (C-4), 142.3 (C-3), 151.1, 161.7 (ArCO_2), 187.1 (CH_3CO); MS (70 eV) 281 (M^+ , 4), 150 (100), 134 (22), 120 (15), 104 (22), 92 (11), 76 (14).

(Z)-4-Ethylsulfanyl-3-(p-nitrobenzoyloxy)-3-buten-2-one (10b). According to method A with 0.349 g (4.16 mmol) of **9b**, afforded 0.84 g (90 %) of **10b** as a pale yellow oil. According to method B with 0.258 g (4.16 mmol) of **9d**, gave 0.73 g (78 %) of **10b**. According to method C with 0.289 g (4.67 mmol) of **9d**, furnished 0.83 g (74%) of **10b**: R_f 0.22 (hexane/EtOAc, 8:2); IR (CH_2Cl_2) 1746, 1675, 1588, 1529, 1423, 1348, 1275, 1246, 1217, 1049, 1013 cm^{-1} ; ^1H NMR (300 MHz, CDCl_3) δ 1.43 (t, $J = 7.1$ Hz, 3H, $\text{CH}_3\text{CH}_2\text{S}$), 2.35 (s, 3H, CH_3CO), 2.92 (q, $J = 7.1$ Hz, 3H, $\text{CH}_3\text{CH}_2\text{S}$), 7.49 (s, 1H, HC=), 8.45-8.47 (m, 4H, Ar-H); ^{13}C NMR (75.4 MHz, CDCl_3) δ 15.4 ($\text{CH}_3\text{CH}_2\text{S}$), 24.7 (CH_3CO), 28.4 ($\text{CH}_3\text{CH}_2\text{S}$), 123.6, 131.4, 134.1, 135.1 (C-4), 142.5 (C-3), 151.0, 161.7 (ArCO_2), 187.1 (CH_3CO). Anal. Calcd for $\text{C}_{13}\text{H}_{13}\text{NO}_5\text{S}$: C, 52.87; H, 4.44. Found: C, 52.80; H, 4.68.

(Z)-4-Isopropylsulfanyl-3-(p-nitrobenzoyloxy)-3-buten-2-one (10c). According to method A with 0.408 g (4.16 mmol) of **9c**, afforded 0.88 g (90 %) of **10c** as a pale yellow oil. According to method B with 0.358 g (4.16 mmol) of **9e**, gave 0.80 g (81 %) of **10c**. According to method C with 0.402 g (4.67 mmol) of **9e**, furnished 0.90 g (77 %) of **10c**: R_f 0.30 (hexane/EtOAc, 8:2); IR (CH_2Cl_2) 1746, 1674, 1587, 1530, 1453, 1427, 1348, 1314, 1218, 1157, 1092, 1015, 958, 898, 870, 847 cm^{-1} ; ^1H NMR (300 MHz, CDCl_3) δ 1.38 (d, $J = 7.3$ Hz, 6H, $(\text{CH}_3)_2\text{CHS}$), 2.35 (s, 3H, CH_3CO), 3.33 (sept, $J = 7.3$ Hz, 3H, $(\text{CH}_3)_2\text{CHS}$), 7.50 (s, 1H, HC=), 8.38-8.36 (m, 4H, Ar-H); ^{13}C NMR (75.4 MHz, CDCl_3) δ 23.4 ($(\text{CH}_3)_2\text{CHS}$), 24.5 (CH_3CO), 38.4 ($(\text{CH}_3)_2\text{CHS}$), 123.4, 131.1, 133.9, 134.1 (C-4), 141.9 (C-3), 150.7, 161.5 (ArCO_2), 187.0 (CH_3CO). Anal. Calcd for $\text{C}_{14}\text{H}_{15}\text{NO}_5\text{S}$: C, 54.36; H, 4.89. Found: C, 54.30; H, 5.00.

(Z)-4-tert-Butylsulfanyl-3-(p-nitrobenzoyloxy)-3-buten-2-one (10d). According to method B with 0.375 g (4.13 mmol) of **9f**, afforded 0.84 g (82 %) of **10d** as a pale yellow oil. According to method C with 0.514 g (4.67 mmol) of **9f**, furnished 0.86 g (70%) of **10d**: R_f 0.26 (hexane/EtOAc, 8:2); IR (CH_2Cl_2) 1744, 1673, 1584, 1529, 1349, 1248, 1093, 769, 751 cm^{-1} ; ^1H NMR (300 MHz, CDCl_3) δ 1.48 (s, 9H, $(\text{CH}_3)_3\text{CS}$), 2.35 (s, 3H, CH_3CO), 7.60 (s, 1H, HC=), 8.33 (br s, 4H, Ar-H); ^{13}C NMR (75.4 MHz, CDCl_3) δ 24.8 (CH_3CO), 31.1 ($(\text{CH}_3)_3\text{CS}$), 45.7 ($(\text{CH}_3)_3\text{CS}$), 123.7, 131.4, 131.6 (C-4), 134.2, 142.3 (C-3), 151.0, 162.4 (ArCO_2), 198.8 (CH_3CO).

Anal. Calcd for $\text{C}_{15}\text{H}_{17}\text{NO}_5\text{S}$: C, 55.71; H, 5.30. Found: C, 55.50; H, 5.30.

(Z)-4-Benzylsulfanyl-3-(p-nitrobenzoyloxy)-3-buten-2-one (10e). According to method B with 0.516 g (4.13 mmol) of **9g**, afforded 1.02 g (90 %) of **10e** as a pale yellow oil. According to method C with 0.579 g (4.67 mmol) of **9g**, furnished 1.18 g (87 %) of **10e**: R_f 0.28 (hexane / EtOAc, 8:2); IR (CH_2Cl_2) 1740, 1677, 1606, 1530, 1348, 1248, 1093 cm^{-1} ; ^1H NMR (300 MHz, CDCl_3) δ 2.25 (s, 3H, CH_3CO), 4.08 (s, 2H, CH_2Ph), 7.27-7.46 (m, 6H, HC= , Ph-H), 8.28-8.39 (m, 4H, Ar-H); ^{13}C NMR (75.4 MHz, CDCl_3) δ 24.7 (CH_3CO), 38.0 (CH_2Ph), 123.7, 128.2, 129.0, 129.1, 131.4, 134.0, 134.2 (C-4), 136.7, 142.4 (C-3), 150.9, 161.7 (ArCO_2), 187.1 (CH_3CO).

(Z)-4-(4-Bromophenylsulfanyl)-3-(4-nitrobenzoyloxy)-3-buten-2-one (10f). According to method B with 0.786 g (4.16 mmol) of **9h**, afforded 1.29 g (96 %) of **10f** as pale yellow crystals. According to method C with 0.883 g (4.67 mmol) of **9h**, furnished 1.45 g (91 %) of **10f**: mp 174-175 $^\circ\text{C}$; R_f 0.16 (hexane/EtOAc, 8:2); IR (CH_2Cl_2) 1747, 1679, 1531, 1419, 1348, 1243, 1090, 1008 cm^{-1} ; ^1H NMR (300 MHz, CDCl_3) δ 2.35 (s, 3H, CH_3CO), 7.36-7.40 (m, 2H, Ar-H), 7.46 (s, 1H, HC=), 7.51-7.55 (m, 2H, Ar-H), 8.30-8.38 (m, 4H, Ar-H); ^{13}C NMR (75.4 MHz, CDCl_3) δ 24.9 (CH_3CO), 123.6, 123.7, 130.8, 131.5, 132.9, 133.1, 133.7, 134.1 (C-4), 142.6 (C-3), 151.2, 161.7 (ArCO_2), 187.3 (CH_3CO). Anal. Calcd for $\text{C}_{17}\text{H}_{12}\text{BrNO}_5\text{S}$: C, 48.36; H, 2.86; N 3.32; S, 7.58. Found: C, 48.04; H, 2.80; N, 3.29; S, 7.41.

(Z)-4-(4-Methoxyphenylsulfanyl)-3-(4-nitrobenzoyloxy)-3-buten-2-one (10g). According to method B with 0.582 g (4.16 mmol) of **9i**, afforded 0.95 g (80 %) of **10g** as pale yellow crystals. According to method C with 0.654 g (4.67 mmol) of **9i**, furnished 1.25 g (93 %) of **10g**: mp 122-123 $^\circ\text{C}$; R_f 0.20 (hexane/EtOAc, 8:2); IR (CH_2Cl_2) 1743, 1674, 1592, 1546, 1531, 1492, 1421, 1346, 1264, 1093 cm^{-1} ; ^1H NMR (300 MHz, CDCl_3) δ 2.33 (s, 3H, CH_3CO), 3.82 (s, 3H, MeO), 6.88-6.94 (m, 2H, Ar-H), 7.44-7.49 (m, 2H, Ar-H), 7.45 (s, 1H, HC=), 8.34 (s, 4H, Ar-H); ^{13}C NMR (75.4 MHz, CDCl_3) δ 24.8 (CH_3CO), 55.4 (CH_3O), 115.3, 122.0, 123.6, 131.5, 134.0, 134.2, 137.2 (C-4), 141.5 (C-3), 150.9, 160.7, 161.7 (ArCO_2), 187.2 (CH_3CO). Anal. Calcd for $\text{C}_{18}\text{H}_{15}\text{NO}_6\text{S}$: C, 57.90; H, 4.05; N, 3.75; S, 8.59. Found: C, 58.08; H, 4.12; N 3.74; S, 8.70.

(Z)-4-(4-Methylphenylsulfanyl)-3-(4-nitrobenzoyloxy)-3-buten-2-one (10h). According to method B with 0.516 g (4.16 mmol) of **9j**, afforded 0.92 g (81 %) of **10h** as pale yellow crystals. According to method C with 0.58 g (4.67 mmol) of **9j**, furnished 1.15 g (90 %) of **10h**: mp 113-115 $^\circ\text{C}$; R_f 0.22 (hexane/EtOAc, 8:2); IR (CH_2Cl_2) 1747, 1677, 1589, 1531, 1348, 1091, 897, 847, 807 cm^{-1} ; ^1H NMR (500 MHz, CDCl_3) δ 2.33 (s, 3H, CH_3CO), 2.37 (s, 3H, MeAr), 7.20-7.22 (m, 2H, Ar-H), 7.39-7.41 (m, 2H, Ar-H), 7.50 (s, 1H, HC=), 8.34 (s, 4H, Ar-H); ^{13}C NMR (125 MHz, CDCl_3) δ 21.1 (CH_3Ar), 24.8 (CH_3CO), 123.7, 128.2, 130.5, 131.5, 131.9, 134.0, 136.3

(C-4), 138.5, 141.9 (C-3), 151.0, 161.7 (ArCO₂), 187.3 (CH₃CO). Anal. Calcd for C₁₈H₁₅NO₅S: C, 60.49; H, 4.23. Found: C, 60.27; H, 4.50.

Single-Crystal X-Ray Crystallography [22]. Olefin **10f** was obtained as pale yellow crystals. These were mounted in glass fibers. Crystallographic measurements were performed on a Siemens P4 diffractometer with Mo K α radiation ($\lambda = 0.7107$ Å; graphite monochromator) at room temperature. Two standard reflections were monitored periodically; they showed no change during data collection. Unit cell parameters were obtained from least-squares refinement of 26 reflections in the range $2 < 2\theta < 20^\circ$. Intensities were corrected for Lorentz and polarization effects. No absorption correction was applied. Anisotropic temperature factors were introduced for all non-hydrogen atoms. Hydrogen atoms were placed in idealized positions and their atomic coordinates refined. Unit weights were used in the refinement. The structure was solved using SHELXTL on a personal computer [23]. Data of **10f**: Formula: C₁₇H₁₂BrNO₅S; molecular weight: 422.25; cryst. syst.: monoclinic; space group: P2₁/c; unit cell parameters: *a*, 6.4121 (9), *b*, 12.8041 (11), *c*, 20.998 (2) (Å); α , 90, β , 94.884 (9), γ , 90 (deg); temp. (°K): 293 (2); Z: 4; No. of reflections collected: 4641; no. of independent reflections: 3335; no. of observed reflections: 3290; *R*: 0.0429; GOF: 1.012.

Calculations. The ab initio SCF/HF calculations were carried out with the 6-31G* basis sets using Gaussian 94 [24] and MacSpartan [24]. Geometries were fully optimized by the AM1 semiempirical method [25] and these were employed as starting point for optimization, at the 6-31G* level.

Acknowledgments

We thank Fernando Labarrios for his help in spectrometric measurements. J.T. would like to acknowledge DEPI/IPN (Grants 921769 and 200410) and CONACyT (Grants 1570P and 32273-E) for financial support. H.A.J.-V. thanks CONACyT (Grant 3251P) for financial support. J.M. and R.H. are grateful to CONACyT (Grants 86038 and 91187) for graduate fellowships, to PIFI-IPN program for a scholarship, and to the Ludwig K. Hellweg Foundation for a partial scholarship. J.L. is grateful to Secretaría de Relaciones Exteriores, México (Grant DAC-III 811.5/(510)/137) for a research fellowship. J.T. and H.A.J.-V. are fellows of the EDD/IPN and COFAA/IPN programs.

References and notes

- (a) Viehe, H. G.; Janousek, Z.; Merényi, R.; Stella, L. *Acc. Chem. Res.* **1985**, *18*, 148-154. (b) Cativiela, C.; Fraile, J. M.; García, J. I.; Mayoral, J. A.; Pires, E.; Royo, A. J.; Figueras, F.; de Ménorval, L. C. *Tetrahedron* **1993**, *49*, 4073-4084. (c) Boucher, J.-L.; Stella, L. *Tetrahedron* **1986**, *42*, 3871-3885. (d) Seerden, J.-P. G.; Scheeren, H. W. *Tetrahedron Lett.* **1993**, *34*, 2669-2672. (e) Döpp, D.; Libera, H. *Tetrahedron Lett.* **1983**, *24*, 885-888. (f) Rulev, A. Y. *Russ. Chem. Rev.* **2002**, *71*, 195-221. (g) Seneci, P.; Leger, I.; Souchet, M.; Nadler, G. *Tetrahedron* **1997**, *53*, 17097-17114. (h) Moody, C. J.; Hughes, R. A.; Thompson, S. P.; Alcaraz, L. *Chem. Commun.* **2002**, 1760-1761. (i) Kozmin, S. A.; Iwama, T.; Huang, Y.; Rawal, V. H. *J. Am. Chem. Soc.* **2002**, *104*, 4628-4641. (j) Ferreira, P. M. T.; Maia, H. L. S.; Monteiro, L. S. *Tetrahedron Lett.* **2002**, *43*, 4491-4493. (k) Ferreira, P. M. T.; Maia, H. L. S.; Monteiro, L. S. *Tetrahedron Lett.* **2002**, *43*, 4495-4497. (l) Huang, C.-G.; Chang, B.-R.; Chang, N.-C. *Tetrahedron Lett.* **2002**, *43*, 2721-2723.
- (a) Reyes, A.; Aguilar, R.; Muñoz, A. H.; Zwick, J.-C.; Rubio, M.; Escobar, J.-L.; Soriano, M.; Toscano, R.; Tamariz, J. *J. Org. Chem.* **1990**, *55*, 1024-1034. (b) Aguilar, R.; Reyes, A.; Tamariz, J.; Birbaum, J.-L. *Tetrahedron Lett.* **1987**, *28*, 865-868. (c) García de Alba, O.; Chanona, J.; Delgado, F.; Zepeda, G.; Labarrios, F.; Bates, R. W.; Bott, S.; Juaristi, E.; Tamariz, J. *Anal. Quím., Int. Ed.* **1996**, *92*, 108-117.
- (a) Nagarajan, A.; Zepeda, G.; Tamariz, J. *Tetrahedron Lett.* **1996**, *37*, 6835-6838. (b) Jiménez, R.; Pérez, L.; Tamariz, J.; Salgado, H. *Heterocycles* **1993**, *35*, 591-598. (c) Herrera, R.; Nagarajan, A.; Morales, M. A.; Méndez, F.; Jiménez-Vázquez, H. A.; Zepeda, L. G.; Tamariz, J. *J. Org. Chem.* **2001**, *66*, 1252-1263.
- (a) Andrade, R. M.; Muñoz, A. H.; Tamariz, J. *Synth. Commun.* **1992**, *22*, 1603-1609. (b) Orduña, A.; Zepeda, G.; Tamariz, J. *Synthesis* **1993**, 375-377. (c) Dienes, Z.; Vogel, P. *J. Org. Chem.* **1996**, *61*, 6958-6970. (d) Ochoa, M. E.; Arias, A. S.; Aguilar, R.; Delgado, F.; Tamariz, J. *Tetrahedron* **1999**, *55*, 14535-14546. (e) Aguilar, R.; Reyes, A.; Orduña, A.; Zepeda, G.; Bates, R. W.; Tamariz, J. *Rev. Soc. Quím. Méx.* **2000**, *44*, 91-96.
- Herrera, R.; Jiménez-Vázquez, H. A.; Modelli, A.; Jones, D.; Söderberg, B. C.; Tamariz, J. *Eur. J. Org. Chem.* **2001**, 4657-4669.
- Peralta, J.; Bullock, J. P.; Bates, R. W.; Bott, S.; Zepeda, G.; Tamariz, J. *Tetrahedron* **1995**, *51*, 3979-3996.
- (a) Rappoport, Z., Ed., *The Chemistry of Enamines*. Wiley, Chichester, **1994**. (b) Greenhill, J. V. *Chem. Soc. Rev.* **1977**, *6*, 277-294. (c) Granik, V. G. *Russ. Chem. Rev.* **1984**, *53*, 383-404.
- (a) Scott, K. R.; Edafiogho, I. O.; Richardson, E. L.; Farrar, V. A.; Moore, J. A.; Tietz, E. I.; Hinko, C. N.; Chang, H.; El-Assadi, A.; Nicholson, J. M. *J. Med. Chem.* **1993**, *36*, 1947-1955. (b) Kesten, S. J.; Degnan, M. J.; Hung, J.; McNamara, D. J.; Ortwine, D. F.; Uhlendorf, S. E.; Werbel, L. M. *J. Med. Chem.* **1992**, *35*, 3429-3447.
- For some recent examples: (a) Bhathia, G. S.; Lowe, R. F.; Pritchard, R. G.; Stoodley, R. *J. Chem. Commun.* **1997**, 1981-1982. (b) Abbiati, G.; Clerici, F.; Gelmi, M. L.; Gambini, A.; Pilati, T. *J. Org. Chem.* **2001**, *66*, 6299-6304. (c) Clerici, F.; Gelmi, M. L.; Pocar, D.; Pilati, T. *Tetrahedron: Asymmetry* **2001**, *12*, 2663-2669. (d) Buñuel, E.; Gil, A. M.; Díaz-de-Villegas, M. D.; Cativiela, C. *Tetrahedron* **2001**, *57*, 6417-6427. (e) Clerici, F.; Gelmi, M. L.; Gambini, A.; Nava, D. *Tetrahedron* **2001**, *57*, 6429-6438. (f) Maekawa, K.; Igarashi, T.; Kubo, K.; Sakurai, T. *Tetrahedron* **2001**, *57*, 5515-5526. (g) Caine, D. *Tetrahedron* **2001**, *57*, 2643-2684. (h) Stanovnik, B.; Svete, J. *Synlett* **2000**, 1077-1091. (i) Yonehara, K.; Ohe, K.; Uemura, S. *J. Org. Chem.* **1999**, *64*, 9381-9385.
- (a) Sustmann, R. *Tetrahedron Lett.* **1971**, 2721-2724. (b) Houk, K. N.; Sims, J.; Watts, C. R.; Luskus, L. J. *J. Am. Chem. Soc.* **1973**, *95*, 7301-7315.
- Jiménez-Vázquez, H. A.; Ochoa, M. E.; Zepeda, G.; Modelli, A.; Jones, D.; Mendoza, J. A.; Tamariz, J. *J. Phys. Chem. A* **1997**, *101*, 10082-10089.
- (a) Guseinov, F. I.; Yudina, N. A.; Burangulova, R. N.; Ryzhikova, T. Ya.; Valliullina, R. Zh. *Chem. Heterocyclic Comp.* **2002**, *38*, 496-497. (b) Toplak, R.; Svete, J.; Stanovnik, B.; Grdadolnik, S. G. *J. Heterocyclic Chem.* **1999**, *36*, 225-235. (c) Pizzioli, L.; Ornik, B.; Svete, J.; Stanovnik, B. *Helv. Chim. Acta* **1998**, *81*, 231-235.

13. Zhuo, J.-C. *Magn. Reson. Chem.* **1998**, 36, 565-572.
14. (a) Fleming, I. *Frontier Orbitals and Organic Chemical Reactions*. John Wiley & Sons, Chichester, **1976**. (b) Houk, K. N. *Acc. Chem. Res.* **1975**, 8, 361-369. (c) Eisenstein, O.; Lefour, J. M.; Anh, N. T.; Hudson, R. F. *Tetrahedron* **1977**, 33, 523-531. (d) Spino, C.; Pesant, M. Dory, Y. *Angew. Chem., Int. Ed.* **1998**, 37, 3262-3265.
15. (a) Carruthers, W. *Cycloaddition Reactions in Organic Synthesis*. Pergamon Press, Oxford, **1990**, 269-331. (b) Houk, K. N.; Chang, Y.-M.; Strozier, R. W.; Caramella, P. *Heterocycles* **1977**, 7, 793-799. (c) Huisgen, R., in: *1,3-Dipolar Cycloaddition Chemistry*, Vol. 1, Padwa, A., Ed., Wiley-Interscience, New York, **1984**. (d) Jarosková, L.; Fisera, L.; Matejková, I.; Ertl, P.; Prónayová, N. *Monatsh. Chem.* **1994**, 125, 1413-1425. (e) Houk, K. N.; Sims, J.; Duke, Jr., R. E.; Strozier, R. W.; George, J. K. *J. Am. Chem. Soc.* **1973**, 95, 7287-7301.
16. (a) Sauer, J.; Sustmann, R. *Angew. Chem., Int. Ed. Engl.* **1980**, 19, 779-807. (b) Anh, N. T.; Canadell, E.; Eisenstein, O. *Tetrahedron* **1978**, 34, 2283-2288.
17. (a) Exner, O. *J. Phys. Org. Chem.* **1999**, 12, 265-274. (b) Charton, M. *J. Phys. Org. Chem.* **1999**, 12, 275-282. (c) Galkin, V. I. *J. Phys. Org. Chem.* **1999**, 12, 283-288.
18. Zhuo, J.-C.; Schenk, K. *Helv. Chim. Acta* **1997**, 80, 2137-2147.
19. Allen, F. H.; Kennard, E.; Watson, D. G.; Brammer, L.; Orpen, A. G.; Taylor, R. *J. Chem. Soc., Perkin Trans. 2* **1987**, S1-S19.
20. (a) Fox, M. A.; Cardona, R.; Kiwiet, N. J. *J. Org. Chem.* **1987**, 52, 1469-1474. (b) Bachler, V.; Mark, F. *Theoret. Chim. Acta* **1976**, 43, 121-135. (c) Tripathy, R.; Franck, R. W.; Onan, K. D. *J. Am. Chem. Soc.* **1988**, 110, 3257-3262. (d) Padwa, A.; Kline, D. N.; Koehler, K. F.; Matzinger, M.; Venkatramanan, M. K. *J. Org. Chem.* **1987**, 52, 3909-3917.
21. Kahn, S. D.; Pau, C. F.; Overman, L. E.; Hehre, W. J. *J. Am. Chem. Soc.* **1986**, 108, 7381-7396.
22. The authors have deposited the atomic coordinates for this structure with the Cambridge Crystallographic Data Centre. The coordinates can be obtained, on request, from the Director Cambridge Crystallographic Data Centre, 12 Union Road, Cambridge, CB2 1EZ, UK.
23. SHELXTL, v. 5.03, Siemens Energy & Automation, Germany, **1995**.
24. Calculated with Gaussian 94, Revision E.2: Frisch, M. J.; Trucks, G. W.; Schlegel, H. B.; Gill, P. M. W.; Johnson, B. G.; Robb, M. A.; Cheeseman, J. R.; Keith, T.; Petersson, G. A.; Montgomery, J. A.; Raghavachari, K.; Al-Laham, M. A.; Zakrzewski, V. G.; Ortiz, J. V.; Foresman, J. B.; Cioslowski, J.; Stefanov, B. B.; Nanayakkara, A.; Challacombe, M.; Peng, C. Y.; Ayala, P. Y.; Chen, W.; Wong, M. W.; Andres, J. L.; Replogle, E. S.; Gomperts, R.; Martin, R. L.; Fox, D. J.; Binkley, J. S.; Defrees, D. J.; Baker, J.; Stewart, J. P.; Head-Gordon, M.; Gonzalez, C.; Pople, J. A. Gaussian, Inc., Pittsburgh, PA, **1995**; and MacSpartan, v. 1.0, WaveFunction Inc., 18401 VonKarman, Suite 370, Irvine, CA 92715.
25. Dewar, M. J. S.; Zoebisch, E. G.; Healy, E. F.; Stewart, J. J. P. *J. Am. Chem. Soc.* **1985**, 107, 3902-3909.