



Journal of the Mexican Chemical Society

ISSN: 1870-249X

editor.jmcs@gmail.com

Sociedad Química de México

México

Ortega, María C.; López Cortés, José G.; Toscano, Rubén A.; Alvarez Toledano, Cecilio  
Electronic Effects on the Reactivity of Cross-Conjugated Carbonyl Systems with  $\text{Fe}_2(\text{CO})_9$

Journal of the Mexican Chemical Society, vol. 49, núm. 2, 2005, pp. 124-133

Sociedad Química de México

Distrito Federal, México

Available in: <http://www.redalyc.org/articulo.oa?id=47549209>

- How to cite
- Complete issue
- More information about this article
- Journal's homepage in redalyc.org

redalyc.org

Scientific Information System

Network of Scientific Journals from Latin America, the Caribbean, Spain and Portugal

Non-profit academic project, developed under the open access initiative

## Electronic Effects on the Reactivity of Cross-Conjugated Carbonyl Systems with $\text{Fe}_2(\text{CO})_9$

María C. Ortega-Alfaro<sup>\*,a</sup>, José G. López-Cortés<sup>b</sup>, Rubén A. Toscano<sup>b</sup> and Cecilio Alvarez-Toledano<sup>b</sup>

<sup>a</sup> Facultad de Química, Ed. A, Lab. 4-D, UNAM, Circuito Exterior, Ciudad Universitaria, Coyoacán 04510, México D.F., México

<sup>b</sup> Instituto de Química-UNAM, Circuito Exterior, Ciudad Universitaria, Coyoacán 04510, México D.F., México

Neste trabalho foram estudados os efeitos eletrônicos induzidos por p-substituintes no anel fenílico de diversos compostos carbonílicos conjugados [4-*p*-R-fenil-1-(1,3-ditiolano-2-ilideno)-3-*E*-buteno-2-ona, onde R=NO<sub>2</sub>, Cl, Br, OMe, NEt<sub>2</sub> (**2-6**), e sua reatividade frente a  $\text{Fe}_2(\text{CO})_9$ , usando o diagrama de Hammett. A estrutura dos compostos (**3** e **4b**) foi totalmente estabelecida por análise de difração de raio-X.


The electronic effects induced by *p*-substituents on the phenyl ring of several novel cross-conjugated carbonyl compounds [4-*p*-R-phenyl-1-(1,3-dithiolane-2-ylidene)-3-*E*-butene-2-one, where R= NO<sub>2</sub>, Cl, Br, OMe, NEt<sub>2</sub> (**2-6**),] and their reactivity towards  $\text{Fe}_2(\text{CO})_9$  have been studied using a Hammett plot. The structure of two compounds (**3** and **4b**) was fully established by X-ray diffraction analysis.

**Keywords:** iron(0) complexes, C-S bond activation, reactivity, Hammett plot, X-ray diffraction

## Introduction

The  $\alpha,\beta$ -unsaturated carbonyl compounds react with  $\text{Fe}_2(\text{CO})_9$  forming  $\eta^2$ -[ $\text{Fe}(\text{CO})_4$ ] and the most thermodynamically stable  $\eta^4$ -[ $\text{Fe}(\text{CO})_3$ ] complexes,<sup>1</sup> which have shown different applications such as transfer agent of  $\text{Fe}(\text{CO})_3$ ,<sup>2</sup> protecting group,<sup>3</sup> stereodirector group,<sup>4</sup> etc. Recently, we have described how reactivity changes drastically when heteroatoms (nitrogen and sulfur) in  $\beta$ -position are located on the system leading to novel an

unexpected dinuclear complexes.<sup>5</sup> In this subject, we have reported that the reactivity of cross-conjugated carbonyl systems  $\beta$ -positioned by sulfur atoms towards  $\text{Fe}_2(\text{CO})_9$ , principally produce partial and total C-S bond activation in addition to the formation of  $\eta^4$ -[ $\text{Fe}(\text{CO})_3$ ] complexes (Scheme 1).<sup>6</sup> As a part of our constant interest in the understanding of the coordination patterns of this kind of systems, we herein report the study of electronic effects induced by *p*-substituents on the phenyl ring on the mechanism of reaction of cross-conjugated carbonyls




Scheme 1.



## Results


The structural arrangement for **3** was fully established by a single-crystal X-ray diffraction analysis (Figure 1). The PhCH=CHCO moiety presents a *s-trans* conformation. The ylidenebutenone backbone of the ligand is quasi-planar with the phenyl ring tilted 18.6° out of main plane. The bond distance C1-C11 in comparison with C3-C4 bond is longer [0.069 Å] (Table1) and the distance [S1---O1, 2.705Å] is shorter than the  $\Sigma$  of Van der Waals radii. These features indicate that an interaction between the sulfur and oxygen atoms through the conjugated system is present, in agreement with the obtained results from IR technique and with described in literature.<sup>6,7</sup> The 1,3-dithiolane ring adopts a half-chair conformation with C<sub>2</sub> axe over C11. Additionally, this moiety presents disorder generating two conformers in 46/54 ratio, only the major contributors are shown in Figure 1.




**Table 1.** Selected bond lengths (Å) and angles (°) for compound

S1-C11	1.735(3)	S2-C11	1.739(3)
O1-C2	1.215(3)	C1-C11	1.347(4)
C3-C4	1.278(4)	C4-C5	1.506(4)
S1-C13	1.882(12)	S2-C12	1.814(10)
S1-C13B	1.789(11)	S2-C12B	1.785(9)
C1-C2	1.455(4)	C2-C3	1.491(4)
C11-S1-C13	94.4(4)	C11-S2-C12	97.1(3)
C11-S1-C13B	95.9(4)	C11-S2-C12B	97.7(3)
C1-C11-S1	124.3(2)	C1-C11-S2	121.5(2)
C12-C13-S1	105.7(10)	O1-C2-C3	117.8(3)
C11-C1-C2	122.1(3)	O1-C2-C1	121.4(3)
C4-C3-C2	122.1(3)	C3-C4-C5	124.4(3)
S1-C11-S2	114.2(15)	C13-C12-S2	104.9(10)
C1-C2-C3	120.9(3)	C13B-C12B-S2	107.7(9)

When the coordination reactions were carried out with ligands **3** and **4** (Scheme 4), the formation of  $\eta^2$ -[Fe(CO)] complexes was observed. The expected  $\eta^2$ -[Fe(CO)]



Scheme 3.




Scheme 4.

and **4a**) and  $\eta^4$ -[Fe(CO)<sub>3</sub>] (**3b** and **4b**) complexes were obtained as orange crystalline solids. These complexes displayed in IR spectra 3 bands around 2000 cm<sup>-1</sup> that correspond to  $\nu$ (M-CO) region. In MS the corresponding peaks assigned to the fragments [M<sup>+</sup>-28] were observed. The <sup>1</sup>H NMR spectra for these complexes exhibit the peaks corresponding to the AB system shifted upfield, indicating the coordination of the iron complex to the C=C  $\pi$ -system. The chemical shift of the other signals was very similar to that of the free-ligands. The <sup>13</sup>C NMR spectra of these complexes show that the corresponding signals to the C=C double bond are shifted to higher frequencies ( $\Delta\delta$  around 67). Signals around 200 and 208 ppm were assigned for M-CO group. The most salient difference between  $\eta^2$ -[Fe(CO)<sub>4</sub>] and  $\eta^4$ -[Fe(CO)<sub>4</sub>] complexes is that for  $\eta^4$ -[Fe(CO)<sub>3</sub>] complexes (**3b** and **4b**), the ketone group signal was shifted from 184 ppm at 155 ppm and this fact indicates the coordination of CO group to iron atom.

The X-ray diffraction analysis for **4b** (Figure 2), showed that the [Fe(CO)<sub>3</sub>] fragment was bonded in  $\eta^4$ -coordination mode to the enone moiety O4-C5-C6-C7, thus the coordination polyhedron can be best described as a distorted trigonal bipyramid, with two carbonyl groups

and the C3-C4 double bond in the equatorial positions, with the O4 oxygen atom and the remaining CO group in apical positions with a bond angle of 164.5(2)° (Table 2). This arrangement is in agreement with other  $\eta^4$ -[Fe(CO)<sub>3</sub>] complexes described in literature.<sup>10</sup> One O4---S1 interaction (2.757 Å) was also present and was in the same order than with the obtained crystallographic data from **3**. The 1,3-dithiolane ring adopts an envelope conformation with C16(C16B) as flap, the crystalline structure showed disorder in this moiety leading two conformers in 77/23 ratio, only the major contributor are shown in Figure 2.

Additionally, the dinuclear complexes **3c** and **4c** were obtained in moderate yields. They have the same structural arrangement as the complex **1c**, previously reported.<sup>6,11</sup> The IR spectra of these compounds show the typical bands of  $\nu$ (M-CO) in 2058 and 2017 cm<sup>-1</sup>. The <sup>1</sup>H NMR spectra of **3c** and **4c** exhibit changes in the assigned signals to the methylene groups, which shifted upfield with respect to ligands. In <sup>13</sup>C NMR spectra, as key signals, we observed that the alpha CH and the carbon attached to sulfur and iron atoms were shifted to higher frequencies (12 and 25 ppm respectively). In both reactions, complex **7** was obtained in a yield of 5%.



**Figure 2.** ORTEP drawing of ligand **4b**. Thermal ellipsoids at 30% probability level.

The reaction with ligands **5** and **6** lead to the formation of the  $\eta^4$ - $[\text{Fe}(\text{CO})_3]$  complexes (**5b** and **6b**) in low yield (Scheme 5), being the major products the dinuclear complexes **5c** and **6c**, these complexes displayed a similar structural arrangement to those of **1c**, **3c** and **4c** *vide supra*. In both experiments, complex **7** was also obtained in low yield and for the ligand **6**, the  $\kappa^N$ - $[\text{Fe}(\text{CO})_4]$  complex (**10**) was additionally obtained in traces. For **6b**, an additionally  $\sigma$ -N coordination of  $[\text{Fe}(\text{CO})_4]$  fragment towards amine group was evidenced by  $^1\text{H}$  and  $^{13}\text{C}$  NMR, as well as by a peak in 599  $m/z$  in MS.

We have carried out some experiments exploring different condition reaction (Table 4), which evidenced that the coordination reactions were sensitive to the temperature and the stability of the complexes. Thus, the

$\eta^2$ - $[\text{Fe}(\text{CO})_4]$  complexes are intermediates in the formation of most thermodynamic stable  $\eta^4$ - $[\text{Fe}(\text{CO})_3]$  complexes, whose formation depends strongly on the electronic characteristics of the C-C double bond.<sup>12</sup> The dinuclear complexes **3c-6c** are the intermediates in the formation of complex **7**, resulting from the total C-S bond activation of the corresponding ligands, as we have previously proposed.<sup>6</sup>

**Table 2.** Selected bond lengths (Å) and angles (°) for compound


Fe1-C3	1.745(6)	Fe1-C1	1.795(6)
Fe1-C2	1.830(6)	Fe1-O4	2.014(3)
Fe1-C6	2.037(4)	Fe1-C5	2.112(5)
Fe1-C7	2.134(4)	S1-C14	1.726(5)
S1-C15	1.798(12)	S2-C16	1.740(8)
S2-C14	1.748(5)	O1-C1	1.153(6)
O2-C2	1.124(5)	O3-C3	1.155(6)
O4-C5	1.299(5)	C4-C14	1.347(6)
C4-C5	1.449(6)	C5-C6	1.416(6)
C6-C7	1.416(5)	C15-C16	1.397(14)
S1-C15B	1.83(4)	S2-C16B	1.92(3)
C15B-C16B	1.57(5)	C1-Fe1-C2	104.6(2)
C3-Fe1-C1	96.8(3)	C3-Fe1-C2	90.5(2)
C3-Fe1-O4	164.5(2)	C3-Fe1-C6	97.8(2)
C1-Fe1-O4	97.7(2)	C2-Fe1-O4	91.15(19)
C1-Fe1-C6	131.3(2)	C3-Fe1-C5	127.9(2)
C1-Fe1-C5	132.3(2)	C6-Fe1-C5	39.9(16)
C3-Fe1-C7	95.9(2)	O4-Fe1-C7	77.96(15)
C6-Fe1-C7	39.6(14)	C16-S2-C14	96.5(4)
C2-Fe1-C6	121.44(2)	O4-Fe1-C6	63.36(15)
C2-Fe1-C5	92.0(2)	O4-Fe1-C5	36.62(13)
C1-Fe1-C7	92.8(2)	C2-Fe1-C7	160.7(2)
C5-Fe1-C7	69.67(17)	C14-S1-C15	92.8(4)
C14-C4-C5	123.3(5)	C14-S1-C15B	108.0(14)
O1-C1-Fe1	179.1(5)	O2-C2-Fe1	179.3(7)
C5-O4-Fe1	75.8(3)	O3-C3-Fe1	177.6(6)
O4-C5-C6	113.8(4)	O4-C5-C4	121.3(4)

**Table 3.** Summary of Crystal data, data collection, and refinement details

Compound	<b>3</b>	<b>4b</b>
Formula	$\text{C}_{13}\text{H}_{11}\text{ClO}_2\text{S}_2$	$\text{C}_{16}\text{H}_{11}\text{BrFeO}_4\text{S}_2$
Molecular weight (g/mol)	282.79	467.13
Description	Orange prism	Red prism
crystal size (mm)	0.308 x 0.30 x 0.108	0.224 x 0.09 x 0.06
crystalline system	Monoclinic	Monoclinic
Spatial group	$P2_1/n$	$P2_1/c$
Cell parameters (Å, °)	a = 6.1474(4) $\alpha$ = 90 b = 18.839(1) $\beta$ = 95.302(2) c = 11.090(1) $\gamma$ = 90	a = 7.661(1) $\alpha$ = 90 b = 17.609(1) $\beta$ = 91.449(2) c = 13.252(1) $\gamma$ = 90
Volume (Å <sup>3</sup> )	1278.9 (2)	1787.2(3)
Z	4	4
$D_{\text{calc}}$ (Mg/m <sup>3</sup> )	1.469	1.736
No. collect. reflex.	12640	17743
No. Ind. Reflex.	2923	4107
Date /parameters	2923/173	4107/226
$R_{\text{int}}$	0.0452	0.1069
RI, wR2 [ $I \geq 2\sigma(I)$ ]	0.0564, 0.1196	0.0485, 0.0556
RI, wR2 [all the date]	0.0859, 0.1315	0.1479, 0.0718
GOOF in $F^2$	0.993	0.957

**Table 4.** Reaction of ligands **3-6** with  $\text{Fe}_2(\text{CO})_9$ 

Condition reaction	Ligand	$\eta^2\text{-}[\text{Fe}(\text{CO})_4]$	$\eta^4\text{-}[\text{Fe}(\text{CO})_3]$	$\text{LFe}_2(\text{CO})_5$	$(\text{CH}_2\text{S})_2\text{Fe}_2(\text{CO})_6$ ( <b>7</b> )	$\text{LFe}(\text{CO})_4$
Ether, RT, 5h	<b>3</b>	<b>3a</b> : 25%	<b>3b</b> : 10%	<b>3c</b> : 8%	< 5%	
	<b>4</b>	<b>4a</b> : 18%	<b>4b</b> : 20%	<b>4c</b> : 18%		
THF, RT, 24h	<b>3</b>	<b>3a</b> : —	<b>3b</b> : 25%	<b>3c</b> : 7%	20%	
	<b>4</b>	<b>4a</b> : —	<b>4b</b> : 32%	<b>4c</b> : 10%		
Ether, RT, 5h	<b>5</b>	<b>5a</b> : —	<b>5b</b> : 20%	<b>5c</b> : 28%	< 5%	<b>10</b> : 2%
	<b>6</b>	<b>6a</b> : —	<b>6b</b> : 7%	<b>6c</b> : 25%		
THF, RT, 24h	<b>5</b>	<b>5a</b> : —	<b>5b</b> : 18%	<b>5c</b> : 12%	18%	<b>10</b> : —
	<b>6</b>	<b>6a</b> : —	<b>6b</b> : 5%	<b>6c</b> : 10%		
Refluxing THF, 24h	<b>6</b>	<b>15a</b> : —	<b>6b</b> : detected by TLC	<b>6c</b> : <5%	27%	<b>10</b> : —
Ether, 5°C, 24h	<b>6</b>	<b>15a</b> : —	<b>6b</b> : 13%	<b>6c</b> : 28%	—	<b>10</b> : 5%


**Scheme 5.**

Screening the reactivity of  $\text{Fe}_2(\text{CO})_9$  toward this kind of systems, the chemical shift of C-3 on  $^{13}\text{C}$  NMR have been analyzed using correlation with Hammett type substituent constant ( $\sigma_p$ )<sup>13</sup> (Table 5) giving a lineal correlation as shown in Figure 3.


The positive value of ( $\rho$ ) indicates that the electronic effect of the group on the aromatic ring increased the electronic density over C3 as a long distance  $\beta$  effect (Scheme 6).<sup>16</sup>

As a result of this study, and the several possibility of coordination ability of Fe(0) towards cross-conjugated compounds, we have observed that  $[\text{Fe}(\text{CO})_4]$  fragment has a preference to react with electronic deficient double bond in agreement with G. Reichenbach,<sup>17</sup> who proposed that the  $[\text{Fe}(\text{CO})_4]$  fragment is an electron-relaxing. Thus, when there are electron-withdrawing groups on the aromatic ring, the reaction produce the  $\eta^2\text{-}[\text{Fe}(\text{CO})_4]$  and  $\eta^4\text{-}[\text{Fe}(\text{CO})_4]$  complexes as the main compounds, in agreement with the Hammett plot.

Nevertheless, with electron-donating groups (Scheme 5), it might be expected that the transmission of electronic effects should increased by mesomeric effect over C3=C4 and then, the reaction proceeds differently leading as major products the dinuclear complexes resulting from the carbon-sulfur bond activation.

**Figure 3.** Hammett correlation,  $\delta = 7.1594 \sigma + 126.93$ ,  $r = 0.9993$ .**Table 5.** Chemical shift on  $^{13}\text{C}$  NMR from C3 and  $\sigma_p$ 

Ligand	R	$\delta$ C3	$\sigma_p$
2	<i>p</i> -NO <sub>2</sub>	132.5	0.78 <sup>14</sup>
3	<i>p</i> -Br	128.8	0.26 <sup>15</sup>
4	<i>p</i> -Cl	128.8	0.24 <sup>15</sup>
1	<i>p</i> -H	126.9	0
5	<i>p</i> -OMe	124.74	-0.28 <sup>15</sup>
6	<i>p</i> -N(Et) <sub>2</sub>	121.9	-0.72 <sup>14</sup>



Scheme 6.

## Conclusion

In this work, we have evidenced that the reactivity of  $\text{Fe}_2(\text{CO})_9$  with cross-conjugated carbonyl ligands depends on the electronic density of the heterodienic moiety, thus, electron-withdrawing groups promote the coordination reaction towards the formation of  $\eta^2\text{-}[\text{Fe}(\text{CO})_4]$  and  $\eta^4\text{-}[\text{Fe}(\text{CO})_3]$  complexes, while electron-donor groups aim the reactivity of  $\text{Fe}_2(\text{CO})_9$  towards the heterodienic moiety  $\beta$ -positioned by sulfur atoms leading the formation of dinuclear iron complexes, coming from the partial and total carbon-sulfur bonds activation.

## Experimental

$^1\text{H}$  NMR and  $^{13}\text{C}$  NMR spectra were recorded on a JEOL 300 spectrometer, using  $\text{CDCl}_3$ ,  $(\text{CD}_3)_2\text{SO}$  and  $\text{C}_6\text{D}_6$  as solvents and TMS as internal reference. IR spectra were performed on a Perkin-Elmer 283 B or 1420 spectrometer. The FAB spectra were obtained on a JEOL JMS SX 102A mass spectrometer operated at an accelerating voltage of 10 Kv. Samples were desorbed from a nitrobenzyl alcohol matrix using a 6 kev Xenon atoms. The electronic impact (EI) ionization mass spectra were acquired on a JEOL JMS-AX505 HA Mass spectrometer operated in the positive ion mode. The acquisition conditions were ion source temperature 230 °C, ionization energy 70 eV, emission current 0.14 mA and ionization current 100 mA. Mass measurements in FAB are performed at 10000 resolution using electrical field scans and the polyethylene glycol ions as reference material. Melting points were measured using a Mel-Temp II apparatus and are uncorrected. All reagents were obtained from commercial suppliers and used as received. Reactions were performed under nitrogen atmosphere in carefully dried glassware. THF and ether were distilled from sodium-benzophenone under argon and/or nitrogen atmosphere. Column chromatography was performed with Merck silica gel (70-230 mesh) or alumina using ethyl acetate: hexane in different ratios as eluent.

### Synthesis of ligands

To a solution of 5 mmol of 1-(1,3-dithiolane-2-yliden)acetone<sup>6</sup> in 20 mL of ethanol was added 10 mL of an

alcoholic solution of NaOH (5.5 mmol) at 0 °C. The solution of corresponding aldehyde (5 mmol) in 5 mL of ethanol was added and the reaction was keeping magnetic stirring during 18 h at room temperature. The solvent was eliminated for distillation at reduced pressure then Ethyl acetate (50 mL) and water (2 x 50 mL) were added, extracting the organic phase, drying with  $\text{Na}_2\text{SO}_4$  and removing the volatile *in vacuo*. The reaction mixture was purified by column chromatography on alumina, using hexane/AcOEt in different ratios as eluent.

### Synthesis of complexes

A solution of the ligand **2-6** (1 mmol) in anhydrous THF or Ether (40 mL) was treated with  $\text{Fe}_2(\text{CO})_9$  (3 mmol) with magnetic stirring at room temperature for 5 h under inert atmosphere. After the reaction was complete, the complex was filtered off through an alumina column and the solvent was evaporated under vacuum. The reaction mixture was chromatographed on silica gel and elution with hexane: ethyl acetate in different ratio gave the corresponding  $\eta^2$  (0) complexes. The yields were based on the pure product isolated.

### Ligands

**4-[p-Nitrophenyl]-1-(1,3-dithiolan-2-ylidene)buten-2-one, (2).** Purification: CC, Hexane/AcOEt, Yellow solid, (40%), m.p. 128 °C.  $^1\text{H}$  NMR (300 MHz,  $(\text{CD}_3)_2\text{SO}$ ):  $\delta$  3.48 (m, 4H,  $\text{CH}_2\text{S}$ ), 6.46 (s, 1H,  $\text{CH}=\text{CH}$ ), 7.13 and 7.63 (AB system, 2H,  $J$  16.10 Hz,  $\text{CH}=\text{CH}$ ), and 8.23 (AA'BB' system, 4H,  $J$  8.81 Hz,  $\text{CH}_{\text{arom}}$ ).  $^{13}\text{C}$  NMR (75 MHz,  $(\text{CD}_3)_2\text{SO}$ ):  $\delta$  35.5 ( $\text{CH}_2\text{S}$ ), 40.4 ( $\text{CH}_2\text{S}$ ), 112.0 ( $\text{CHCH}$ ), 112.0 ( $\text{CHCSS}$ ), 124.9 ( $\text{CH}_o\text{C}_i\text{NO}_2$ ), 129.5 ( $\text{CH}_o\text{C}_i\text{CH}$ ), 132.5 ( $\text{CHCO}$ ), 139.6 ( $\text{PhCH}$ ), 146.8 ( $\text{C}_i\text{N}$ ), 169.5 ( $\text{CSS}$ ), 184.3 ( $\text{CO}$ ). IR ( $\text{CHCl}_3$ )  $\nu_{\text{max}}$ /cm<sup>-1</sup>: 1594 ( $\text{C}=\text{C}$ ), 1499 ( $\text{C}=\text{C}$ ), 1440 ( $\text{NO}_2$ ). M.S. (E.I. 70 eV)  $m/z$  [ $\text{M}^+$ ](293), 265(90), 145(45).

**4-(p-Chlorophenyl)-1-(1,3-dithiolan-2-ylidene)buten-2-one, (3).** Purification: CC, Hexane/AcOEt, Pale yellow solid, (78%), m.p. 119 °C.  $^1\text{H}$  NMR (300 MHz,  $\text{CDCl}_3$ ):  $\delta$  3.48 (m, 2H,  $\text{CH}_2\text{S}$ ), 3.94 (m, 2H,  $\text{CH}_2\text{S}$ ), 6.81 (1H,  $\text{CHCSS}$ ), 6.75 and 7.57 (AB system, 2H,  $J$  15.9 Hz,  $\text{CH}=\text{CH}$ ), 7.34 and 7.48 (AA'BB' system, 4H,  $J$  8.5 Hz,  $\text{CH}_{\text{arom}}$ ).  $^{13}\text{C}$  NMR (75 MHz,  $\text{CDCl}_3$ ):  $\delta$  35.5 ( $\text{CH}_2\text{S}$ ), 40.4 ( $\text{CH}_2\text{S}$ ), 112.0 ( $\text{CHCH}$ ), 112.0 ( $\text{CHCSS}$ ), 124.9 ( $\text{CH}_o\text{C}_i\text{NO}_2$ ), 129.5 ( $\text{CH}_o\text{C}_i\text{CH}$ ), 132.5 ( $\text{CHCO}$ ), 139.6 ( $\text{PhCH}$ ), 146.8 ( $\text{C}_i\text{N}$ ), 169.5 ( $\text{CSS}$ ), 184.3 ( $\text{CO}$ ). IR ( $\text{CHCl}_3$ )  $\nu_{\text{max}}$ /cm<sup>-1</sup>: 1594 ( $\text{C}=\text{C}$ ), 1499 ( $\text{C}=\text{C}$ ), 1440 ( $\text{NO}_2$ ). M.S. (E.I. 70 eV)  $m/z$  [ $\text{M}^+$ ](293), 265(90), 145(45).

$\text{CH}_{\text{arom}}$ ).  $^{13}\text{C}$  NMR (75 MHz,  $\text{CDCl}_3$ ):  $\delta$  35.5 ( $\text{CH}_2\text{S}$ ), 39.1 ( $\text{CH}_2\text{S}$ ), 111.9 ( $\text{CHCSS}$ ), 127.3 ( $\text{CHCO}$ ), 129.2 ( $\text{CH}_o\text{C}_i\text{Cl}$ ), 129.4 ( $\text{CH}_o\text{C}_i\text{CH}$ ), 135.9 ( $\text{C}_i\text{CH}$ ), 137.8 ( $\text{C}_i\text{Cl}$ ), 140.1 ( $\text{PhCH}$ ), 167.5 ( $\text{CSS}$ ), 184.3 ( $\text{CO}$ ). IR ( $\text{CHCl}_3$ )  $\nu_{\text{max}}/\text{cm}^{-1}$ : 1633 ( $\text{C=O}$ ), 1493 ( $\text{C=C}$ ). M.S. (FAB<sup>+</sup> 70 eV)  $m/z$  (%):  $[\text{M}^+ + 1]$  283(100), 154(5), 136(35).

4-(*p*-Bromophenyl)-1-(1,3-dithiolan-2-ylidene)-3E-buten-2-one, (**4**). Purification: CC, Hexane/AcOEt, 6:4. Pale yellow solid, (80%) m.p. 145-7 °C.  $^1\text{H}$  NMR (300 MHz,  $(\text{CD}_3)_2\text{SO}$ ):  $\delta$  3.47 (m, 4H,  $\text{CH}_2\text{S}$ ), 6.42 (s, 1H,  $\text{CHCSS}$ ), 6.97 and 7.52 (AB system, 2H,  $J$  16.1 Hz,  $\text{CH=CH}$ ), 7.18 y 7.41 (AA'BB' system, 4H,  $J$  8.26 Hz,  $\text{CH}_{\text{arom}}$ ).  $^{13}\text{C}$  NMR (75 MHz,  $(\text{CD}_3)_2\text{SO}$ ):  $\delta$  35.8 ( $\text{CH}_2\text{S}$ ), 39.3 ( $\text{CH}_2\text{S}$ ), 111.6 ( $\text{CHCSS}$ ), 124.2 ( $\text{CBr}$ ), 128.7 ( $\text{CHCO}$ ), 130.6 ( $\text{CH}_o\text{C}_i\text{CH}$ ), 132.5 ( $\text{CH}_o\text{C}_i\text{Br}$ ), 134.8 ( $\text{C}_i\text{CH}$ ), 139.7 ( $\text{PhCH}$ ), 168.1 ( $\text{CSS}$ ), 184.5 ( $\text{CO}$ ). IR ( $\text{CHCl}_3$ )  $\nu_{\text{max}}/\text{cm}^{-1}$ : 1633 ( $\text{C=O}$ ), 1489 ( $\text{C=C}$ ). M.S. (E.I. 70 eV)  $m/z$  (%): 328 $[\text{M}^+]$ (62), 300(52), 216(46), 145(100).

4-(*p*-Methoxyphenyl)-1-(1,3-dithiolan-2-ylidene)-3E-buten-2-one, (**5**). Purification: CC, Hexane/AcOEt, 85:15. Yellow crystalline solid (64%), m.p. 101-103 °C.  $^1\text{H}$  NMR (300 MHz,  $\text{CDCl}_3$ ):  $\delta$  3.34 (m, 2H,  $\text{CH}_2\text{S}$ ), 3.44 (m, 2H,  $\text{CH}_2\text{S}$ ), 3.82 (s, 3H,  $\text{CH}_3\text{O}$ ), 6.82 (s, 1H,  $\text{CHCSS}$ ), 6.65 and 7.58 (AB system, 2H,  $J$  15.82 Hz,  $\text{CH=CH}$ ), 6.88 and 7.49 (AA'BB' system, 4H,  $J$  8.81 Hz,  $\text{CH}_{\text{arom}}$ ).  $^{13}\text{C}$  NMR (75 MHz,  $\text{CDCl}_3$ ):  $\delta$  35.5 ( $\text{CH}_2\text{S}$ ), 39.0 ( $\text{CH}_2\text{S}$ ), 55.5 ( $\text{CH}_3\text{O}$ ), 112.1 ( $\text{CHCSS}$ ), 114.4 ( $\text{CH}_o\text{C}_i\text{OMe}$ ), 124.7 ( $\text{CHCO}$ ), 127.9 ( $\text{C}_i\text{CH}$ ), 129.9 ( $\text{CH}_o\text{C}_i\text{CH}$ ), 141.4 ( $\text{PhCH}$ ), 161.3 ( $\text{C}_i\text{OCH}_3$ ), 166.1 ( $\text{CSS}$ ), 184.8 ( $\text{CO}$ ). IR ( $\text{CHCl}_3$ )  $\nu_{\text{max}}/\text{cm}^{-1}$ : 1642 ( $\text{C=O}$ ), 1588, 1573 and 1507 ( $\text{C=C}$ ). M.S. (E.I. 70 eV)  $m/z$  (%): 278 $[\text{M}^+]$ (100), 250(55), 206(20), 166(95).

4-[*p*-(*N*,*N*-diethylamino)-phenyl]-1-(1,3-dithiolan-2-ylidene)-3E-buten-2-one, (**6**). Purification: CC, Hexane/AcOEt, 8:2. Red crystalline solid (68%), m.p. 126-127 °C.  $^1\text{H}$  NMR (300 MHz,  $\text{CDCl}_3$ ):  $\delta$  1.17 (t, 6H,  $\text{CH}_3\text{CH}_2\text{N}$ ), 3.38 (q, 4H,  $\text{CH}_2\text{N}$ ), 3.38 (m, 4H,  $\text{CH}_2\text{S}$ ), 6.82 (s, 1H,  $\text{CHCSS}$ ), 6.57 and 7.57 (AB system, 2H,  $J$  15.8 Hz,  $\text{CH=CH}$ ), 6.62 and 7.42 (AA'BB' system, 4H,  $J$  8.8 Hz,  $\text{CH}_{\text{arom}}$ ).  $^{13}\text{C}$  NMR (75 MHz,  $\text{CDCl}_3$ ):  $\delta$  12.7 ( $\text{CH}_3\text{CH}_2\text{N}$ ), 35.2 ( $\text{CH}_2\text{S}$ ), 39.2 ( $\text{CH}_2\text{S}$ ), 44.8 ( $\text{CH}_2\text{N}$ ), 111.6 ( $\text{CH}_o\text{C}_i\text{N}$ ), 112.7 ( $\text{CHCSS}$ ), 121.9 ( $\text{CHCO}$ ), 122.4 ( $\text{C}_i\text{CH}$ ), 130.5 ( $\text{CH}_o\text{C}_i\text{CH}$ ), 142.6 ( $\text{PhCH}$ ), 149.3 ( $\text{C}_i\text{N}$ ), 167.9 ( $\text{CSS}$ ), 185.1 ( $\text{CO}$ ). IR ( $\text{CHCl}_3$ )  $\nu_{\text{max}}/\text{cm}^{-1}$ : 1632.4 ( $\text{C=O}$ ), 1565, 1517 y 1496 ( $\text{C=C}$ ). M.S. (E.I. 70 eV)  $m/z$  (%): 319 $[\text{M}^+]$  (100), 304(90), 291(20).

### Complexes

$\eta^2$ -[4-(*p*-Nitrophenyl)-1-(1,3-dithiolan-2-ylidene)-3E-buten-2-one] $\text{Fe}(\text{CO})_3$  (**2a**). Purification: CC, Hexane/AcOEt, 98:2. Orange cristalin solid (2%), m.p. 130 °C (dec). IR ( $\text{CHCl}_3$ )  $\nu_{\text{max}}/\text{cm}^{-1}$ : 2928 ( $\text{Csp}^2\text{-H}$ ), 2068, 2011 y 1990

( $\text{CO-M}$ ), 1598 ( $\text{C=O}$ ), 1520 ( $\text{C=C}$ ), 1343 ( $\text{NO}_2$ ). M.S.-E.I. (70 eV)  $m/z$  (%): 461 $[\text{M}^+]$ (5), 433(8) ( $\text{M}^+ - \text{CO}$ ), 377(10) ( $\text{M}^+ - 3\text{CO}$ ), 293(100)  $[(\text{M}^+ - \text{Fe}(\text{CO})_4)]$ .

$\eta^2$ -[4-(*p*-Clorophenyl)-1-(1,3-dithiolan-2-ylidene)-3E-buten-2-one] $\text{Fe}(\text{CO})_3$  (**3a**). Purification: CC, Hexane/AcOEt, 98:2. Red cristalin solid (25%), m.p.:95 °C (dec).  $^1\text{H}$  NMR (300 MHz,  $\text{CDCl}_3$ ):  $\delta$  3.43 (m, 4H,  $\text{CH}_2\text{S}$ ), 5.85 (s, 2H,  $\text{CHCHCO}$ ), 6.71 (s, 1H,  $\text{CHCSS}$ ), 7.34 and 7.50 (d, 4H,  $\text{CH}_{\text{arom}}$ ).  $^{13}\text{C}$  NMR (75 MHz,  $\text{CDCl}_3$ ):  $\delta$  35.9 and 39.4 ( $\text{SCH}_2$ ), 59.6 ( $\text{CHCHCO}$ ), 74.1 ( $\text{CHCHCO}$ ), 105.4 ( $\text{CHCSS}$ ), 127.9 ( $\text{CH}_o\text{C}_i\text{Cl}$ ), 129.0 ( $\text{CH}_o\text{C}_i\text{CH}$ ), 131.7 ( $\text{C}_i\text{Cl}$ ), 138.4 ( $\text{C}_{\text{ipso}}\text{CH}$ ), 182.4 ( $\text{CO}$ ), 202.9 ( $\text{M-CO}$ ). IR ( $\text{CHCl}_3$ )  $\nu_{\text{max}}/\text{cm}^{-1}$ : 2095, 2019, 1989 ( $\text{M-CO}$ ), 1608 ( $\text{CO}$ ). M.S.-FAB<sup>+</sup> ( $m/z$ ): 422 ( $\text{M}^+ - \text{CO}$ ), 394 ( $\text{M}^+ - 2\text{CO}$ ), 282  $[\text{M}^+ - \text{Fe}(\text{CO})_4]$ . H.R.-M.S. (FAB<sup>+</sup>)  $\text{C}_{16}\text{H}_{12}\text{O}_4\text{ClS}_2\text{Fe}$ , Exp.: 422.9207, Calc.: 422.9215.

$\eta^4$ -[4-(*p*-Clorophenyl)-1-(1,3-dithiolan-2-ylidene)-3E-buten-2-one] $\text{Fe}(\text{CO})_3$  (**3b**). Purification: CC, Hexane/AcOEt, 96:4. Orange cristalin solid (10%), m.p.:128-130 °C (dec).  $^1\text{H}$  NMR (300 MHz,  $\text{C}_6\text{D}_6$ ):  $\delta$  2.39 (m, 4H,  $\text{CH}_2\text{S}$ ), 3.25 (s, 1H,  $\text{CHCHCO}$ ), 5.21 (s, 1H,  $\text{CHCHCO}$ ), 6.39 (s, 1H,  $\text{CH=CSS}$ ), 6.74 and 6.92 (d, 4H,  $J$  7.29 Hz  $\text{CH}_{\text{arom}}$ ).  $^{13}\text{C}$  NMR (75 MHz,  $\text{C}_6\text{D}_6$ ):  $\delta$  35.2 and 38.8 ( $\text{SCH}_2$ ), 59.6 ( $\text{CHCHCO}$ ), 73.2 ( $\text{CHCHCO}$ ), 105.1 ( $\text{CHCSS}$ ), 128.0 ( $\text{CH}_o\text{C}_i\text{Cl}$ ), 128.8 ( $\text{CH}_o\text{C}_i\text{CH}$ ), 132.1 ( $\text{C}_i\text{Cl}$ ), 138.2 ( $\text{C}_i\text{CH}$ ), 143.0 ( $\text{CSS}$ ), 155.4 ( $\text{CO}$ ), 212.2 ( $\text{M-CO}$ ). IR ( $\text{CHCl}_3$ )  $\nu_{\text{max}}/\text{cm}^{-1}$ : 2064, 2006, 1986 ( $\text{M-CO}$ ), 1601 ( $\text{CO}$ ). M.S.-E.I. ( $m/z$ ): 394 ( $\text{M}^+ - \text{CO}$ ), 338 ( $\text{M}^+ - 3\text{CO}$ ), 282  $[\text{M}^+ - \text{Fe}(\text{CO})_3]$ .

[4-(*p*-Clorophenyl)-1-(1,3-dithiolan-2-ylidene)-3E-buten-2-one] $\text{Fe}_2(\text{CO})_5$  (**3c**). Purification: CC, Hexane/AcOEt, 9:1. Deep red solid (8%), m.p.:118 °C (dec).  $^1\text{H}$  NMR (300 MHz,  $\text{CDCl}_3$ ):  $\delta$  2.34 (m, 2H,  $\text{FeSCH}_2$ ), 3.34 (m, 2H,  $\text{SCH}_2$ ), 7.08 (s, 1H,  $\text{CH=CSFe}$ ), 6.81 and 7.57 (AB system, 2H,  $J$  15.98 Hz,  $\text{CHCHCO}$ ), 7.53 and 7.71 (AA'BB' system, 4H,  $J$  8.92 Hz  $\text{CH}_{\text{arom}}$ ).  $^{13}\text{C}$  NMR (75 MHz,  $\text{CDCl}_3$ ):  $\delta$  30.4 ( $\text{FeSCH}_2$ ), 32.0 ( $\text{SCH}_2$ ), 132.6 ( $\text{C}_i\text{-Cl}$ ), 114.2 ( $\text{CHCSFe}$ ), 128.9 and 129.2 ( $\text{CH}_{\text{ar}}$ ), 127.8 ( $\text{CHCHCO}$ ), 139.4 ( $\text{CHCHCO}$ ), 133.9 ( $\text{C}_i\text{CH}$ ), 188.7 ( $\text{CSFe}$ ), 189.4 ( $\text{CO}$ ), 211.3 and 213.9 ( $\text{M-CO}$ ) ppm. IR ( $\text{CHCl}_3$ )  $\nu_{\text{max}}/\text{cm}^{-1}$ : 2065, 2027 ( $\text{M-CO}$ ). M.S.-FAB<sup>+</sup> ( $m/z$ ): 478 ( $\text{M}^+ - 2\text{CO}$ ), 339  $[(\text{M}^+ + 1) - \text{Fe}(\text{CO})_5]$ .

$\eta^2$ -[4-(*p*-Bromophenyl)-1-(1,3-dithiolan-2-ylidene)-3E-buten-2-one] $\text{Fe}(\text{CO})_3$  (**4a**). Purification: CC, Hexane/AcOEt, 99:1. Red solid (18%), m.p.:156-8 °C (dec);  $^1\text{H}$  NMR (300 MHz,  $\text{CDCl}_3$ ):  $\delta$  3.38 (m, 4H,  $\text{CH}_2\text{S}$ ); 5.85 (AB system, 1H,  $J$  11.25 Hz,  $\text{CHCHCO}$ ), 6.72 (s, 1H,  $\text{CH=CSS}$ ), 7.19 and 7.38 (AA'BB' system, 4H,  $J$  6.33 Hz  $\text{CH}_{\text{arom}}$ ).  $^{13}\text{C}$  NMR (75 MHz,  $\text{CDCl}_3$ ):  $\delta$  31.1 and 36.3 ( $\text{SCH}_2$ ), 56.4 ( $\text{CHCHCO}$ ), 65.9 ( $\text{CHCHCO}$ ), 105.6 ( $\text{CHCSS}$ ), 123.3 ( $\text{C}_i\text{Br}$ ), 129.6 ( $\text{CH}_o\text{C}_i\text{CH}$ ), 132.1 ( $\text{CH}_o\text{C}_i\text{Br}$ ), 138.7 ( $\text{C}_i\text{CH}$ ), 163.8 ( $\text{CSS}$ ), 190.8 ( $\text{CO}$ ), 200.5 and 207.5 ( $\text{M-CO}$ ). IR ( $\text{CHCl}_3$ )  $\nu_{\text{max}}/\text{cm}^{-1}$ : 2098, 2016, 1985 ( $\text{M-CO}$ ), 1589 ( $\text{CO}$ ).



M.S.-FAB<sup>+</sup> ( $m/z$ ): 495 [ $\text{M}^+$ ], 467 and 469 ( $\text{M}^+-\text{CO}$ ), 411 and 412 [ $\text{M}^+-3\text{CO}$ ], 327 and 329 [ $\text{M}^+-\text{Fe}(\text{CO})_4$ ].

$\eta^4$ -[4-(*p*-Bromophenyl)-1-(1,3-dithiolan-2-ylidene)-3*E*-buten-2-one]  $\text{Fe}(\text{CO})_5$ , (**4b**). Purification: CC, Hexane/AcOEt, 97:3. Orange solid (20%), p.f.: 83 °C (dec). <sup>1</sup>H NMR (300 MHz,  $\text{C}_6\text{D}_6$ ):  $\delta$  2.36 (m, 4H,  $\text{CH}_2$ ), 3.22 and 5.18 (AB system, 1H,  $J$  8.67 Hz,  $\text{CHCHCO}$ ), 6.38 (s, 1H,  $\text{CH}=\text{CSS}$ ), 6.64 and 7.07 (AA'BB' system, 4H,  $J$  7.29 Hz  $\text{CH}_{\text{arom}}$ ). <sup>13</sup>C NMR (75 MHz,  $\text{C}_6\text{D}_6$ ):  $\delta$  35.3 and 38.8 ( $\text{SCH}_2$ ), 59.7 ( $\text{CHCHCO}$ ), 73.2 ( $\text{CHCHCO}$ ), 105.1 ( $\text{CHCSS}$ ), 120.2 ( $\text{C}_i\text{Br}$ ), 128.1 ( $\text{CH}_o\text{C}_i\text{CH}$ ), 131.7 ( $\text{CH}_o\text{C}_i\text{Br}$ ), 138.7 ( $\text{C}_i\text{CH}$ ), 143.1 ( $\text{CSS}$ ), 155.5 ( $\text{CO}$ ), 200.1 ( $\text{M-CO}$ ). IR (KBr)  $\nu_{\text{max}}/\text{cm}^{-1}$ : 2060, 2002, 1982 ( $\text{M-CO}$ ), 1543 ( $\text{CO}$ ). M.S.-FAB<sup>+</sup> ( $m/z$ ): 469 and 467 [ $\text{M}^++1$ ], 440 and 438 ( $\text{M}^+-\text{CO}$ ); 411 and 412 [ $(\text{M}^++1)-2\text{CO}$ ], 383 and 382 ( $\text{M}^+-3\text{CO}$ ), 327 and 329 [ $\text{M}^+-\text{Fe}(\text{CO})_3$ ].

[4-(*p*-Bromophenyl)-1-(1,3-dithiolan-2-ylidene)-3*E*-buten-2-one]  $\text{Fe}_2(\text{CO})_9$ , (**4c**). Purification: CC, Hexane/AcOEt, 9:1. Deep red solid (18%), m.p.: 99 °C (dec). <sup>1</sup>H NMR (300 MHz,  $\text{C}_6\text{D}_6$ ):  $\delta$  1.48 (m, 2H,  $\text{FeSCH}_2$ ), 2.26 (m, 2H,  $\text{SCH}_2$ ), 6.92 (s, 1H,  $\text{CH}=\text{CSFe}$ ), 6.67 and 7.01 (AA'BB' system, 4H,  $J$  8.64 Hz  $\text{CH}_{\text{arom}}$ ), 6.57 and 7.36 (AB system, 2H,  $J$  16.08 Hz,  $\text{CHCHCO}$ ). <sup>13</sup>C NMR (75 MHz,  $\text{C}_6\text{D}_6$ ):  $\delta$  30.5 ( $\text{FeSCH}_2$ ), 35.6 ( $\text{SCH}_2$ ), 122.8 ( $\text{C}_i\text{Br}$ ), 124.5 ( $\text{CHCSFe}$ ), 128.8 ( $\text{CH}_o\text{C}_i\text{CH}$ ), 129.4 ( $\text{CHCHCO}$ ), 131.7 ( $\text{CH}_o\text{C}_i\text{Br}$ ), 134.7 ( $\text{CH-CH-CO}$ ), 135.3 ( $\text{C}_i\text{CH}$ ), 188.6 ( $\text{CSFe}$ ), 189.0 ( $\text{CO}$ ), 211.6 and 213.5 ( $\text{M-CO}$ ). IR (KBr)  $\nu_{\text{max}}/\text{cm}^{-1}$ : 2059, 2017 ( $\text{M-CO}$ ); 1641 ( $\text{CO}$ ). M.S.-FAB<sup>+</sup> ( $m/z$ ): 579 and 580 [ $\text{M}^++1$ ], 552 ( $\text{M}^+-\text{CO}$ ), 524 ( $\text{M}^+-2\text{CO}$ ), 495 and 497 [ $(\text{M}^++1)-3\text{CO}$ ], 440 and 438 [ $\text{M}^+-\text{Fe}(\text{CO})_3$ ].

$\eta^4$ -[4-(*p*-Methoxyphenyl)-1-(1,3-dithiolan-2-ylidene)-3*E*-buten-2-one]  $\text{Fe}(\text{CO})_5$ , (**5b**). Purification: CC, Hexane/AcOEt, 96:4. Red solid (20%), m.p.: 176 °C (dec.). <sup>1</sup>H NMR (300 MHz,  $\text{C}_6\text{D}_6$ ):  $\delta$  2.40 (m, 4H,  $\text{CH}_2$ ), 3.20 (s, 3H,  $\text{CH}_3\text{O}$ ), 3.59 (d, 1H,  $J$  8.94 Hz,  $\text{CHCHCO}$ ), 5.44 (d, 1H,  $J$  8.94 Hz,  $\text{CHCHCO}$ ), 6.45 (s, 1H,  $\text{CH}=\text{CSS}$ ), 6.63 and 7.03 (AA'BB' system, 4H,  $J$  8.26 Hz  $\text{CH}_{\text{arom}}$ ). <sup>13</sup>C NMR (75 MHz,  $\text{CDCl}_3$ ):  $\delta$  35.3 and 38.8 ( $\text{SCH}_2$ ), 54.5 ( $\text{CH}_3\text{O}$ ), 62.4 ( $\text{CHCHCO}$ ), 74.2 ( $\text{CHCHCO}$ ), 105.8 ( $\text{CHCSS}$ ), 114.4 ( $\text{CH}_o\text{C}_i\text{OMe}$ ), 128.5 ( $\text{CH}_o\text{C}_i\text{CH}$ ), 131.3 ( $\text{C}_i\text{CH}$ ), 141.4 ( $\text{C}_i\text{OMe}$ ), 154.0 ( $\text{CSS}$ ), 158.9 ( $\text{CO}$ ), 214.1 ( $\text{M-CO}$ ). IR (KBr)  $\nu_{\text{max}}/\text{cm}^{-1}$ : 2056, 1998 and 1979 ( $\text{M-CO}$ ), 1543 ( $\text{CO}$ ). M.S.-FAB<sup>+</sup> ( $m/z$ ): 419 [ $\text{M}^++1$ ], 391 [ $(\text{M}^++1)-\text{CO}$ ], 363 [ $(\text{M}^++1)-2\text{CO}$ ], 334 ( $\text{M}^+-3\text{CO}$ ), 279 ( $\text{M}^+-\text{Fe}(\text{CO})_3$ ).

[4-(*p*-Methoxyphenyl)-1-(1,3-dithiolan-2-ylidene)-3*E*-buten-2-one]  $\text{Fe}_2(\text{CO})_9$ , (**5c**). Purification: CC, Hexane/AcOEt, 9:1. Deep red solid (28%), m.p.: 164-168 °C (dec.). RMN <sup>1</sup>H (300 MHz,  $\text{C}_6\text{D}_6$ ):  $\delta$  1.54 and 2.39 (m, 2H,  $\text{FeSCH}_2$ ), 3.18 (s, 3H,  $\text{CH}_3\text{O}$ ), 3.22 and 4.27 (m, 2H,  $\text{SCH}_2$ ), 6.58 and 7.13 (AA'BB' system, 4H,  $J$  8.79 Hz  $\text{CH}_{\text{arom}}$ ), 6.77 and 7.67 (AB system, 2H,  $J$  15.8 Hz,  $\text{CHCHCO}$ ), 7.03 (s, 1H,  $\text{CH}=\text{CSFe}$ ). <sup>13</sup>C NMR (75 MHz,  $\text{C}_6\text{D}_6$ ):  $\delta$  30.0 ( $\text{FeSCH}_2$ ),

35.3 ( $\text{SCH}_2$ ), 54.6 ( $\text{CH}_3\text{O}$ ), 114.2 ( $\text{CH}_o\text{C}_i\text{OMe}$ ), 120.8 ( $\text{CSFe}$ ), 124.0 ( $\text{CHCHCO}$ ), 129.0 ( $\text{CH}_o\text{C}_i\text{CH}$ ), 129.7 ( $\text{C}_i\text{CH}$ ), 136.5 ( $\text{CHCHCO}$ ), 161.4 ( $\text{C}_i\text{OMe}$ ), 176 ( $\text{CSFe}$ ), 191.2 ( $\text{CO}$ ), 219.5 ( $\text{M-CO}$ ). IR (KBr)  $\nu_{\text{max}}/\text{cm}^{-1}$ : 2060, 2020 ( $\text{M-CO}$ ), 1603 ( $\text{CO}$ ). M.S.-FAB<sup>+</sup> ( $m/z$ ): 531 [ $\text{M}^++1$ ], 474 ( $\text{M}^+-2\text{CO}$ ), 446 ( $\text{M}^+-3\text{CO}$ ); 419 [ $(\text{M}^++1)-4\text{CO}$ ].

$\eta^4$ ,  $\kappa^N$ -[4-[*p*-(*N,N*-diethylamino)-phenyl]-1-(1,3-dithiolan-2-ylidene)-3*E*-buten-2-one]  $\text{Fe}_2(\text{CO})_9$ , (**6c**). Purification: CC, Hexane/AcOEt, 95:5. Deep red oil, (25%). <sup>1</sup>H NMR (300 MHz,  $\text{CDCl}_3$ ):  $\delta$  0.89 (q, 4H,  $\text{CH}_2\text{N}$ ), 1.2 (t, 3H,  $\text{CH}_3$ ), 3.43 (m, 4H,  $\text{CH}_2\text{S}$ ), 6.7 (s, 1H,  $\text{CH}=\text{CSS}$ ), 5.34 (AB system, 2H,  $\text{CH}=\text{CH}$ ), 6.66 and 7.32 (AA'BB' system, 4H,  $J$  8.1 Hz,  $\text{CH}_{\text{arom}}$ ). IR ( $\text{CHCl}_3$ )  $\nu_{\text{max}}/\text{cm}^{-1}$ : 2072, 2016 and 2003, ( $\text{M-CO}$ ). M.S. (FAB<sup>+</sup>)  $m/z$ : 599 ( $\text{M}^++1$ ), 571 ( $\text{M}^+-2\text{CO}$ ), 543 ( $\text{M}^+-3\text{CO}$ ), 431 [ $\text{M}^+-\text{Fe}(\text{CO})_3$ ], 403 [ $\text{M}^+-\text{Fe}(\text{CO})_4$ ].

[4-[*p*-(*N,N*-diethylamino)-phenyl]-1-(1,3-dithiolan-2-ylidene)-3*E*-buten-2-one]  $\text{Fe}_2(\text{CO})_9$ , (**6c**). Purification: Hexane/AcOEt, 92:8. Deep red oil, (25%). <sup>1</sup>H NMR (300 MHz,  $\text{CDCl}_3$ ):  $\delta$  1.26 (t, 6H,  $\text{CH}_3\text{CH}_2\text{N}$ ), 3.37 (q, 4H,  $\text{CH}_2\text{S}$ ), 3.16 and 1.53 (2m, 4H,  $\text{CH}_2\text{S}$ ), 7.08 (s, 1H,  $\text{CHCSFe}$ ), and 7.22 (s, 2H,  $\text{CH}=\text{CH}$ ), 6.61 and 7.37 (AA'BB' system, 4H,  $J$  8.6 Hz,  $\text{CH}_{\text{arom}}$ ). <sup>13</sup>C NMR (75 MHz,  $\text{CDCl}_3$ ):  $\delta$  31.0 ( $\text{CH}_3\text{CH}_2\text{N}$ ), 31.0 ( $\text{CH}_2\text{SFe}$ ), 36.1 ( $\text{CH}_2\text{S}$ ), 44.5 ( $\text{CH}_3$ ), 111.4 ( $\text{CH}_o\text{C}_i\text{N}$ ), 118.5 ( $\text{CHCSFe}$ ), 123.1 ( $\text{CHCO}$ ), 129.3 ( $\text{CH}_o\text{C}_i\text{CH}$ ), 136.9 ( $\text{PhCH}$ ), 148.6 ( $\text{C}_i\text{N}$ ), 191.5 ( $\text{CSFe}$ ), 191.5 ( $\text{CO}$ ), 207.0 ( $\text{CO-M}$ ). IR ( $\text{CHCl}_3$ )  $\nu_{\text{max}}/\text{cm}^{-1}$ : 2003, 2016 ( $\text{M-CO}$ ), 1632 ( $\text{C=O}$ ). M.S. (FAB<sup>+</sup>)  $m/z$ : 543 ( $\text{M}^++1$ ), 543 ( $\text{M}^+-\text{CO}$ ), 515 ( $\text{M}^+-2\text{CO}$ ), 487 ( $\text{M}^+-3\text{CO}$ ), 403 [ $\text{M}^+-\text{Fe}(\text{CO})_4$ ].

4-(*p*-aminophenyl)-1-(1,3-dithiolan-2-ylidene)-3*E*-buten-2-one, (**8**) Purification: CC, Hexane/AcOEt, 9:1. Orange solid (75%), m.p.: 233-235 °C; <sup>1</sup>H NMR (300 MHz,  $\text{CDCl}_3$ ):  $\delta$  3.38 (m, 4H,  $\text{CH}_2\text{S}$ ), 6.57 (s, 2H,  $\text{NH}_2$ ), 6.8 (s, 1H,  $\text{CHCSS}$ ), 6.59 and 7.54 (AB system, 2H,  $J$  15.81 Hz,  $\text{CH}=\text{CH}$ ), 6.63 and 7.37 (AA'BB' system, 4H,  $J$  8.4 Hz,  $\text{CH}_{\text{arom}}$ ). <sup>13</sup>C NMR (75 MHz,  $\text{CDCl}_3$ ):  $\delta$  35.3 ( $\text{CH}_2\text{S}$ ), 35.3 ( $\text{CH}_2\text{S}$ ), 112.3 ( $\text{CHCSS}$ ), 114.9 ( $\text{CH}_o\text{C}_i\text{NH}_2$ ), 123.6 ( $\text{CHCO}$ ), 125.4 ( $\text{C}_i\text{CH}$ ), 130.1 ( $\text{CH}_o\text{C}_i\text{CH}$ ), 142.4 ( $\text{PhCH}$ ), 168.4 ( $\text{C}_i\text{NH}_2$ ), 168.4 ( $\text{CSS}$ ), 184.5 ( $\text{CO}$ ). IR ( $\text{CHCl}_3$ )  $\nu_{\text{max}}/\text{cm}^{-1}$ : 3408 (N-H), 1620 ( $\text{C=O}$ ), 1498 ( $\text{C=C}$ ). M.S.-E.I. (70 eV)  $m/z$  (%): 263 [ $\text{M}^+$ ] (100), 235 (40), 146 (25).

$\kappa^N$ -[4-(*p*-Aminophenyl)-1-(1,3-dithiolan-2-ylidene)-3*E*-buten-2-one]  $\text{Fe}(\text{CO})_5$ , (**9**). Purification: CC, Hexane/AcOEt, 9:1. Brown solid, (2%), m.p.: 166 °C (dec.). IR ( $\text{CHCl}_3$ )  $\nu_{\text{max}}/\text{cm}^{-1}$ : 3409 (N-H), 2061 and 2019 ( $\text{CO}$ ), 1600 ( $\text{C=O}$ ), 1498 ( $\text{C=C}$ ). M.S.-E.I. (70 eV)  $m/z$  (%): 431 [ $\text{M}^+$ ] (15), 403 (10) ( $\text{M}^+-\text{CO}$ ), 375 (5) ( $\text{M}^+-2\text{CO}$ ), 340 ( $\text{M}^+-3\text{CO}$ ), 263 (100) [ $\text{M}^+-\text{Fe}(\text{CO})_4$ ].

$\kappa^N$ -[4-[*p*-(*N,N*-diethylamino)-phenyl]-1-(1,3-dithiolan-2-ylidene)-3*E*-buten-2-one]  $\text{Fe}(\text{CO})_5$ , (**10**). Purification:

CC, Hexane: AcOEt, 99:1. Red solid, (2%) m.p.: 98–101 °C (dec). <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>): δ 0.86 (q, 4H, CH<sub>2</sub>N), 1.17 (t, 6H, CH<sub>3</sub>CH<sub>2</sub>N), 3.37 (m, 4H, CH<sub>2</sub>S), 6.82 (s, 1H, CHCSS), 6.56 and 7.57 (AB system, 2H, *J* 15.66 Hz, CH=CH), 6.62 and 7.39 (AA'BB' system, 4H, *J* 8.1 Hz, CH<sub>arom</sub>). IR (CHCl<sub>3</sub>)  $\nu_{\text{max}}$ /cm<sup>-1</sup>: 2003, 2016 (M-CO) 1632(C=O). E.M. (FAB<sup>+</sup>) *m/z*: 487(M<sup>+</sup>-28), 431(M<sup>+</sup>-2CO), 403(M<sup>+</sup>-3CO), 319[M<sup>+</sup>-(Fe(CO)<sub>4</sub>].

#### *X-ray crystal structure determinations of compounds 3 and 4b*

Data collection and refinement parameters are summarized in Table 3. The diffraction data for **3** and **4b** were collected on a Bruker Smart Apex CCD diffractometer with MoK $\alpha$  radiation,  $\lambda = 0.71063$  Å. Each data set was corrected for Lorentz and polarization effects and analytical absorption corrections based on face indexed were applied. The structures were solved by direct methods<sup>18</sup> and each structure was refined by full-matrix least squares on *F*<sup>2</sup> using all data with the all non-hydrogen atoms assigned anisotropic displacement parameters and hydrogen atoms bound to carbon atoms inserted at calculated position with isotropic temperature factor 1.2 times the U<sub>iso</sub> of the parent carbon atom. The program used in the final refinements was SHELXL-97.<sup>19</sup> Selected bond lengths and bond angles are shown in Tables 1–2.

### Supplementary Information

Crystallographic data (excluding structure factors) for the structures in this paper have been deposited with the Cambridge Crystallographic Data Centre as supplementary publication no CCDC No. 251031 for ligand **3** and No. 251032 for complex **4b**. Copies of the data can be obtained, free of charge via [www.ccdc.cam.ac.uk/conts/retrieving.html](http://www.ccdc.cam.ac.uk/conts/retrieving.html) (or from the Cambridge Crystallographic Data Centre, CCDC, 12 Union Road, Cambridge CB2 1EZ, UK ; fax: +44 1223 336033 ; or e-mail: [deposit@ccdc.cam.ac.uk](mailto:deposit@ccdc.cam.ac.uk)).

### Acknowledgements

We thank DGAPA for financial support (Project # IN242004). The technical assistance of Rocio Patiño, Luis Velasco and Javier Pérez is gratefully acknowledged.

### References

1. Hegedus L.S.; *Transition Metals in the Synthesis of Complex Organic Molecules*, University Science Books: CA, 1994, pp. 131–142, 220–225.

2. Howell, J.A. S.; Johnson, B.F. G.; Josty, P.L.; Lewis, J.; *J. Organomet. Chem.* **1972**, 39, 329; Domingos, J. P.; Howell, J. S. A.; Johnson, B. F. G.; Lewis, J.; *Inorg. Synth.* **1976**, 16, 103; Fleckner, H.; Grevels, F.W.; Hess, D.; *J. Am. Chem. Soc.* **1984**, 106, 2027; Knölker, H. J.; Baum, G.; Foitzik, N.; Goesmann, H.; Gonser, P.; Jones, P. G.; Röttele, H.; *Eur. J. Inorg. Chem.* **1998**, 993.
3. Brandsna, L.; Vasilesky, S. F.; Verbré, H. D.; *Application of Transition Metal Catalyst in Organic Chemistry*, Springer: New York, 1999, p.254–307; Barton D. H. R.; Gunatilaka, A. A. L.; Nakanishi, T.; Patin, H.; Widdowson, D. A.; Worth, B. R.; *J. Chem. Soc. Perkin Trans.1* **1976**, 821; Evans, G.; Johnson, B. F. G.; Lewis, J.; *J. Organomet. Chem.* **1975**, 102, 507.
4. Cox, L. R.; Ley, S. V.; *Chem. Soc. Rev.* **1998**, 27, 301; Takemoto, Y.; Baba, Y.; Yoshikawa, N.; Iwata, C.; Tanaka, T.; Ibuka, T.; *Chem. Commun.* **1998**, 1911; Knölker, H. J.; *Chem. Soc. Rev.* **1999**, 28, 151; Frank-Neuman, M.; Olson, P. J.; *Synlett* **1991**, 891; Donaldson, W. A.; Bell, P. T.; Wang, Z.; Bennett, D. W.; *Tetrahedron Lett.* **1994**, 32, 5892; Prahald, V.; Donaldson, W. A.; *Tetrahedron Lett.* **1996**, 37, 9169; Benvegnu, T. J.; Toupet, L. J.; Grée, R.; *Tetrahedron* **1996**, 52, 11811; Benvegnu, T. J.; Grée, R.; *Tetrahedron* **1996**, 52, 11821. Iwata, C.; Takemoto, Y.; *Chem. Commun.* **1996**, 2497.
5. Alvarez, C.; Gutiérrez, R.; Toscano, R. A.; Moya, M.; Velasco, L.; Rosas, R.; Tapia, R.; Penieres, G.; *J. Coord. Chem.* **1999**, 48, 383; Alvarez-Toledano, C.; Gutiérrez-Pérez, R.; Toscano, R. A.; Moya-Cabrera, M.; Haquet, T.; Ortega, M. C.; Cabrera, A.; *Polyhedron* **2001**, 20, 215; Alvarez-Toledano, C.; Enríquez, J.; Toscano, R. A.; Martínez-García, M.; Cortés-Cortés, E.; Osornio, Y.; García, O.; Gutierrez-Pérez, R.; *J. Organomet. Chem.* **1999**, 577, 38; Ortega, C.; Gutiérrez, R.; Sharma, P.; Gómez, E.; Toscano, A.; López, J. G.; Penieres, G.; Alvarez C.; *Trends Organomet. Chem.* **2002**, 4, 19.
6. Ortega-Jiménez, F.; Ortega-Alfaro, M. C.; López-Cortés, J. G.; Gutiérrez-Pérez, R.; Toscano, R. A.; Velasco-Ibarra, L.; Peña-Cabrera, E.; Alvarez-Toledano, C.; *Organometallics* **2000**, 19, 4127.
7. Karl Dieter, R.; *Tetrahedron* **1986**, 42, 3029; Yates, P.; Lynch, T. R.; Moore, D. R.; *Can. J. Chem.* **1971**, 49, 1467.
8. Shriver, D. F.; Whitmire K. H. In *Comprehensive Organometallic Chemistry. The Synthesis, Reactions and Structures of Organometallic Compounds*; Wilkinson, G.; Stone, F. G.; Abel, E. W., eds.; Pergamon Press: Oxford, 1982, vol. 4, p.251,252.
9. Hugues, D. L.; Leigh, G. L.; Paulson, D.; *Inorg. Chim. Acta* **1986**, 120, 191; King, R. B.; *J. Am. Chem. Soc.* **1963**, 85, 1584.
10. Bernès, S.; Toscano, R. A.; Cano, A. C.; Mellado, O. G.; Alvarez-Toledano, C.; Rudler, H.; Daran, J.; *J. Organomet. Chem.* **1995**, 498, 15; Alvarez-Toledano, C.; Hernández-Ortega, S.; Bernès, S.; Gutiérrez-Pérez, R.; García-Mellado, O.; *J. Organomet. Chem.* **1997**, 549, 49.



11. Ortega-Alfaro, M. C.; Hernández, N.; Cerna, I.; López-Cortés, J. G.; Gómez, E.; Toscano, R. A.; Alvarez-Toledano, C.; *J. Organomet. Chem.* **2004**, 689, 885.
12. Howell, J. A. S.; Johnson, B. F.; Josty, P. L.; Lewis, J.; *J. Organomet. Chem.* **1972**, 39, 329.
13. Geribaldi, S.; Azzaro, M.; *Spectrochim. Acta* **1982**, 38A, 779; Ewing, D. F. In *Correlation Analysis in Chemistry: Recent Advances*; Chapman, N. B.; Shorter, J., eds.; Plenum: NY, 1978.
14. Hansch, C.; Leo, A.; Taft, R. W.; *Chem. Rev.* **1991**, 91, 165.
15. Smith, M. B.; March, J.; *Advanced Organic Chemistry. Reactions, Mechanisms and Structure*. 5<sup>th</sup> ed., Wiley: USA, 2001, p 363-380.
16. Pearson, A.J.; *Iron Compounds in Organic Chemistry*, Academic Press: CA, 1994, p. 199-270.
17. Reichenbach, G.; Cardaci, G.; Aloisi, G. G.; *J. Organomet. Chem.* **1977**, 134, 47.
18. Altomare, A.; Cascarano, G.; Giacovazzo, C.; Guagliardi, A.; Burla, M. C.; Polidori, G.; Canalli, M.; *J. Appl. Cryst.* **1997**, 27, 435.
19. Sheldrick, G. M.; *SHELXL-97 Program for Refinement of Crystal Structures*, Univ. of Goettingen, Germany, 1997.

Received: October 13,

Published on the web: March 15,